LOCAL EXTREMA IN RANDOM TREES

LANE CLARK

Received 23 March 2004 and in revised form 8 November 2005

The number of local maxima (resp., local minima) in a tree $T \in \mathcal{T}_n$ rooted at $r \in [n]$ is denoted by $M_r(T)$ (resp., by $m_r(T)$). We find exact formulas as rational functions of n for the expectation and variance of $M_1(T)$ and $m_n(T)$ when $T \in \mathcal{T}_n$ is chosen randomly according to a uniform distribution. As a consequence, a.a.s. $M_1(T)$ and $m_n(T)$ belong to a relatively small interval when $T \in \mathcal{T}_n$.

1. Introduction

The extension of permutation statistics to labelled trees is the subject of a number of articles. Generating functions for the number of labelled trees of several types according to the number of ascents and descents are given in [4]. A functional equation satisfied by the generating function for the number of labelled trees according to the number of descents and leaves is given in [5]. Central and local limit theorems for the number of ascents or of descents in uniformly random labelled trees are given in [1]. A functional equation satisfied by the generating function for the number of labelled trees according to the number of inversions is given in [7]. Related results are contained in [6]. A formula for the expected number of inversions of a uniformly random labelled tree is given in [9]. Formulas for the expectation and variance of the number of inversions of a uniformly random labelled tree are given in [2].

Local extrema (in the literature as local maxima and local minima; peaks and troughs; collectively turning points; related to phases) in permutations have a long history; see [10] and references there in. The examination of local maxima (equivalently, local minima) in permutations is more recent. A recurrence relation and a generating function for the number of permutations according to the number of local maxima are given in [10]. A central limit theorem for the number of local maxima in a uniformly random permutation also is given in [10]. In this note, we extend local extrema in permutations to labelled trees and examine local maxima (equivalently, local minima) in uniformly random labelled trees.

For $n \geq 2$, let $\mathcal{T}_n$ denote the set of trees with vertex set $[n] := \{1, \dots, n\}$. When $T_1, T_2 \in \mathcal{T}_n$, $T_1 = T_2$ if and only if T_1 and T_2 have the same edge set. Let $T \in \mathcal{T}_n$, $r \in [n]$, and

Table 1.1

	$T_1(n, k) = t_n(n, k)$	k					
		0	1	2	3	4	...
n	2	1	0	0	0	0	...
	3	2	1	0	0	0	...
	4	6	9	1	0	0	...
	5	24	73	27	1	0	...

distinct $i, j, k \in [n]$. We say T rooted at r has a *local maximum at path ijk* if and only if $j > i, k$ and the path in T from r to k contains the path ijk . Similarly, T rooted at r has a *local minimum at path ijk* if and only if $j < i, k$ and the path in T from r to k contains the path ijk . Let $M_r(T) = M_{r,n}(T)$ (resp., $m_r(T) = m_{r,n}(T)$) denote the number of local maxima (resp., local minima) of $T \in \mathcal{T}_n$ rooted at r . Then $M_r(T), m_r(T) \in \{0, \dots, n-2\}$. Let $T_r(n, k)$ (resp., $t_r(n, k)$) denote the number of trees in $\mathcal{T}_n$ rooted at r with precisely k local maxima (resp., k local minima). Then $T_r(n, k) = t_r(n, k) = 0$ for $k \notin \{0, \dots, n-2\}$ and $\sum_{k=0}^{n-2} T_r(n, k) = \sum_{k=0}^{n-2} t_r(n, k) = n^{n-2}$. As with the other statistics extended to labelled trees, roots $r = 1, n$ are appropriate. The values of $T_1(n, k) = t_n(n, k)$ (see Lemma 2.1) are given in Table 1.1 for $2 \leq n \leq 5$ and $0 \leq k \leq n-2$.

We work in the probability space Ω_n consisting of all trees in $\mathcal{T}_n$, where each tree is chosen randomly according to a uniform distribution. Hence, $\Pr(T) = 1/n^{n-2}$ for $T \in \mathcal{T}_n$. A property Q of trees in $\{\mathcal{T}_n\}$ holds *asymptotically almost surely* (a.a.s.) on $\{\Omega_n\}$ if and only if $\lim_{n \rightarrow \infty} \Pr(T \in \mathcal{T}_n : T \text{ has property } Q) \rightarrow 1$ as $n \rightarrow \infty$.

The parameters M_r and m_r are then random variables on Ω_n whose exact expectations $E(M_r)$, $E(m_r)$ and variances $\sigma^2(M_r)$, $\sigma^2(m_r)$ we find as rational functions of n ($r = 1, n$). From Theorem 2.5,

$$\begin{aligned} E(M_1) &= E(m_n) = \frac{2n^3 - 3n^2 - 5n + 6}{6n^2}, \\ \sigma^2(M_1) &= \sigma^2(m_n) = \frac{7n^5 - 20n^4 + 75n^3 - 40n^2 - 322n + 300}{60n^4}. \end{aligned} \quad (1.1)$$

As a consequence, a.a.s. on $\{\Omega_n\}$,

$$E(M_1) - \omega(n)\sigma(M_1) < M_1, m_n < E(M_1) + \omega(n)\sigma(M_1), \quad (1.2)$$

where $\omega(n) \rightarrow \infty$ arbitrarily slowly as $n \rightarrow \infty$. (See Corollary 2.6 for this and further results.)

We mention that should $(M_1 - E(M_1))/\sigma(M_1) \xrightarrow{d} N(0, 1)$, we could only conclude the above inequality for M_1, m_n a.a.s. on $\{\Omega_n\}$. Of course, asymptotic normality of M_1, m_n gives more information about the distribution of M_1, m_n than their a.a.s. properties.

Let $\mathbb{N}$ denote the nonnegative integers and let $\mathbb{R}$ denote the real numbers. The expectation of a random variable X is denoted by $E(X)$ and its variance by $\text{Var}(X)$. We refer the reader to Moon [8] for trees and Durrett [3] for probability.

2. Results

We first show that local maxima in trees rooted at r and local minima in trees rooted at $n+1-r$ are equidistributed.

LEMMA 2.1. For $r \in [n]$,

$$T_r(n, k) = t_{n+1-r}(n, k) \quad (0 \leq k \leq n-2). \quad (2.1)$$

Proof. The bijection $i \mapsto n+1-i$ ($i \in [n]$) induces a bijection $T \in \mathcal{T}_n \mapsto T' \in \mathcal{T}_n$, where $T \simeq T'$. Then $r, \dots, i, j, k$ (with $j > i, k$) is a path in T if and only if $n+1-r, \dots, n+1-i, n+1-j, n+1-k$ (with $n+1-j < n+1-i, n+1-k$) is a path in T' . Hence, $M_r(T) = m_{n+1-r}(T')$. Consequently, $T_r(n, k) = t_{n+1-r}(n, k)$ for $0 \leq k \leq n-2$. $\square$

In view of Lemma 2.1, we consider only M_1 .

Let $(x)_0 = x^0 = 1$ ($x \in \mathbb{R}$) and $(x)_k = (x) \cdot \dots \cdot (x - k + 1)$ ($k \geq 1, x \in \mathbb{R}$). For $n \in \mathbb{N}$ and $a, x \in \mathbb{R}$, let

$$\begin{aligned} E_n(x) &= \sum_{k=0}^n \frac{x^k}{k!}, & P_n(x) &= \sum_{k=0}^n (n)_k x^k, & Q_{n,a}(x) &= \sum_{k=0}^n (n)_k (k+a) x^k, \\ R_n(x) &= \sum_{k=0}^n (n)_k (k+1)(k+7) x^k. \end{aligned} \quad (2.2)$$

We require the following technical result which allows us to calculate the exact expectation and variance of M_1 as rational functions of n .

LEMMA 2.2. For $n, m-1 \in \mathbb{N}$,

$$P_n\left(\frac{1}{m}\right) = \frac{n!}{m^n} E_n(m), \quad (2.3)$$

for $n-1, m-1 \in \mathbb{N}$, and $a \in \mathbb{R}$,

$$Q_{n,a}\left(\frac{1}{m}\right) = \frac{n!}{m^n} (n+a) E_n(m) - \frac{n!}{m^{n-1}} E_{n-1}(m), \quad (2.4)$$

and for $n-2, m-1 \in \mathbb{N}$,

$$R_n\left(\frac{1}{m}\right) = \frac{n!}{m^n} (n^2 + 8n + 7) E_n(m) - \frac{n!}{m^{n-1}} (2n+7) E_{n-1}(m) + \frac{n!}{m^{n-2}} E_{n-2}(m). \quad (2.5)$$

Proof. (All derivatives are with respect to real x). First,

$$\frac{x^n P_n(x^{-1})}{n!} = \sum_{k=0}^n \frac{x^{n-k}}{(n-k)!} = E_n(x) \quad (2.6)$$

so that

$$P_n(x) = n! x^n E_n(x^{-1}), \quad (2.7)$$

and hence

$$P_n\left(\frac{1}{m}\right) = \frac{n!}{m^n} E_n(m). \quad (2.8)$$

Next, (2.7) gives

$$\begin{aligned} Q_{n,a}(x) &= xP'_n(x) + aP_n(x) \\ &= n!x^n(n+a)E_n(x^{-1}) - n!x^{n-1}E_{n-1}(x^{-1}), \end{aligned} \quad (2.9)$$

and hence

$$Q_{n,a}\left(\frac{1}{m}\right) = \frac{n!}{m^n}(n+a)E_n(m) - \frac{n!}{m^{n-1}}E_{n-1}(m). \quad (2.10)$$

Finally, (2.7) gives

$$\begin{aligned} R_n(x) &= x^2P''_n(x) + 9xP'_n(x) + 7P_n(x) \\ &= n!x^n(n^2 + 8n + 7)E_n(x^{-1}) - n!x^{n-1}(2n + 7)E_{n-1}(x^{-1}) \\ &\quad + n!x^{n-2}E_{n-2}(x^{-1}), \end{aligned} \quad (2.11)$$

and hence

$$R_n\left(\frac{1}{m}\right) = \frac{n!}{m^n}(n^2 + 8n + 7)E_n(m) - \frac{n!}{m^{n-1}}(2n + 7)E_{n-1}(m) + \frac{n!}{m^{n-2}}E_{n-2}(m). \quad (2.12)$$

□

COROLLARY 2.3. For $j, n-1 \in \mathbb{N}$ with $0 \leq j \leq n$,

$$Q_{n-j,j}\left(\frac{1}{n}\right) = n. \quad (2.13)$$

Proof. For $0 \leq j \leq n-1$, our result follows from Lemma 2.2. For $j = n$, our result follows from the definition of $Q_{n-j,j}(x)$. □

We require the following result of Moon [8].

THEOREM 2.4 (Moon [8]). Let F be a forest with vertex set $[n]$ having ω components of orders $p_1, \dots, p_\omega$. Then the number of distinct trees in $\mathcal{T}_n$ containing F is $pn^{\omega-2}$, where $p = p_1 \cdots p_\omega$.

We now give our main result. Here $M_1 = M_{1,n}$.

THEOREM 2.5. For Ω_n ($n \geq 2$),

$$\begin{aligned} E(M_1) &= \frac{2n^3 - 3n^2 - 5n + 6}{6n^2}, \\ E(M_1^2) &= \frac{20n^6 - 39n^5 - 115n^4 + 495n^3 - 175n^2 - 1266n + 1080}{180n^4}, \end{aligned} \quad (2.14)$$

hence,

$$\sigma^2(M_1) = \text{Var}(M_1) = \frac{7n^5 - 20n^4 + 75n^3 - 40n^2 - 322n + 300}{60n^4}. \quad (2.15)$$

Proof. The theorem can be seen to be true for $2 \leq n \leq 5$ using Table 1.1. Assume $n \geq 6$. Let $I_{n,1} = \{(i, j, k) : 1 \leq k < i < j \leq n\}$, $I_{n,2} = \{(i, j, k) : 1 \leq i < k < j \leq n\}$, and $I_n = I_{n,1} \cup I_{n,2}$. For $(i, j, k) \in I_n$ and $T \in \mathcal{T}_n$, let

$$X_{(i,j,k)}(T) = \begin{cases} 1, & \text{if } ijk \text{ is a local maximum in } T \text{ rooted at } 1; \\ 0, & \text{otherwise;} \end{cases} \quad (2.16)$$

hence,

$$M_1 = \sum_{(i,j,k) \in I_n} X_{(i,j,k)}. \quad (2.17)$$

We remind the reader that ij , jk are always edges in a tree by using thick lines in our diagrams.

Expectation of M_1 . We consider the following two cases according to the path S of T from 1 through ijk . Only $E(X_{(i,j,k)}) \neq 0$ need to be considered.

Case 1 ($i \neq 1$). Here

There are $(a+4)n^{n-a-5}$ trees in $\mathcal{T}_n$ containing a specific tree S by Theorem 2.4; there are $(n-4)_a$ specific trees containing a vertices between 1, i ; and there are $2^{\binom{n-1}{3}}$ choices for (i, j, k) . Hence,

$$\sum_{\substack{(i,j,k) \in I_n \\ 1 \neq i}} E(X_{(i,j,k)}) = \frac{(n-1)_3}{3n^3} \sum_{a=0}^{n-4} (n-4)_a \frac{a+4}{n^a} = \frac{(n-1)_3}{3n^2} \quad (2.18)$$

by Lemma 2.2.

Case 2 ($i = 1$). Here

There are $3n^{n-4}$ trees in $\mathcal{T}_n$ containing a specific tree S by Theorem 2.4; and there are $\binom{n-1}{2}$ choices for (j, k) . Hence,

$$\sum_{(1,j,k) \in I_n} E(X_{(i,j,k)}) = \frac{3(n-1)_2}{2n^2}. \quad (2.19)$$

From (2.18), (2.19),

$$E(M_1) = \frac{(n-1)_3}{3n^2} + \frac{3(n-1)_2}{2n^2} = \frac{2n^3 - 3n^2 - 5n + 6}{6n^2}. \quad (2.20)$$

Variance of M_1 . Here

$$M_1^2 = \left(\sum_{(i,j,k) \in I_n} X_{(i,j,k)} \right)^2 = M_1 + \sum_{((i_1,j_1,k_1),(i_2,j_2,k_2)) \in I_n^*} X_{(i_1,j_1,k_1)} X_{(i_2,j_2,k_2)}, \quad (2.21)$$

where $I_n^* = \{((i_1, j_1, k_1), (i_2, j_2, k_2)) \in I_n \times I_n : (i_1, j_1, k_1) \neq (i_2, j_2, k_2)\}$.

First, we describe how we calculate $E(M_1^2) - E(M_1)$.

We first consider $3 \cdot 2 = 6$ cases according to $\#\{i_1, j_1, k_1, i_2, j_2, k_2\} = 6, 5$, or 4 , and, whether $1 \notin \{i_1, j_1, k_1, i_2, j_2, k_2\}$ or $1 \in \{i_1, j_1, k_1, i_2, j_2, k_2\}$. In each of these six cases, we further partition as described below.

For $((i_1, j_1, k_1), (i_2, j_2, k_2)) \in I_n^*$, we consider the possible subtrees $S = S_{((i_1,j_1,k_1),(i_2,j_2,k_2))}$ of $[n]$ determined by the path from 1 to the second coordinate $i_2 j_2 k_2$ relative to the path from 1 to the first coordinate $i_1 j_1 k_1$. The possible subtrees $S' = S'_{((i_2,j_2,k_2),(i_1,j_1,k_1))}$ are included above by definition. Only $E(X_{(i_1,j_1,k_1)} X_{(i_2,j_2,k_2)}) \neq 0$ need to be considered. This gives nine types of subtrees of $[n]$ total among these six cases.

For the symmetric types $1, 3, 5, 7$, and 9 , S “looks like” S' . We count the number t_S of trees $T \in \mathcal{T}_n$ containing $S = S_{((i_1,j_1,k_1),(i_2,j_2,k_2))}$ and the number i_S of such $((i_1, j_1, k_1), (i_2, j_2, k_2))$. The product $i_S t_S$ counts each tree $T \in \mathcal{T}_n$ containing S twice; once for S and once for S' . For each such tree T , $X_{(i_1,j_1,k_1)}(T) X_{(i_2,j_2,k_2)}(T) = 1 = X_{(i_2,j_2,k_2)}(T) X_{(i_1,j_1,k_1)}(T)$.

For the asymmetric types $2, 4, 6$, and 8 , S “looks different” than S' . We introduce subtypes $S^x_{((i_1,j_1,k_1),(i_2,j_2,k_2))}$ and $S^y_{((i_1,j_1,k_1),(i_2,j_2,k_2))}$ so that $T \in \mathcal{T}_n$ contains $S^x = S^x_{((i_1,j_1,k_1),(i_2,j_2,k_2))}$ if and only if T contains $S^y = S^y_{((i_2,j_2,k_2),(i_1,j_1,k_1))}$; note the different orders. We count the number t_{S^z} of trees $T \in \mathcal{T}_n$ containing S^z and the number i_{S^z} of such $((i_1, j_1, k_1), (i_2, j_2, k_2))$ for $z = x, y$. The sum $i_{S^x} t_{S^x} + i_{S^y} t_{S^y}$ counts each tree $T \in \mathcal{T}_n$ containing S^x and S^y ; once for S^x and once for S^y . For each such tree T , $X_{(i_1,j_1,k_1)}(T) X_{(i_2,j_2,k_2)}(T) = 1 = X_{(i_2,j_2,k_2)}(T) X_{(i_1,j_1,k_1)}(T)$.

For each type, the above count(s) are divided by n^{n-2} then simplified using Lemma 2.2 and Corollary 2.3. Summing over the types of a particular case i ($1 \leq i \leq 6$) gives

$$\sum_{\substack{((i_1,j_1,k_1),(i_2,j_2,k_2)) \in I_n^* \\ \text{case } i}} E(X_{(i_1,j_1,k_1)} X_{(i_2,j_2,k_2)}). \quad (2.22)$$

The sum over all six cases is then $E(M_1^2) - E(M_1)$.

In what follows, $(n-1)_6 = E_{n-7}(n) = 0$ for $n = 6$ and $E_{n-9}(n) = 0$ for $n = 6, 7, 8$ as usual. All cases appear for $n \geq 9$.

Case 3 ($\#\{i_1, j_1, k_1, i_2, j_2, k_2\} = 6, 1 \notin \{i_1, j_1, k_1, i_2, j_2, k_2\}$).

Type 1. Here

There are $(a + b + 7)n^{n-a-b-8}$ trees in $\mathcal{T}_n$ containing a specific tree S^1 by Theorem 2.4; there are $(n - 7)_{a+b}$ specific trees containing a vertices between 1, i_1 and b vertices between x , i_2 ; there are $a + 1$ choices for x ; and there are $2\binom{n-1}{3} \cdot 2\binom{n-4}{3}$ such pairs $((i_1, j_1, k_1), (i_2, j_2, k_2))$. Observe that T contains $S^1_{((i_1, j_1, k_1), (i_2, j_2, k_2))}$ for a, b, x if and only if T contains $S^1_{((i_2, j_2, k_2), (i_1, j_1, k_1))}$ for a', b', x . Hence, (each such pair appears once)

$$\begin{aligned}
 \sum_{\substack{((i_1, j_1, k_1), (i_2, j_2, k_2)) \in I_n^* \\ T \text{ Type 1}}} E(X_{(i_1, j_1, k_1)} X_{(i_2, j_2, k_2)}) &= \frac{(n-1)_6}{9n^6} \sum_{\substack{(a,b) \in \mathbb{N}^2 \\ 0 \leq a+b \leq n-7}} (n-7)_{a+b} \frac{(a+1)(a+b+7)}{n^{a+b}} \\
 &= \frac{(n-1)_6}{9n^6} \sum_{a=0}^{n-7} (n-7)_a \frac{a+1}{n^a} \sum_{b=0}^{n-a-7} (n-a-7)_b \frac{a+b+7}{n^b} \\
 &= \frac{(n-1)_6}{9n^5} \sum_{a=0}^{n-7} (n-7)_a \frac{a+1}{n^a} \\
 &= \frac{(n-1)_6}{9n^5} \left\{ n - \frac{6(n-7)!}{n^{n-7}} E_{n-7}(n) \right\} \\
 &= \frac{(n-1)_6}{9n^4} - \frac{2(n-1)!}{3n^{n-2}} E_{n-7}(n)
 \end{aligned} \tag{2.23}$$

by Lemma 2.2, and Corollary 2.3.

Type 2. First subtypes $2_1, 2_2, 2_3$ are

In each of the subcases, we have replaced one of the $a + 1$ edges uv between $1, i_1$, with the path $ux = i_2v$, $ui_2x = j_2v$ or $ui_2j_2x = k_2v$, where the rest of the path $i_2j_2k_2$ is as indicated. In each of these three subcases, there are $(a + 7)n^{n-a-8}$ trees in $\mathcal{T}_n$ containing a specific tree $S^{2_1}, S^{2_2}, S^{2_3}$ by Theorem 2.4; there are $(n - 7)_a$ specific trees containing a other vertices between $1, i_1$; there are $a + 1$ choices for x , equivalently, uv ; and there are $2\binom{n-1}{3} \cdot 2\binom{n-4}{3}$ such pairs $((i_1, j_1, k_1), (i_2, j_2, k_2))$. Next, subtype 2_4 is

There are $(a + b + 7)n^{n-a-b-8}$ trees in $\mathcal{T}_n$ containing a specific tree S^{2_4} by Theorem 2.4; there are $(n - 7)_{a+b}$ specific trees containing a vertices between $1, i_1$ and b vertices between x, i_2 ; there are 3 choices for $x = i_1, j_1$, or k_1 ; and there are $2\binom{n-1}{3} \cdot 2\binom{n-4}{3}$ such pairs $((i_1, j_1, k_1), (i_2, j_2, k_2))$. Observe that T contains $S^{2_{1,2 \text{ or } 3}}_{((i_1, j_1, k_1), (i_2, j_2, k_2))}$ for a, x if and only if T contains $S^{2_4}_{((i_2, j_2, k_2), (i_1, j_1, k_1))}$ for a', b', x . Hence, (each such pair appears once)

$$\sum_{\substack{((i_1, j_1, k_1), (i_2, j_2, k_2)) \in I_n^* \\ T \text{ Type } 2}} E(X_{(i_1, j_1, k_1)} X_{(i_2, j_2, k_2)})$$

$$= \frac{(n-1)_6}{9n^6} \left\{ 3 \sum_{a=0}^{n-7} (n-7)_a \frac{(a+1)(a+7)}{n^a} + 3 \sum_{\substack{(a,b) \in \mathbb{N}^2 \\ 0 \leq a+b \leq n-7}} (n-7)_{a+b} \frac{a+b+7}{n^{a+b}} \right\}$$

elements j_1, j_2 , and there are $2 \cdot 2 = 4$ pairs with $j_1 > k_1 > j_2$ or $j_2 > k_2 > j_1$). Observe that T contains $S^3_{((i_1, j_1, k_1), (i_2, j_2, k_2))}$ if and only if T contains $S^3_{((i_2, j_2, k_2), (i_1, j_1, k_1))}$. Hence, (each such pair appears once)

$$\sum_{\substack{((i_1, j_1, k_1), (i_2, j_2, k_2)) \in I_n^* \\ T \text{ Type 3}}} E(X_{(i_1, j_1, k_1)} X_{(i_2, j_2, k_2)}) = \frac{2(n-1)_5}{15n^5} \sum_{a=0}^{n-6} (n-6)_a \frac{a+6}{n^a} = \frac{2(n-1)_5}{15n^4} \quad (2.26)$$

by Corollary 2.3.

Type 4. First subtypes $4_1, 4_2$ are

In either subcase, there are $(a+6)n^{n-a-7}$ trees in $\mathcal{T}_n$ containing a specific tree S^{4_1}, S^{4_2} by Theorem 2.4; there are $(n-6)_a$ specific trees containing a vertices between 1, i_1 ; there are $24 \binom{n-1}{5}$ such pairs $((i_1, j_1, k_1), (i_2, j_2, k_2))$ total (for 5 elements in $\{2, \dots, n\}$; there are $6+2=8$ pairs with $j_2 > i_2 = j_1$ or $j_2 > k_2 > i_2 = j_1$ for 4_1 ; there are $2 \cdot 6 = 12$ pairs with largest elements j_1, j_2 ; and there are $2 \cdot 2 = 4$ pairs with $j_1 > i_1 > j_2$ or $j_2 > k_2 > j_1$ for 4_2). Next subtypes $4_3, 4_4$ are

In either subcase, there are $(a + 6)n^{n-a-7}$ trees in $\mathcal{T}_n$ containing a specific tree S^4_3 , S^4_4 by Theorem 2.4; there are $(n - 6)_a$ specific trees containing a vertices between 1, i_2 ; and there are $24 \binom{n-1}{5}$ such pairs $((i_1, j_1, k_1), (i_2, j_2, k_2))$ total (for 5 elements in $\{2, \dots, n\}$, there are $6 + 2 = 8$ pairs with $j_1 > i_1 = j_2$ or $j_1 > k_1 > i_1 = j_2$ for 4_3 ; there are $2 \cdot 6 = 12$ pairs with largest elements j_1, j_2 , and there are $2 \cdot 2 = 4$ pairs with $j_2 > i_2 > j_1$ or $j_1 > k_1 > j_2$ for 4_4). Observe that T contains $S^4_i_{((i_1, j_1, k_1), (i_2, j_2, k_2))}$ if and only if T contains $S^{4_{i+2}}_{((i_2, j_2, k_2), (i_1, j_1, k_1))}$ for $i = 1, 2$. Hence, (each such pair appears once)

$$\sum_{\substack{((i_1, j_1, k_1), (i_2, j_2, k_2)) \in I_n^* \\ T \text{ Type 4}}} E(X_{(i_1, j_1, k_1)} X_{(i_2, j_2, k_2)}) = \frac{2(n-1)_5}{5n^5} \sum_{a=0}^{n-6} (n-6)_a \frac{a+6}{n^a} = \frac{2(n-1)_5}{5n^4} \quad (2.27)$$

by Corollary 2.3. (The number of subcases has been accounted for.) Summing (2.26), (2.27) gives the following equation:

$$\sum_{\substack{((i_1, j_1, k_1), (i_2, j_2, k_2)) \in I_n^* \\ \#\{i_1, j_1, k_1, i_2, j_2, k_2\} = 5, 1 \notin \{i_1, j_1, k_1, i_2, j_2, k_2\}}} E(X_{(i_1, j_1, k_1)} X_{(i_2, j_2, k_2)}) = \frac{2(n-1)_5}{15n^4} + \frac{2(n-1)_5}{5n^4} = \frac{8(n-1)_5}{15n^4}. \quad (2.28)$$

Case 5 ($\#\{i_1, j_1, k_1, i_2, j_2, k_2\} = 4, 1 \notin \{i_1, j_1, k_1, i_2, j_2, k_2\}$).

Type 5. Here

There are $(a + 5)n^{n-a-6}$ trees in $\mathcal{T}_n$ containing a specific tree S^5 by Theorem 2.4; there are $(n - 5)_a$ specific trees containing a vertices between 1, i_1 ; and there are $6 \binom{n-1}{4}$ such pairs

$((i_1, j_1, k_1), (i_2, j_2, k_2))$. Observe that T contains $S_{((i_1, j_1, k_1), (i_2, j_2, k_2))}^5$ if and only if T contains $S_{((i_2, j_2, k_2), (i_1, j_1, k_1))}^5$. Hence, (each such pair appears once)

$$\sum_{\substack{((i_1, j_1, k_1), (i_2, j_2, k_2)) \in I_n^* \\ \# \{i_1, j_1, k_1, i_2, j_2, k_2\} = 4, 1 \notin \{i_1, j_1, k_1, i_2, j_2, k_2\}}} E(X_{(i_1, j_1, k_1)} X_{(i_2, j_2, k_2)}) = \frac{(n-1)_4}{4n^4} \sum_{a=0}^{n-5} (n-5)_a \frac{a+5}{n^a} = \frac{(n-1)_4}{4n^3} \quad (2.29)$$

by Corollary 2.3.

Case 6 ($\# \{i_1, j_1, k_1, i_2, j_2, k_2\} = 6, 1 \in \{i_1, j_1, k_1, i_2, j_2, k_3\}$).

Type 6. First subtypes $6_1, 6_2, 6_3$ are

In each of these three subcases, there are $(a+6)n^{n-a-7}$ trees in $\mathcal{T}_n$ containing a specific tree $S^{6_1}, S^{6_2}, S^{6_3}$ by Theorem 2.4; there are $(n-6)_a$ specific trees containing a vertices between x, i_1 ; and there are $\binom{n-1}{2} \cdot 2 \binom{n-3}{3}$ such pairs $((i_1, j_1, k_1), (1, i_2, j_2))$. Next

subtype 6_4 is

There are $(a+6)n^{n-a-7}$ trees in $\mathcal{T}_n$ containing a specific tree S^{6_4} by Theorem 2.4; there are $(n-6)_a$ specific trees containing a vertices between x , i_2 ; there are 3 choices for $x = 1, j_1$, or k_1 ; and there are $\binom{n-1}{2} \cdot 2\binom{n-3}{3}$ such pairs $((1, j_1, k_1), (i_2, j_2, k_2))$. Observe that T contains $S_{((i_1,j_1,k_1),(1,j_2,k_2))}^{6_{1,2,3}}$ for a, x if and only if T contains $S_{((1,j_2,k_2),(i_1,j_1,k_1))}^{6_4}$ for a, x . Hence, (each such pair appears once)

$$\begin{aligned}
 & \sum_{\substack{((i_1,j_1,k_1),(i_2,j_2,k_2)) \in I_n^* \\ \#\{i_1,j_1,k_1,i_2,j_2,k_2\}=6, 1 \in \{i_1,j_1,k_1,i_2,j_2,k_2\}}} E(X_{(i_1,j_1,k_1)} X_{(i_2,j_2,k_2)}) \\
 &= \frac{(n-1)_5}{6n^5} \left\{ 3 \sum_{a=0}^{n-6} (n-6)_a \frac{a+6}{n^a} + 3 \sum_{a=0}^{n-6} (n-6)_a \frac{a+6}{n^a} \right\} \\
 &= \frac{(n-1)_5}{n^4}
 \end{aligned} \tag{2.30}$$

by Corollary 2.3. (The first 3 and second 3 above are the number of subcases, i.e., choices for x .)

Case 7 ($\#\{i_1, j_1, k_1, i_2, j_2, k_2\} = 5, 1 \in \{i_1, j_1, k_1, i_2, j_2, k_2\}$).

Type 7. Here

There are $5n^{n-6}$ trees in $\mathcal{T}_n$ containing a specific tree S^7 by Theorem 2.4; and there are $\binom{n-1}{2} \cdot \binom{n-3}{2}$ such pairs $((1, j_1, k_1), (1, j_2, k_2))$. Observe that T contains $S_{((1,j_1,k_1),(1,j_2,k_2))}^7$ if and only if T contains $S_{((1,j_2,k_2),(1,j_1,k_1))}^7$. Hence, (each such pair occurs once)

$$\sum_{\substack{((1,j_1,k_1),(1,j_2,k_2)) \in I_n^* \\ T \text{ Type 7}}} E(X_{(1,j_1,k_1)} X_{(1,j_2,k_2)}) = \frac{5(n-1)_4}{4n^4}. \tag{2.31}$$

Type 8. First subtypes $8_1, 8_2$ are

In either subcase, there are $5n^{n-6}$ trees in $\mathcal{T}_n$ containing a specific tree S^{8_1}, S^{8_2} by Theorem 2.4; and there are $8\binom{n-1}{4}$ such pairs $((1, j_1, k_1), (i_2, j_2, k_2))$ total (for 4 elements in $\{2, \dots, n\}$, there are $2 + 1 = 3$ pairs with $j_2 > j_1 = i_2$ or $j_2 > k_2 > i_2 = j_1$ for 8_1 ; and there are $2 + 2 + 1 = 5$ pairs with largest elements j_1, j_2 or $j_2 > k_2 > j_1$ for 8_2). Next subtypes $8_3, 8_4$ are

In either subcase, there are $5n^{n-6}$ trees in $\mathcal{T}_n$ containing a specific tree S^{8_3}, S^{8_4} by Theorem 2.4; and there are $8\binom{n-1}{4}$ such pairs $((i_1, j_1, k_1), (1, j_2, k_2))$ total (for 4 elements in $\{2, \dots, n\}$, there are $2 + 1 = 3$ pairs with $j_1 > j_2 = i_1$ or $j_1 > k_1 > i_1 = j_2$ for 8_3 ; and there are $2 + 2 + 1 = 5$ pairs with largest elements j_1, j_2 or $j_1 > k_1 > j_2$ for 8_4). Observe that T contains $S_{((1,j_1,k_1),(i_2,j_2,k_2))}^{8_i}$ if and only if T contains $S_{((i_2,j_2,k_2),(1,j_1,k_1))}^{8_{i+2}}$ for $i = 1, 2$. Hence, (each such pair occurs once)

$$\sum_{\substack{((i_1,j_1,k_1),(i_2,j_2,k_2)) \in I_n^* \\ T \text{ Type } 8}} E(X_{(i_1,j_1,k_1)} X_{(i_2,j_2,k_2)}) = \frac{5(n-1)_4}{3n^4} + \frac{5(n-1)_4}{3n^4} = \frac{10(n-1)_4}{3n^4}. \quad (2.32)$$

(The number of subcases has been accounted for.) Summing (2.31), (2.32) gives the following equation:

$$\sum_{\substack{((i_1, j_1, k_1), (i_2, j_2, k_2)) \in I_n^* \\ \#\{i_1, j_1, k_1, i_2, j_2, k_2\} = 5, 1 \in \{i_1, j_1, k_1, i_2, j_2, k_2\}}} E(X_{(i_1, j_1, k_1)} X_{(i_2, j_2, k_2)}) \quad (2.33)$$

$$= \frac{5(n-1)_4}{4n^4} + \frac{10(n-1)_4}{3n^4} = \frac{55(n-1)_4}{12n^4}.$$

Case 8 ($\#\{i_1, j_1, k_1, i_2, j_2, k_2\} = 4, 1 \in \{i_1, j_1, k_1, i_2, j_2, k_2\}$).

Type 9. Here

There are $4n^{n-5}$ trees in $\mathcal{T}_n$ containing a specific tree S^9 by Theorem 2.4; and there are $2\binom{n-1}{3}$ such pairs $((1, j_1, k_1), (1, j_2, k_2))$. Observe that T contains $S^9_{((1, j_1, k_1), (1, j_2, k_2))}$ if and only if T contains $S^9_{((1, j_2, k_2), (1, j_1, k_1))}$. Hence, (each such pair occurs once)

$$\sum_{\substack{((i_1, j_1, k_1), (i_2, j_2, k_2)) \in I_n^* \\ \#\{i_1, j_1, k_1, i_2, j_2, k_2\} = 4, 1 \in \{i_1, j_1, k_1, i_2, j_2, k_2\}}} E(X_{(i_1, j_1, k_1)} X_{(i_2, j_2, k_2)}) = \frac{4(n-1)_3}{3n^3}. \quad (2.34)$$

After all this preparation, we are now able to find the second moment and the variance of M_1 . From (2.21), summing (2.20), (2.25), (2.28)–(2.30), (2.33), (2.34) gives

$$\begin{aligned} E(M_1^2) &= \frac{2n^3 - 3n^2 - 5n + 6}{6n^2} + \frac{(n-1)_6}{9n^4} + \frac{8(n-1)_5}{15n^4} + \frac{(n-1)_4}{4n^3} \\ &\quad + \frac{(n-1)_5}{n^4} + \frac{55(n-1)_4}{12n^4} + \frac{4(n-1)_3}{3n^3} \\ &= \frac{20n^6 - 39n^5 - 115n^4 + 495n^3 - 175n^2 - 1266n + 1080}{180n^4}. \end{aligned} \quad (2.35)$$

Hence, (2.20), (2.35) give

$$\sigma^2(M_1) = \text{Var}(M_1) = \frac{7n^5 - 20n^4 + 75n^3 - 40n^2 - 322n + 300}{60n^4}. \quad (2.36)$$

□

As a consequence of Theorem 2.5, a.a.s. on $\{\Omega_n\}$, $M_1(T)$ and $m_n(T)$ belong to a relatively small interval for $T \in \mathcal{T}_n$. Again, $M_1 = M_{1,n}$.

COROLLARY 2.6. For $\{\Omega_n\}$,

$$\Pr(|M_1 - E(M_1)| < \omega(n)\sigma(M_1)) \longrightarrow 1 \quad \text{as } n \longrightarrow \infty, \quad (2.37)$$

where $\omega(n) \rightarrow \infty$ arbitrarily slowly as $n \rightarrow \infty$. Hence, a.a.s. on $\{\Omega_n\}$,

$$\frac{n}{3} - \omega(n)n^{0.5} < M_1 < \frac{n}{3} + \omega(n)n^{0.5}, \quad (2.38)$$

where $\omega(n) \rightarrow \infty$ arbitrarily slowly as $n \rightarrow \infty$.

Proof. By Chebyshev's inequality,

$$\Pr(|M_1 - E(M_1)| \geq \omega(n)\sigma(M_1)) \leq \frac{1}{\omega^2(n)} \rightarrow 0 \quad \text{as } n \rightarrow \infty, \quad (2.39)$$

provided that $\omega(n) \rightarrow \infty$ as $n \rightarrow \infty$. This implies our result. $\square$

Acknowledgment

I wish to thank both referees for comments and suggestions. The detailed report by one referee in particular led to this much improved version of the note.

References

- [1] L. H. Clark, *Ascents and descents in random trees*, preprint, 2004.
- [2] ———, *Inversions in random trees*, preprint, 2004.
- [3] R. Durrett, *Probability. Theory and Examples*, The Wadsworth & Brooks/Cole Statistics/Probability Series, Wadsworth & Brooks/Cole Advanced Books & Software, California, 1991.
- [4] Ö. Eğecioğlu and J. B. Remmel, *Bijections for Cayley trees, spanning trees, and their q -analogues*, J. Combin. Theory Ser. A **42** (1986), no. 1, 15–30.
- [5] I. M. Gessel, *Counting forests by descents and leaves*, Electron. J. Combin. **3** (1996), no. 2, Research Paper 8, 1–5.
- [6] I. M. Gessel, B. E. Sagan, and Y.-N. Yeh, *Enumeration of trees by inversions*, J. Graph Theory **19** (1995), no. 4, 435–459.
- [7] C. L. Mallows and J. Riordan, *The inversion enumerator for labeled trees*, Bull. Amer. Math. Soc. **74** (1968), 92–94.
- [8] J. W. Moon, *Counting Labelled Trees*, Canadian Mathematical Monographs, no. 1, Canadian Mathematical Congress, Quebec, 1970.
- [9] ———, *The expected number of inversions in a random tree*, Proc. Louisiana Conf. on Combinatorics, Graph Theory and Computing (Louisiana State Univ., Baton Rouge, La, 1970), Louisiana State University, Louisiana, 1970, pp. 375–382.
- [10] D. Warren and E. Seneta, *Peaks and Eulerian numbers in a random sequence*, J. Appl. Probab. **33** (1996), no. 1, 101–114.

Lane Clark: Department of Mathematics, College of Science, Southern Illinois University Carbondale, Carbondale, IL 62901-4408, USA

E-mail address: lclark@math.siu.edu