ON f -DERIVATIONS OF BCI-ALGEBRAS

JIANMING ZHAN AND YONG LIN LIU

Received 14 December 2004 and in revised form 16 May 2005

The notion of left-right (resp., right-left) f -derivation of a BCI-algebra is introduced, and some related properties are investigated. Using the idea of regular f -derivation, we give characterizations of a p -semisimple BCI-algebra.

1. Introduction and preliminaries

In the theory of rings and near-rings, the properties of derivations are an important topic to study, see [2, 3, 7, 10]. In [6], Jun and Xin applied the notions in rings and near-rings theory to BCI-algebras, and obtained some related properties. In this paper, the notion of left-right (resp., right-left) f -derivation of a BCI-algebra is introduced, and some related properties are investigated. Using the idea of regular f -derivation, we give characterizations of a p -semisimple BCI-algebra.

By a BCI-algebra we mean an algebra $(X; *, 0)$ of type $(2,0)$ satisfying the following conditions:

- (I) $((x * y) * (x * z)) * (z * y) = 0$;
- (II) $(x * (x * y)) * y = 0$;
- (III) $x * x = 0$;
- (IV) $x * y = 0$ and $y * x = 0$ imply that $x = y$;

for all $x, y, z \in X$.

In any BCI-algebra X , one can define a partial order “ $\leq$ ” by putting $x \leq y$ if and only if $x * y = 0$.

A subset S of a BCI-algebra X is called *subalgebra* of X if $x * y \in S$ for all $x, y \in S$. A subset I of a BCI-algebra X is called an *ideal* of X if it satisfies (i) $0 \in I$; (ii) $x * y \in I$ and $y \in I$ imply that $x \in I$ for all $x, y \in X$.

A mapping f of a BCI-algebra X into itself is called an *endomorphism* of X if $f(x * y) = f(x) * f(y)$ for all $x, y \in X$. Note that $f(0) = 0$. Especially, f is *monic* if for any $x, y \in X$, $f(x) = f(y)$ implies that $x = y$.

A BCI-algebra X has the following properties:

- (1) $x * 0 = x$;
- (2) $(x * y) * z = (x * z) * y$;

- (3) $x \leq y$ implies that $x * z \leq y * z$ and $z * y \leq z * x$;
- (4) $x * (x * (x * y)) = x * y$;
- (5) $(x * z) * (y * z) \leq x * y$;
- (6) $0 * (x * y) = (0 * x) * (0 * y)$;
- (7) $x * 0 = 0$ implies that $x = 0$.

For a BCI-algebra X , denote by X_+ (resp., $G(X)$) the BCK-part (resp., the BCI-G part) of X , that is, $X_+ = \{x \in X \mid 0 \leq x\}$ (resp., $G(X) = \{x \in X \mid 0 * x = x\}$). Note that $G(X) \cap X_+ = \{0\}$. If $X_+ = \{0\}$, then X is called a p -semisimple BCI-algebra.

In a p -semisimple BCI-algebra X , the following hold:

- (8) $(x * z) * (y * z) = x * y$;
- (9) $0 * (0 * x) = x$;
- (10) $x * (0 * y) = y * (0 * x)$;
- (11) $x * y = 0$ implies that $x = y$;
- (12) $x * a = x * b$ implies that $a = b$;
- (13) $a * x = b * x$ implies that $a = b$;
- (14) $a * (a * x) = x$.

Let X be a p -semisimple BCI-algebra. We define addition “+” as $x + y = x * (0 * y)$ for all $x, y \in X$. Then $(X, +)$ is an abelian group with identity 0 and $x - y = x * y$. Conversely, let $(X, +)$ be an abelian group with identity 0 and let $x * y = x - y$. Then X is a p -semisimple BCI-algebra and $x + y = x * (0 * y)$ for all $x, y \in X$ (see [5]).

For a BCI-algebra X , we denote $x \wedge y = y * (y * x)$, in particular, $0 * (0 * x) = a_x$, and $L_p(X) = \{a \in X \mid x * a = 0 \Rightarrow x = a \text{ for any } x \in X\}$. We call the elements of $L_p(X)$ the p -atoms of X . For any $a \in X$, let $V(a) = \{x \in X \mid a * x = 0\}$, which is called the branch of X with respect to a . It follows that $x * y \in V(a * b)$ whenever $x \in V(a)$ and $y \in V(b)$ for all $x, y \in X$ and $a, b \in L_p(X)$. Note that $L_p(X) = \{x \in X \mid a_x = x\}$, which is the p -semisimple part of X , and X is a p -semisimple BCI-algebra if and only if $L_p(X) = X$ (see [6]). Note also that $a_x \in L_p(X)$, that is, $0 * (0 * a_x) = a_x$, which implies that $a_x * y \in L_p(X)$ for all $y \in X$. It is clear that $G(X) \subseteq L_p(X)$, $x * (x * a) = a$, and $a * x \in L_p(X)$ for all $a \in L_p(X)$ and $x \in X$. For more details, refer to [1, 8, 11].

Definition 1.1 [9]. A BCI-algebra X is said to be *commutative* if $x = x \wedge y$ whenever $x \leq y$ for all $x, y \in X$.

Definition 1.2 [4]. A BCI-algebra X is said to be *branchwise commutative* if $x \wedge y = y \wedge x$ for all $x, y \in V(a)$ and all $a \in L_p(X)$.

LEMMA 1.3 [6]. A BCI-algebra X is commutative if and only if it is branchwise commutative.

Definition 1.4 [6]. Let X be a BCI-algebra. By a *left-right derivation* (briefly, (l, r) -derivation) of X , a self-map d of X satisfying the identity $d(x * y) = (d(x) * y) \wedge (x * d(y))$ for all $x, y \in X$ is meant. If d satisfies the identity $d(x * y) = (x * d(y)) \wedge (d(x) * y)$ for all $x, y \in X$, then it is said that d is a *right-left derivation* (briefly, (r, l) -derivation) of X . Moreover, if d is both an (r, l) - and an (l, r) -derivation, it is said that d is a *derivation*.

2. f -derivations

In what follows, let f be an endomorphism of X unless otherwise specified.

Definition 2.1. Let X be a BCI-algebra. By a *left-right f -derivation* (briefly, *(l,r) - f -derivation*) of X , a self-map d_f of X satisfying the identity $d_f(x * y) = (d_f(x) * f(y)) \wedge (f(x) * d_f(y))$ for all $x, y \in X$ is meant, where f is an endomorphism of X . If d_f satisfies the identity $d_f(x * y) = (f(x) * d_f(y)) \wedge (d_f(x) * f(y))$ for all $x, y \in X$, then it is said that d_f is a *right-left f -derivation* (briefly, *(r,l) - f -derivation*) of X . Moreover, if d_f is both an (r,l) - and an (l,r) - f -derivation, it is said that d_f is an *f -derivation*.

Example 2.2. Let $X = \{0, 1, 2, 3, 4, 5\}$ be a BCI-algebra with the following Cayley table:

$*$	0	1	2	3	4	5
0	0	0	2	2	2	2
1	1	0	2	2	2	2
2	2	2	0	0	0	0
3	3	2	1	0	0	0
4	4	2	1	1	0	1
5	5	2	1	1	1	0

Define a map $d_f : X \rightarrow X$ by

$$d_f(x) = \begin{cases} 2 & \text{if } x = 0, 1, \\ 0 & \text{otherwise,} \end{cases} \quad (2.1)$$

and define an endomorphism f of X by

$$f(x) = \begin{cases} 0 & \text{if } x = 0, 1, \\ 2 & \text{otherwise.} \end{cases} \quad (2.2)$$

Then it is easily checked that d_f is both derivation and f -derivation of X .

Example 2.3. Let X be a BCI-algebra as in Example 2.2. Define a map $d_f : X \rightarrow X$ by

$$d_f(x) = \begin{cases} 2 & \text{if } x = 0, 1, \\ 0 & \text{otherwise.} \end{cases} \quad (2.3)$$

Then it is easily checked that d_f is a derivation of X .

Define an endomorphism f of X by

$$f(x) = 0, \quad \forall x \in X. \quad (2.4)$$

Then d_f is not an f -derivation of X since

$$d_f(2 * 3) = d_f(0) = 2, \quad (2.5)$$

but

$$(d_f(2) * f(3)) \wedge (f(2) * d_f(3)) = (0 * 0) \wedge (0 * 0) = 0 \wedge 0 = 0, \quad (2.6)$$

and thus $d_f(2 * 3) \neq (d_f(2) * f(3)) \wedge (f(2) * d_f(3))$.

Remark 2.4. From Example 2.3, we know that there is a derivation of X which is not an f -derivation of X .

Example 2.5. Let $X = \{0, 1, 2, 3, 4, 5\}$ be a BCI-algebra with the following Cayley table:

$*$	0	1	2	3	4	5
0	0	0	3	2	3	2
1	1	0	5	4	3	2
2	2	2	0	3	0	3
3	3	3	2	0	2	0
4	4	2	1	5	0	3
5	5	3	4	1	2	0

Define a map $d_f : X \rightarrow X$ by

$$d_f(x) = \begin{cases} 0 & \text{if } x = 0, 1, \\ 2 & \text{if } x = 2, 4, \\ 3 & \text{if } x = 3, 5, \end{cases} \quad (2.7)$$

and define an endomorphism f of X by

$$f(x) = \begin{cases} 0 & \text{if } x = 0, 1, \\ 2 & \text{if } x = 2, 4, \\ 3 & \text{if } x = 3, 5. \end{cases} \quad (2.8)$$

Then it is easily checked that d_f is both derivation and f -derivation of X .

Example 2.6. Let X be a BCI-algebra as in Example 2.5. Define a map $d_f : X \rightarrow X$ by

$$d_f(x) = \begin{cases} 0 & \text{if } x = 0, 1, \\ 2 & \text{if } x = 2, 4, \\ 3 & \text{if } x = 3, 5. \end{cases} \quad (2.9)$$

Then it is easily checked that d_f is a derivation of X .

Define an endomorphism f of X by

$$f(0) = 0, \quad f(1) = 1, \quad f(2) = 3, \quad f(3) = 2, \quad f(4) = 5, \quad f(5) = 4. \quad (2.10)$$

Then d_f is not an f -derivation of X since

$$d_f(2 * 3) = d_f(3) = 3, \quad (2.11)$$

but

$$(d_f(2) * f(3)) \wedge (f(2) * d_f(3)) = (2 * 2) \wedge (3 * 3) = 0 \wedge 0 = 0, \quad (2.12)$$

and thus $d_f(2 * 3) \neq (d_f(2) * f(3)) \wedge (f(2) * d_f(3))$.

Example 2.7. Let X be a BCI-algebra as in Example 2.5. Define a map $d_f : X \rightarrow X$ by

$$d_f(0) = 0, \quad d_f(1) = 1, \quad d_f(2) = 3, \quad d_f(3) = 2, \quad d_f(4) = 5, \quad d_f(5) = 4. \quad (2.13)$$

Then d_f is not a derivation of X since

$$d_f(2 * 3) = d_f(3) = 2, \quad (2.14)$$

but

$$(d_f(2) * 3) \wedge (2 * d_f(3)) = (3 * 3) \wedge (2 * 2) = 0 \wedge 0 = 0, \quad (2.15)$$

and thus $d_f(2 * 3) \neq (d_f(2) * 3) \wedge (2 * d_f(3))$.

Define an endomorphism f of X by

$$f(0) = 0, \quad f(1) = 1, \quad f(2) = 3, \quad f(3) = 2, \quad f(4) = 5, \quad f(5) = 4. \quad (2.16)$$

Then it is easily checked that d_f is an f -derivation of X .

Remark 2.8. From Example 2.7, we know that there is an f -derivation of X which is not a derivation of X .

For convenience, we denote $f_x = 0 * (0 * f(x))$ for all $x \in X$. Note that $f_x \in L_P(X)$.

THEOREM 2.9. *Let d_f be a self-map of a BCI-algebra X defined by $d_f(x) = f_x$ for all $x \in X$. Then d_f is an (l, r) - f -derivation of X . Moreover, if X is commutative, then d_f is an (r, l) - f -derivation of X .*

Proof. Let $x, y \in X$.

Since

$$\begin{aligned} 0 * (0 * (f_x * f(y))) &= 0 * (0 * ((0 * (0 * f(x))) * f(y))) \\ &= 0 * (0 * ((0 * f(y)) * (0 * f(x)))) \\ &= 0 * (0 * (0 * f(y * x))) = 0 * f(y * x) \\ &= 0 * (f(y) * f(x)) = (0 * f(y)) * (0 * f(x)) \\ &= (0 * (0 * f(x))) * f(y) = f_x * f(y), \end{aligned} \quad (2.17)$$

we have $f_x * f(y) \in L_P(X)$, and thus

$$f_x * f(y) = (f(x) * f_y) * ((f(x) * f_y) * (f_x * f(y))). \quad (2.18)$$

It follows that

$$\begin{aligned} d_f(x * y) &= f_{x*y} = 0 * (0 * f(x * y)) = 0 * (0 * (f(x) * f(y))) \\ &= (0 * (0 * f(x))) * (0 * (0 * f(y))) = f_x * f_y \\ &= (0 * (0 * f_x)) * (0 * (0 * f(y))) = 0 * (0 * (f_x * f(y))) \\ &= f_x * f(y) = (f(x) * f_y) * ((f(x) * f_y) * (f_x * f(y))) \\ &= (f_x * f(y)) \wedge (f(x) * f_y) = (d_f(x) * f(y)) \wedge (f(x) * d_f(y)), \end{aligned} \quad (2.19)$$

and so d_f is an (l, r) - f -derivation of X . Now, assume that X is commutative. Using Lemma 1.3, it is sufficient to show that $d_f(x) * f(y)$ and $f(x) * d_f(y)$ belong to the same branch for all $x, y \in X$, we have

$$\begin{aligned} d_f(x) * f(y) &= f_x * f(y) = 0 * (0 * (f_x * f(y))) \\ &= (0 * (0 * f_x)) * (0 * (0 * f(y))) \\ &= f_x * f_y \in V(f_x * f_y), \end{aligned} \quad (2.20)$$

and so $f_x * f_y = (0 * (0 * f(x))) * (0 * (0 * f(y))) = 0 * (0 * (f(x) * f_y)) = 0 * (0 * (f(x) * d_f(y))) \leq f(x) * d_f(y)$, which implies that $f(x) * d_f(y) \in V(f_x * f_y)$. Hence, $d_f(x) * f(y)$ and $f(x) * d_f(y)$ belong to the same branch, and so

$$\begin{aligned} d_f(x * y) &= (d_f(x) * f(y)) \wedge (f(x) * d_f(y)) \\ &= (f(x) * d_f(y)) \wedge (d_f(x) * f(y)). \end{aligned} \quad (2.21)$$

This completes the proof. $\square$

PROPOSITION 2.10. *Let d_f be a self-map of a BCI-algebra X . Then the following hold.*

- (i) *If d_f is an (l, r) - f -derivation of X , then $d_f(x) = d_f(x) \wedge f(x)$ for all $x \in X$.*
- (ii) *If d_f is an (r, l) - f -derivation of X , then $d_f(x) = f(x) \wedge d_f(x)$ for all $x \in X$ if and only if $d_f(0) = 0$.*

Proof. (i) Let d_f be an (l, r) - f -derivation of X . Then,

$$\begin{aligned} d_f(x) &= d_f(x * 0) = (d_f(x) * f(0)) \wedge (f(x) * d_f(0)) \\ &= (d_f(x) * 0) \wedge (f(x) * d_f(0)) = d_f(x) \wedge (f(x) * d_f(0)) \\ &= (f(x) * d_f(0)) * ((f(x) * d_f(0)) * d_f(x)) \\ &= (f(x) * d_f(0)) * ((f(x) * d_f(x)) * d_f(0)) \\ &\leq f(x) * (f(x) * d_f(x)) = d_f(x) \wedge f(x). \end{aligned} \quad (2.22)$$

But $d_f(x) \wedge f(x) \leq d_f(x)$ is trivial and so (i) holds.

(ii) Let d_f be an (r, l) - f -derivation of X . If $d_f(x) = f(x) \wedge d_f(x)$ for all $x \in X$, then for $x = 0$, $d_f(0) = f(0) \wedge d_f(0) = 0 \wedge d_f(0) = d_f(0) * (d_f(0) * 0) = 0$.

Conversely, if $d_f(0) = 0$, then $d_f(x) = d_f(x * 0) = (f(x) * d_f(0)) \wedge (d_f(x) * f(0)) = (f(x) * 0) \wedge (d_f(x) * 0) = f(x) \wedge d_f(x)$, ending the proof. $\square$

PROPOSITION 2.11. *Let d_f be an (l, r) - f -derivation of a BCI-algebra X . Then,*

- (i) $d_f(0) \in L_p(X)$, that is, $d_f(0) = 0 * (0 * d_f(0))$;
- (ii) $d_f(a) = d_f(0) * (0 * f(a)) = d_f(0) + f(a)$ for all $a \in L_p(X)$;
- (iii) $d_f(a) \in L_p(X)$ for all $a \in L_p(X)$;
- (iv) $d_f(a + b) = d_f(a) + d_f(b) - d_f(0)$ for all $a, b \in L_p(X)$.

Proof. (i) The proof follows from Proposition 2.10(i).

(ii) Let $a \in L_p(X)$, then $a = 0 * (0 * a)$, and so $f(a) = 0 * (0 * f(a))$, that is, $f(a) \in L_p(X)$. Hence

$$\begin{aligned}
 d_f(a) &= d_f(0 * (0 * a)) \\
 &= (d_f(0) * f(0 * a)) \wedge (f(0) * d_f(0 * a)) \\
 &= (d_f(0) * f(0 * a)) \wedge (0 * d_f(0 * a)) \\
 &= (0 * d_f(0 * a)) * ((0 * d_f(0 * a)) * (d_f(0) * f(0 * a))) \\
 &= (0 * d_f(0 * a)) * ((0 * (d_f(0) * f(0 * a))) * d_f(0 * a)) \\
 &= 0 * (0 * (d_f(0) * (f(0) * f(a)))) \\
 &= 0 * (0 * (d_f(0) * (0 * f(a)))) \\
 &= d_f(0) * (0 * f(a)) = d_f(0) + f(a).
 \end{aligned} \tag{2.23}$$

(iii) The proof follows directly from (ii).

(iv) Let $a, b \in L_p(X)$. Note that $a + b \in L_p(X)$, so from (ii), we note that

$$\begin{aligned}
 d_f(a + b) &= d_f(0) + f(a + b) \\
 &= d_f(0) + f(a) + d_f(0) + f(b) - d_f(0) = d_f(a) + d_f(b) - d_f(0).
 \end{aligned} \tag{2.24}$$

□

PROPOSITION 2.12. *Let d_f be a (r, l) - f -derivation of a BCI-algebra X . Then,*

- (i) $d_f(a) \in G(X)$ for all $a \in G(X)$;
- (ii) $d_f(a) \in L_p(X)$ for all $a \in G(X)$;
- (iii) $d_f(a) = f(a) * d_f(0) = f(a) + d_f(0)$ for all $a \in L_p(X)$;
- (iv) $d_f(a + b) = d_f(a) + d_f(b) - d_f(0)$ for all $a, b \in L_p(X)$.

Proof. (i) For any $a \in G(X)$, we have $d_f(a) = d_f(0 * a) = (f(0) * d_f(a)) \wedge (d_f(0) * f(a)) = (d_f(0) * f(a)) * ((d_f(0) * f(a)) * (0 * d_f(a))) = 0 * d_f(a)$, and so $d_f(a) \in G(X)$.

(ii) For any $a \in L_p(X)$, we get

$$\begin{aligned}
 d_f(a) &= d_f(0 * (0 * a)) = (0 * d_f(0 * a)) \wedge (d_f(0) * f(0 * a)) \\
 &= (d_f(0) * f(0 * a)) * ((d_f(0) * f(0 * a)) * (0 * d_f(0 * a))) \\
 &= 0 * d_f(0 * a) \in L_p(X).
 \end{aligned} \tag{2.25}$$

(iii) For any $a \in L_p(X)$, we get

$$\begin{aligned}
 d_f(a) &= d_f(a * 0) = (f(a) * d_f(0)) \wedge (d_f(a) * f(0)) \\
 &= d_f(a) * (d_f(a) * (f(a) * d_f(0))) = f(a) * d_f(0) \\
 &= f(a) * (0 * d_f(0)) = f(a) + d_f(0).
 \end{aligned} \tag{2.26}$$

(iv) The proof follows from (iii). This completes the proof.

□

Using Proposition 2.12, we know there is an (l, r) - f -derivation which is not an (r, l) - f -derivation as shown in the following example.

Example 2.13. Let $\mathbb{Z}$ be the set of all integers and “ $-$ ” the minus operation on $\mathbb{Z}$. Then $(\mathbb{Z}, -, 0)$ is a BCI-algebra. Let $d_f : X \rightarrow X$ be defined by $d_f(x) = f(x) - 1$ for all $x \in \mathbb{Z}$. Then,

$$\begin{aligned}(d_f(x) - f(y)) \wedge (f(x) - d_f(y)) &= (f(x) - 1 - f(y)) \wedge (f(x) - (f(y) - 1)) \\ &= (f(x - y) - 1) \wedge (f(x - y) + 1) \\ &= (f(x - y) + 1) - 2 = f(x - y) - 1 \\ &= d_f(x - y).\end{aligned}\tag{2.27}$$

Hence, d_f is an (l, r) - f -derivation of X . But $d_f(0) = f(0) - 1 = -1 \neq 1 = f(0) - d_f(0) = 0 - d_f(0)$, that is, $d_f(0) \notin G(X)$. Therefore, d_f is not an (r, l) - f -derivation of X by Proposition 2.12(i).

3. Regular f -derivations

Definition 3.1. An f -derivation d_f of a BCI-algebra X is said to be *regular* if $d_f(0) = 0$.

Remark 3.2. We know that the f -derivations d_f in Examples 2.5 and 2.7 are regular.

PROPOSITION 3.3. *Let X be a commutative BCI-algebra and let d_f be a regular (r, l) - f -derivation of X . Then the following hold.*

- (i) Both $f(x)$ and $d_f(x)$ belong to the same branch for all $x \in X$.
- (ii) d_f is an (l, r) - f -derivation of X .

Proof. (i) Let $x \in X$. Then,

$$\begin{aligned}0 = d_f(0) &= d_f(a_x * x) \\ &= (f(a_x) * d_f(x)) \wedge (d(a_x) * f(x)) \\ &= (d(a_x) * f(x)) * ((d(a_x) * f(x)) * (f(a_x) * d_f(x))) \\ &= (d(a_x) * f(x)) * ((d(a_x) * f(x)) * (f_x * d_f(x))) \\ &= f_x * d_f(x) \quad \text{since } f_x * d_f(x) \in L_P(X),\end{aligned}\tag{3.1}$$

and so $f_x \leq d_f(x)$. This shows that $d_f(x) \in V(f_x)$. Clearly, $f(x) \in V(f_x)$.

(ii) By (i), we have $f(x) * d_f(y) \in V(f_x * f_y)$ and $d_f(x) * f(y) \in V(f_x * f_y)$. Thus $d_f(x * y) = (f(x) * d_f(y)) \wedge (d_f(x) * f(y)) = (d_f(x) * f(y)) \wedge (f(x) * d_f(y))$, which implies that d_f is an (l, r) - f -derivation of X . $\square$

Remark 3.4. The f -derivations d_f in Examples 2.5 and 2.7 are regular f -derivations but we know that the (l, r) - f -derivation d_f in Example 2.2 is not regular. In the following, we give some properties of regular f -derivations.

Definition 3.5. Let X be a BCI-algebra. Then define $\ker d_f = \{x \in X \mid d_f(x) = 0 \text{ for all } f\text{-derivations } d_f\}$.

PROPOSITION 3.6. *Let d_f be an f -derivation of a BCI-algebra X . Then the following hold:*

- (i) $d_f(x) \leq f(x)$ for all $x \in X$;
- (ii) $d_f(x) * f(y) \leq f(x) * d_f(y)$ for all $x, y \in X$;
- (iii) $d_f(x * y) = d_f(x) * f(y) \leq d_f(x) * d_f(y)$ for all $x, y \in X$;
- (iv) $\ker d_f$ is a subalgebra of X . Especially, if f is monic, then $\ker d_f \subseteq X_+$.

Proof. (i) The proof follows by Proposition 2.10(ii).

(ii) Since $d_f(x) \leq f(x)$ for all $x \in X$, then $d_f(x) * f(y) \leq f(x) * f(y) \leq f(x) * d_f(y)$.

(iii) For any $x, y \in X$, we have

$$\begin{aligned} d_f(x * y) &= (f(x) * d_f(y)) \wedge (d_f(x) * f(y)) \\ &= (d_f(x) * f(y)) * ((d_f(x) * f(y)) * (f(x) * d_f(y))) \\ &= (d_f(x) * f(y)) * 0 = d_f(x) * f(y) \leq d_f(x) * d_f(y), \end{aligned} \quad (3.2)$$

which proves (iii).

(iv) Let $x, y \in \ker d_f$, then $d_f(x) = 0 = d_f(y)$, and so $d_f(x * y) \leq d_f(x) * d_f(y) = 0 * 0 = 0$ by (iii), and thus $d_f(x * y) = 0$, that is, $x * y \in \ker d_f$. Hence, $\ker d_f$ is a subalgebra of X . Especially, if f is monic, and letting $x \in \ker d_f$, then $0 = d_f(x) \leq f(x)$ by (i), and so $f(x) \in X_+$, that is, $0 * f(x) = 0$, and thus $f(0 * x) = f(x)$, which implies that $0 * x = x$, and so $x \in X_+$, that is, $\ker d_f \subseteq X_+$. $\square$

THEOREM 3.7. *Let f be monic of a commutative BCI-algebra X . Then X is p -semisimple if and only if $\ker d_f = \{0\}$ for every regular f -derivation d_f of X .*

Proof. Assume that X is p -semisimple BCI-algebra and let d_f be a regular f -derivation of X . Then $X_+ = \{0\}$, and so $\ker d_f = \{0\}$ by using Proposition 3.6(iv). Conversely, let $\ker d_f = \{0\}$ for every regular f -derivation d_f of X . Define a self-map d_f^* of X by $d_f^*(x) = f_x$ for all $x \in X$. Using Theorem 2.9, d_f^* is an f -derivation of X . Clearly, $d_f^*(0) = f_0 = 0 * (0 * f(0)) = 0$, and so d_f^* is a regular f -derivation of X . It follows from the hypothesis that $\ker d_f^* = \{0\}$. In addition, $d_f^*(x) = f_x = 0 * (0 * f(x)) = f(0 * (0 * x)) = f(0) = 0$ for all $x \in X_+$, and thus $x \in \ker d_f^*$, which shows that $X_+ \subseteq \ker d_f^*$. Hence, by Proposition 3.6(iv), $X_+ = \ker d_f^* = \{0\}$. Therefore X is p -semisimple. $\square$

Definition 3.8. An ideal A of a BCI-algebra X is said to be an f -ideal if $f(A) \subseteq A$.

Definition 3.9. Let d_f be a self-map of a BCI-algebra X . An f -ideal A of X is said to be d_f -invariant if $d_f(A) \subseteq A$.

THEOREM 3.10. *Let d_f be a regular (r, l) - f -derivation of a BCI-algebra X , then every f -ideal A of X is d_f -invariant.*

Proof. By Proposition 2.10(ii), we have $d_f(x) = f(x) \wedge d_f(x) \leq f(x)$ for all $x \in X$. Let $y \in d_f(A)$. Then $y = d_f(x)$ for some $x \in A$. It follows that $y * f(x) = d_f(x) * f(x) = 0 \in A$. Since $x \in A$, then $f(x) \in f(A) \subseteq A$ as A is an f -ideal. It follows that $y \in A$ since A is an ideal of X . Hence $d_f(A) \subseteq A$, and thus A is d_f -invariant. $\square$

THEOREM 3.11. *Let d_f be an f -derivation of a BCI-algebra X . Then d_f is regular if and only if every f -ideal of X is d_f -invariant.*

Proof. Let d_f be a derivation of a BCI-algebra X and assume that every f -ideal of X is d_f -invariant. Then since the zero ideal $\{0\}$ is f -ideal and d_f -invariant, we have $d_f(\{0\}) \subseteq \{0\}$, which implies that $d_f(0) = 0$. Thus d_f is regular. Combining this and Theorem 3.10, we complete the proof. $\square$

Acknowledgments

This work was supported by the Education Committee of Hubei Province (2004Z002, D200529001). The authors would like to thank the Editor-in-Chief and referees for the valuable suggestions and corrections for the improvement of this paper.

References

- [1] M. Aslam and A. B. Thaheem, *A note on p -semisimple BCI-algebras*, Math. Japon. **36** (1991), no. 1, 39–45.
- [2] H. E. Bell and L.-C. Kappe, *Rings in which derivations satisfy certain algebraic conditions*, Acta Math. Hungar. **53** (1989), no. 3-4, 339–346.
- [3] H. E. Bell and G. Mason, *On derivations in near-rings*, Near-Rings and Near-Fields (Tübingen, 1985), North-Holland Math. Stud., vol. 137, North-Holland, Amsterdam, 1987, pp. 31–35.
- [4] M. A. Chaudhry, *Branchwise commutative BCI-algebras*, Math. Japon. **37** (1992), no. 1, 163–170.
- [5] M. Daoji, *BCI-algebras and abelian groups*, Math. Japon. **32** (1987), no. 5, 693–696.
- [6] Y. B. Jun and X. L. Xin, *On derivations of BCI-algebras*, Inform. Sci. **159** (2004), no. 3-4, 167–176.
- [7] K. Kaya, *Prime rings with α -derivations Hacettepe*, Bull. Mater. Sci. Eng. **16-17** (1987-1988), 63–71.
- [8] Y. L. Liu, J. Meng, and X. L. Xin, *Quotient rings induced via fuzzy ideals*, Korean J. Comput. Appl. Math. **8** (2001), no. 3, 631–643.
- [9] J. Meng and X. L. Xin, *Commutative BCI-algebras*, Math. Japon. **37** (1992), no. 3, 569–572.
- [10] E. C. Posner, *Derivations in prime rings*, Proc. Amer. Math. Soc. **8** (1957), 1093–1100.
- [11] L. Tiande and X. Changchang, *p -radical in BCI-algebras*, Math. Japon. **30** (1985), no. 4, 511–517.

Jianming Zhan: Department of Mathematics, Hubei Institute for Nationalities, Enshi 445000, Hubei Province, China

E-mail address: zhanjianming@hotmail.com

Yong Lin Liu: Department of Applied Mathematics, School of Science, Xidian University, Xi'an 710071, Shaanxi, China; Department of Mathematics, Nanping Teachers College, Nanping 353000, Fujian, China

E-mail address: ylliun@tom.com