NEW ERROR INEQUALITIES FOR THE LAGRANGE INTERPOLATING POLYNOMIAL

NENAD UJEVIĆ

Received 30 August 2005

A new representation of remainder of Lagrange interpolating polynomial is derived. Error inequalities of Ostrowski-Grüss type for the Lagrange interpolating polynomial are established. Some similar inequalities are also obtained.

1. Introduction

Many error inequalities in polynomial interpolation can be found in [1, 7]. These error bounds for interpolating polynomials are usually expressed by means of the norms $\|\cdot\|_p$, $1 \leq p \leq \infty$. Some new error inequalities (for corrected interpolating polynomials) are given in [10, 11]. The last mentioned inequalities are similar to error inequalities obtained in recent years in numerical integration and they are known in the literature as inequalities of Ostrowski (or Ostrowski-like, Ostrowski-Grüss) type. For example, in [9] we can find inequalities of Ostrowski-Grüss type for the well-known Simpson's quadrature rule,

$$\left| \int_{x_0}^{x_2} f(t) dt - \frac{h}{3} [f(x_0) + 4f(x_1) + f(x_2)] \right| \leq C_n (\Gamma_n - \gamma_n) h^{n+1}, \quad (1.1)$$

where $x_i = x_0 + ih$, for $h > 0$, $i = 1, 2$, γ_n, Γ_n are real numbers such that $\gamma_n \leq f^{(n)}(t) \leq \Gamma_n$, for all $t \in [x_0, x_2]$, and C_n are constants, $n \in \{1, 2, 3\}$.

The inequalities of Ostrowski type can be also found in [2, 3, 4, 5, 6, 12]. In some of the mentioned papers, we can find estimations for errors of quadrature formulas which are expressed by means of the differences $\Gamma_k - \gamma_k$, $S - \gamma_k$, $\Gamma_k - S$, where Γ_k, γ_k are real numbers such that $\gamma_k \leq f^{(k)}(t) \leq \Gamma_k$, $t \in [a, b]$ (k is a positive integer while $[a, b]$ is an interval of integration) and $S = [f^{(k-1)}(b) - f^{(k-1)}(a)]/(b - a)$. It is shown that the estimations expressed in such a way can be much better than the estimations expressed by means of the norms $\|f^{(k)}\|_p$, $1 \leq p \leq \infty$.

As we know there is a close relationship between interpolation polynomials and quadrature rules. Thus, it is a natural try to establish similar error inequalities in polynomial interpolation.

We first establish general error inequalities, expressed by means of $\|f^{(k)} - P_m\|$, where P_m is any polynomial of degree m and then we obtain inequalities of the above mentioned types. For that purpose, we derive a new representation of remainder of the interpolating polynomial. This is done in Section 2. In Section 3, we obtain the error inequalities of the above-mentioned types. In Section 4, we give some results for derivatives.

Finally, we emphasize that the usual error inequalities in polynomial interpolation (for the Lagrange interpolating polynomial $L_n(x)$) are given by means of the $(n+1)$ th derivative while in this paper we can find these error inequalities expressed by means of the k th derivative for $k = 1, 2, \dots, n$.

2. Representation of remainder

Let $D = \{a = x_0 < x_1 < \dots < x_n = b\}$ be a given subdivision of the interval $[a, b]$ and let $f : [a, b] \rightarrow \mathbb{R}$ be a given function. The Lagrange interpolation polynomial is given by

$$L_n(x) = \sum_{i=0}^n p_{ni}(x) f(x_i), \quad (2.1)$$

where

$$p_{ni}(x) = \frac{(x-x_0) \cdots (x-x_{i-1})(x-x_{i+1}) \cdots (x-x_n)}{(x_i-x_0) \cdots (x_i-x_{i-1})(x_i-x_{i+1}) \cdots (x_i-x_n)}, \quad (2.2)$$

for $i = 0, 1, \dots, n$. We have the Cauchy relations [7, pages 160-161],

$$\sum_{i=0}^n p_{ni}(x) = 1, \quad (2.3)$$

$$\sum_{i=0}^n p_{ni}(x)(x-x_i)^j = 0, \quad j = 1, 2, \dots, n. \quad (2.4)$$

Let $\bar{D} = \{x_0 = a < x_1 < \dots < x_n = b\}$ be a given uniform subdivision of the interval $[a, b]$, that is, $x_i = x_0 + ih$, $h = (b-a)/n$, $i = 0, 1, 2, \dots, n$. Then the Lagrange interpolating polynomial is given by

$$L_n(x) = L_n(x_0 + th) = (-1)^n \frac{t(t-1) \cdots (t-n)}{n!} \sum_{i=0}^n (-1)^i \binom{n}{i} \frac{f(x_i)}{t-i}, \quad (2.5)$$

where $t \notin \{0, 1, 2, \dots, n\}$, $0 < t < n$.

LEMMA 2.1. Let $P_m(t)$ be an arbitrary polynomial of degree $\leq m$ and let $p_{ni}(x)$ be defined by (2.2). Then

$$\sum_{i=0}^n p_{ni}(x) \int_{x_i}^x P_m(t)(t-x_i)^k dt = 0, \quad (2.6)$$

for $0 \leq k+m \leq n-1$ and $x \in [a, b]$.

Proof. Let x be a given real number. Then we have

$$P_m(t) = \sum_{j=0}^m c_j (x-t)^j, \quad (2.7)$$

for some coefficients $c_j = c_j(x)$, $j = 0, 1, 2, \dots, m$. (This is a consequence of the Taylor formula.) Thus,

$$\sum_{i=0}^n p_{ni}(x) \int_{x_i}^x P_m(t) (t-x_i)^k dt = \sum_{j=0}^m c_j \sum_{i=0}^n p_{ni}(x) \int_{x_i}^x (x-t)^j (t-x_i)^k dt. \quad (2.8)$$

Let $\beta(\cdot, \cdot)$ and $\Gamma(\cdot)$ denote the beta and gamma functions, respectively. We now calculate

$$\begin{aligned} \int_{x_i}^x (x-t)^j (t-x_i)^k dt &= \int_0^{x-x_i} (x-x_i-u)^j u^k du \\ &= (x-x_i)^j \int_0^{x-x_i} \left(1 - \frac{u}{x-x_i}\right)^j u^k du \\ &= (x-x_i)^{j+k+1} \int_0^1 (1-v)^j v^k dv \\ &= \beta(j+1, k+1) (x-x_i)^{j+k+1} \\ &= \frac{\Gamma(k+1)\Gamma(j+1)}{\Gamma(k+j+2)} (x-x_i)^{j+k+1} \\ &= \frac{k!j!}{(k+j+1)!} (x-x_i)^{j+k+1}. \end{aligned} \quad (2.9)$$

From (2.8) and (2.9) it follows that

$$\sum_{i=0}^n p_{ni}(x) \int_{x_i}^x P_m(t) (t-x_i)^k dt = \sum_{j=0}^m c_j \frac{k!j!}{(k+j+1)!} \sum_{i=0}^n p_{ni}(x) (x-x_i)^{j+k+1}. \quad (2.10)$$

From (2.10) and (2.4) we conclude that (2.6) holds. $\square$

THEOREM 2.2. Let $f \in C^{n+1}(a, b)$ and let the assumptions of Lemma 2.1 hold. Then

$$f(x) = L_n(x) + R_{k,m}(x), \quad (2.11)$$

where $L_n(x)$ is given by (2.1) and

$$R_{k,m}(x) = \frac{(-1)^k}{k!} \sum_{i=0}^n p_{ni}(x) \int_{x_i}^x [f^{(k+1)}(t) - P_m(t)] (t-x_i)^k dt. \quad (2.12)$$

Proof. We have

$$R_{k,m}(x) = \frac{(-1)^k}{k!} \sum_{i=0}^n p_{ni}(x) \int_{x_i}^x f^{(k+1)}(t) (t-x_i)^k dt - \frac{(-1)^k}{k!} \sum_{i=0}^n p_{ni}(x) \int_{x_i}^x P_m(t) (t-x_i)^k dt. \quad (2.13)$$

From (2.13) and (2.6) it follows that

$$R_{k,m}(x) = R_k(x) = \frac{(-1)^k}{k!} \sum_{i=0}^n p_{ni}(x) \int_{x_i}^x f^{(k+1)}(t)(t-x_i)^k dt. \quad (2.14)$$

For $k = 0$ we have

$$\begin{aligned} R_0(x) &= \sum_{i=0}^n p_{ni}(x) \int_{x_i}^x f'(t) dt \\ &= \sum_{i=0}^n p_{ni}(x) [f(x) - f(x_i)] = f(x) - L_n(x), \end{aligned} \quad (2.15)$$

since (2.3) holds.

We now suppose that $k \geq 1$. Integrating by parts, we obtain

$$\frac{(-1)^k}{k!} \int_{x_i}^x f^{(k+1)}(t)(t-x_i)^k dt = \frac{(-1)^k}{k!} f^{(k)}(x)(x-x_i)^k + \frac{(-1)^{k-1}}{(k-1)!} \int_{x_i}^x f^{(k)}(t)(t-x_i)^{k-1} dt. \quad (2.16)$$

In a similar way we get

$$\begin{aligned} &\frac{(-1)^{k-1}}{(k-1)!} \int_{x_i}^x f^{(k)}(t)(t-x_i)^{k-1} dt \\ &= \frac{(-1)^{k-1}}{(k-1)!} f^{(k-1)}(x)(x-x_i)^{k-1} + \frac{(-1)^{k-2}}{(k-2)!} \int_{x_i}^x f^{(k-1)}(t)(t-x_i)^{k-2} dt. \end{aligned} \quad (2.17)$$

Continuing in this way, we get

$$\begin{aligned} \frac{(-1)^k}{k!} \int_{x_i}^x f^{(k+1)}(t)(t-x_i)^k dt &= \sum_{j=1}^k \frac{(-1)^j}{j!} f^{(j)}(x)(x-x_i)^j + \int_{x_i}^x f'(t) dt \\ &= f(x) - f(x_i) + \sum_{j=1}^k \frac{(-1)^j}{j!} f^{(j)}(x)(x-x_i)^j. \end{aligned} \quad (2.18)$$

From (2.14) and (2.18) it follows that

$$\begin{aligned} R_k(x) &= \sum_{i=0}^n p_{ni}(x) \left[f(x) - f(x_i) + \sum_{j=1}^k \frac{(-1)^j}{j!} f^{(j)}(x)(x-x_i)^j \right] \\ &= f(x) - L_n(x) + \sum_{j=1}^k \frac{(-1)^j}{j!} f^{(j)}(x) \sum_{i=0}^n p_{ni}(x)(x-x_i)^j \\ &= f(x) - L_n(x), \quad k = 1, 2, \dots, n, \end{aligned} \quad (2.19)$$

since (2.3) and (2.4) hold. From (2.14), (2.15), and (2.19) we see that (2.11) holds. $\square$

3. Error inequalities

We now introduce the notations

$$\omega_n(x) = (x - x_0)(x - x_1) \cdots (x - x_n), \quad (3.1)$$

$$C_k(x) = \sum_{i=0}^n \frac{|x - x_i|^k}{|x_i - x_0| \cdots |x_i - x_{i-1}| |x_i - x_{i+1}| \cdots |x_i - x_n|}, \quad (3.2)$$

$$B_k(x) = \sum_{i=0}^n \frac{(S_{ki} - \gamma_{k+1}) |x - x_i|^k}{|x_i - x_0| \cdots |x_i - x_{i-1}| |x_i - x_{i+1}| \cdots |x_i - x_n|}, \quad (3.3)$$

$$D_k(x) = \sum_{i=0}^n \frac{(\Gamma_{k+1} - S_{ki}) |x - x_i|^k}{|x_i - x_0| \cdots |x_i - x_{i-1}| |x_i - x_{i+1}| \cdots |x_i - x_n|}, \quad (3.4)$$

where $S_{ki} = [f^{(k)}(x) - f^{(k)}(x_i)]/(x - x_i)$, $i = 0, 1, \dots, n$, and γ_{k+1} , Γ_{k+1} are real numbers such that $\gamma_{k+1} \leq f^{(k+1)}(t) \leq \Gamma_{k+1}$, $t \in [a, b]$, $k = 0, 1, \dots, n - 1$.

Let $g \in C(a, b)$. As we know among all algebraic polynomials of degree $\leq m$ there exists the only polynomial $P_m^*(t)$ having the property that

$$\|g - P_m^*\|_\infty \leq \|g - P_m\|_\infty, \quad (3.5)$$

where $P_m \in \Pi_m$ is an arbitrary polynomial of degree $\leq m$. We define

$$E_m(g) = \|g - P_m^*\| = \inf_{P_m \in \Pi_m} \|g - P_m\|_\infty. \quad (3.6)$$

THEOREM 3.1. *Under the assumptions of Theorem 2.2,*

$$|f(x) - L_n(x)| \leq \frac{E_m(f^{(k+1)})}{(k+1)!} C_k(x) |\omega_n(x)|, \quad (3.7)$$

where $C_k(\cdot)$ and $E_m(\cdot)$ are defined by (3.2) and (3.6), respectively.

Proof. Let $P_m(t) = P_m^*(t)$, where $P_m^*(t)$ is defined by (3.6) for the function $g(t) = f^{(k+1)}(t)$. We have

$$\begin{aligned} |R_{k,m}(x)| &= \left| \frac{(-1)^k}{k!} \sum_{i=0}^n p_{ni}(x) \int_{x_i}^x [f^{(k+1)}(t) - P_m^*(t)] (t - x_i)^k dt \right| \\ &\leq \frac{\|f^{(k+1)} - P_m^*\|_\infty}{(k+1)!} C_k(x) |\omega_n(x)| \\ &= \frac{E_m(f^{(k+1)})}{(k+1)!} C_k(x) |\omega_n(x)|, \end{aligned} \quad (3.8)$$

since

$$\left| \int_{x_i}^x (t - x_i)^k dt \right| = \frac{|x - x_i|^{k+1}}{k+1}. \quad (3.9)$$

□

Remark 3.2. The above estimate has only theoretical importance, since it is difficult to find the polynomial P^* . In fact, we can find P^* only for some special cases of functions. However, we can use the estimate to obtain some practical estimations—see Theorem 3.3.

THEOREM 3.3. *Let the assumptions of Theorem 2.2 hold. If γ_{k+1} , Γ_{k+1} are real numbers such that $\gamma_{k+1} \leq f^{(k+1)}(t) \leq \Gamma_{k+1}$, $t \in [a, b]$, $k = 0, 1, \dots, n-1$, then*

$$|f(x) - L_n(x)| \leq \frac{\Gamma_{k+1} - \gamma_{k+1}}{2(k+1)!} C_k(x) |\omega_n(x)|, \quad (3.10)$$

where ω_n and $C_k(\cdot)$ are defined by (3.1) and (3.2), respectively. Also

$$\begin{aligned} |f(x) - L_n(x)| &\leq \frac{|\omega_n(x)|}{k!} B_k(x), \\ |f(x) - L_n(x)| &\leq \frac{|\omega_n(x)|}{k!} D_k(x), \end{aligned} \quad (3.11)$$

where $B_k(\cdot)$ and $D_k(\cdot)$ are defined by (3.3) and (3.4), respectively.

Proof. We set $P_m(t) = (\Gamma_{k+1} + \gamma_{k+1})/2$ in (2.12). Then we have

$$|f(x) - L_n(x)| = |R_k(x)| \leq \frac{1}{k!} \sum_{i=0}^n |p_{ni}(x)| \left\| f^{(k+1)} - \frac{\Gamma_{k+1} + \gamma_{k+1}}{2} \right\|_{\infty} \left| \int_{x_i}^x (t - x_i)^k dt \right|. \quad (3.12)$$

We also have

$$\begin{aligned} \left\| f^{(k+1)} - \frac{\Gamma_{k+1} + \gamma_{k+1}}{2} \right\|_{\infty} &\leq \frac{\Gamma_{k+1} - \gamma_{k+1}}{2}, \\ \left| \int_{x_i}^x (t - x_i)^k dt \right| &= \frac{|x - x_i|^{k+1}}{k+1}. \end{aligned} \quad (3.13)$$

From the above three relations we get

$$\begin{aligned} |f(x) - L_n(x)| &\leq \frac{\Gamma_{k+1} - \gamma_{k+1}}{2(k+1)!} \sum_{i=0}^n |p_{ni}(x)| |x - x_i|^{k+1} \\ &= \frac{\Gamma_{k+1} - \gamma_{k+1}}{2(k+1)!} C_k(x) |\omega_n(x)|. \end{aligned} \quad (3.14)$$

The first inequality is proved.

We now set $P_m(t) = \gamma_{k+1}$ in (2.12). Then we have

$$|f(x) - L_n(x)| = |R_k(x)| \leq \frac{1}{k!} \sum_{i=0}^n |p_{ni}(x)| \left| \int_{x_i}^x [f^{(k+1)}(t) - \gamma_{k+1}] (t - x_i)^k dt \right|. \quad (3.15)$$

We also have

$$\begin{aligned} \left| \int_{x_i}^x [f^{(k+1)}(t) - \gamma_{k+1}](t - x_i)^k dt \right| &\leq |x - x_i|^k |f^{(k)}(x) - f^{(k)}(x_i) - \gamma_{k+1}(x - x_i)| \\ &= |x - x_i|^{k+1} (S_{ki} - \gamma_{k+1}). \end{aligned} \quad (3.16)$$

Thus,

$$\begin{aligned} |f(x) - L_n(x)| &\leq \frac{1}{k!} \sum_{i=0}^n |p_{ni}(x)| |x - x_i|^{k+1} (S_{ki} - \gamma_{k+1}) \\ &= \frac{|\omega_n(x)|}{k!} B_k(x). \end{aligned} \quad (3.17)$$

The second inequality is proved. In a similar way we prove that the third inequality holds. $\square$

LEMMA 3.4. Let $D = \{x_0 = a < x_1 < \dots < x_n = b\}$ be a given uniform subdivision of the interval $[a, b]$, that is, $x_i = x_0 + ih$, $h = (b - a)/n$, $i = 0, 1, 2, \dots, n$. If $x \in (x_{j-1}, x_j)$, for some $j \in \{1, 2, \dots, n\}$, then

$$|\omega_n(x)| \leq j!(n - j + 1)!h^{n+1}, \quad (3.18)$$

$$C_k(x) \leq \frac{2^n}{n!} \left\{ \frac{1}{2} [n + 1 + |n - 2j + 1|] \right\}^k h^{k-n}, \quad (3.19)$$

$$C_k(x) |\omega_n(x)| \leq \alpha_{jnk} \frac{n - j + 1}{n} \frac{2^n (b - a)^{k+1}}{\binom{n}{j}}, \quad (3.20)$$

where

$$\alpha_{jnk} = \left[\frac{1}{2n} (n + 1 + |2j - n - 1|) \right]^k. \quad (3.21)$$

This lemma is proved in [10].

Remark 3.5. Note that

$$\alpha_{jnk} \leq 1 \quad (3.22)$$

and $\alpha_{jnk} = 1$ if and only if $j = 1$ or $j = n$. If we choose $x \in [x_j, x_{j+1}]$, $j = 0, 1, \dots, n - 1$, then we get the factor $(j + 1)/n$ instead of the factor $(n - j + 1)/n$ in (3.20).

THEOREM 3.6. Under the assumptions of Lemma 3.4 and Theorem 3.3,

$$|f(x) - L_n(x)| \leq \frac{\Gamma_{k+1} - \gamma_{k+1}}{(k + 1)!} \alpha_{jnk} \frac{n - j + 1}{n} \frac{2^{n-1} (b - a)^{k+1}}{\binom{n}{j}}. \quad (3.23)$$

Proof. The proof follows immediately from Theorem 3.3 and Lemma 3.4. $\square$

4. Results for derivatives

LEMMA 4.1. *Let $1 \leq j \leq n-1$ and $j+1 \leq r \leq n$. Then*

$$\sum_{i=0}^n p_{ni}^{(j)}(x)(x-x_i)^r = 0. \quad (4.1)$$

Proof. We have (see (2.4))

$$A(x) = \sum_{i=0}^n p_{ni}(x)(x-x_i)^r = 0, \quad \text{for } 1 \leq r \leq n. \quad (4.2)$$

Thus,

$$A'(x) = \sum_{i=0}^n p'_{ni}(x)(x-x_i)^r + r \sum_{i=0}^n p_{ni}(x)(x-x_i)^{r-1} = 0, \quad (4.3)$$

if $1 \leq r \leq n$. If $n \geq r-1 \geq 1$, that is, $n+1 \geq r \geq 2$, then

$$r \sum_{i=0}^n p_{ni}(x)(x-x_i)^{r-1} = 0. \quad (4.4)$$

From (4.3) and (4.4) we get

$$\sum_{i=0}^n p'_{ni}(x)(x-x_i)^r = 0, \quad \text{for } 2 \leq r \leq n. \quad (4.5)$$

(Note that $\{r : 1 \leq r \leq n\} \cap \{r : 2 \leq r \leq n+1\} = \{r : 2 \leq r \leq n\}$. Here we use this fact and similar facts without a special mentioning.)

We now suppose that

$$\sum_{i=0}^n p_{ni}^{(j)}(x)(x-x_i)^r = 0, \quad (4.6)$$

for $j = 1, 2, \dots, m$, $m < n-1$ and $j+1 \leq r \leq n$. We wish to prove that

$$\sum_{i=0}^n p_{ni}^{(m+1)}(x)(x-x_i)^r = 0, \quad \text{for } m+2 \leq r \leq n. \quad (4.7)$$

For that purpose, we first calculate

$$\begin{aligned} A^{(m)}(x) &= \sum_{i=0}^n [p_{ni}(x)(x-x_i)^r]^{(m)} \\ &= \sum_{i=0}^n \sum_{k=0}^m \binom{m}{k} p_{ni}^{(k)}(x) \frac{r!}{(r-m+k)!} (x-x_i)^{r-m+k} \\ &= \sum_{k=0}^m \binom{m}{k} \frac{r!}{(r-m+k)!} \sum_{i=0}^n p_{ni}^{(k)}(x)(x-x_i)^{r-m+k}. \end{aligned} \quad (4.8)$$

We have

$$A^{(m)}(x) = 0, \quad \text{for } r \geq m+1, \quad (4.9)$$

by the above assumption. Thus,

$$A^{(m+1)}(x) = 0. \quad (4.10)$$

On the other hand, we have

$$\begin{aligned} A^{(m+1)}(x) &= \frac{d}{dx} A^{(m)}(x) \\ &= \sum_{k=0}^m \binom{m}{k} \frac{r!}{(r-m+k)!} \sum_{i=0}^n p_{ni}^{(k+1)}(x) (x-x_i)^{r-m+k} \\ &\quad + \sum_{k=0}^m \binom{m}{k} \frac{r!}{(r-m+k-1)!} \sum_{i=0}^n p_{ni}^{(k)}(x) (x-x_i)^{r-m+k-1} \\ &= 0. \end{aligned} \quad (4.11)$$

We now rewrite the above relation in the form

$$\begin{aligned} \sum_{i=0}^n p_{ni}^{(m+1)}(x) (x-x_i)^r + \sum_{k=0}^{m-1} \binom{m}{k} \frac{r!}{(r-m+k)!} \sum_{i=0}^n p_{ni}^{(k+1)}(x) (x-x_i)^{r-m+k} \\ + \sum_{k=0}^m \binom{m}{k} \frac{r!}{(r-m+k-1)!} \sum_{i=0}^n p_{ni}^{(k)}(x) (x-x_i)^{r-m+k-1} = 0. \end{aligned} \quad (4.12)$$

For $r-m+k-1 \geq k+1$, that is, $r \geq m+2$, we have

$$\sum_{i=0}^n p_{ni}^{(k)}(x) (x-x_i)^{r-m+k-1} = 0 \quad (4.13)$$

by the above assumption. We also have

$$\sum_{i=0}^n p_{ni}^{(k+1)}(x) (x-x_i)^{r-m+k} = 0, \quad (4.14)$$

if $r-m+k \geq k+2$, that is, $r \geq m+2$. Thus (4.7) holds. This completes the proof. $\square$

THEOREM 4.2. *Let $f \in C^{n+1}(a, b)$ and let $P_r(t)$ be an arbitrary polynomial of degree $\leq r$ and let $0 \leq k \leq n$, $1 \leq m \leq k$. Then*

$$f^{(m)}(x) = L_n^{(m)}(x) + E_{k,r}(x), \quad (4.15)$$

where

$$E_{k,r}(x) = \frac{(-1)^k}{k!} \sum_{i=0}^n p_{ni}^{(m)}(x) \int_{x_i}^x [f^{(k+1)}(t) - P_r(t)] (t-x_i)^k dt. \quad (4.16)$$

Proof. We define

$$\begin{aligned} v_i(x) &= \int_{x_i}^x [f^{(k+1)}(t) - P_r(t)](t - x_i)^k dt \\ &= \int_{x_i}^x g(t)(t - x_i)^k dt, \end{aligned} \quad (4.17)$$

where, obviously, $g(t) = f^{(k+1)}(t) - P_r(t)$. We denote

$$R_{k,r}(x) = f(x) - L_n(x) = \frac{(-1)^k}{k!} \sum_{i=0}^n p_{ni}(x) v_i(x), \quad (4.18)$$

see Theorem 2.2. Then we have

$$\begin{aligned} R_{k,r}^{(m)}(x) &= \frac{(-1)^k}{k!} \sum_{i=0}^n [p_{ni}(x) v_i(x)]^{(m)} \\ &= \frac{(-1)^k}{k!} \sum_{i=0}^n \sum_{j=0}^m \binom{m}{j} p_{ni}^{(j)}(x) v_i^{(m-j)}(x) \\ &= \frac{(-1)^k}{k!} \sum_{i=0}^n p_{ni}^{(m)}(x) v_i(x) + \frac{(-1)^k}{k!} \sum_{i=0}^n \sum_{j=0}^{m-1} \binom{m}{j} p_{ni}^{(j)}(x) v_i^{(m-j)}(x). \end{aligned} \quad (4.19)$$

We introduce the notation

$$B(x) = \frac{(-1)^k}{k!} \sum_{i=0}^n \sum_{j=0}^{m-1} \binom{m}{j} p_{ni}^{(j)}(x) v_i^{(m-j)}(x) \quad (4.20)$$

such that

$$R_{k,r}^{(m)}(x) = \frac{(-1)^k}{k!} \sum_{i=0}^n p_{ni}^{(m)}(x) v_i(x) + B(x). \quad (4.21)$$

We now rewrite $B(x)$ in the form

$$B(x) = \frac{(-1)^k}{k!} \sum_{i=0}^n \sum_{j=0}^{m-2} \binom{m}{j} p_{ni}^{(j)}(x) v_i^{(m-j)}(x) + \frac{(-1)^k}{k!} m \sum_{i=0}^n p_{ni}^{(m-1)}(x) v_i'(x). \quad (4.22)$$

We have

$$v_i'(x) = g(x)(x - x_i)^k \quad (4.23)$$

such that

$$\sum_{i=0}^n p_{ni}^{(m-1)}(x) v_i'(x) = g(x) \sum_{i=0}^n p_{ni}^{(m-1)}(x) (x - x_i)^k = 0, \quad (4.24)$$

for $k \geq m$ —see Lemma 4.1.

We also have

$$v_i^{(m-j)}(x) = \sum_{l=0}^{m-j-1} \binom{m-j-1}{l} g^{(l)}(x) \frac{k!}{(k-m+j+l+1)!} (x-x_i)^{k-m+j+l+1}, \quad (4.25)$$

for $m \geq j+2$ such that

$$\begin{aligned} \sum_{i=0}^n \sum_{j=0}^{m-2} \binom{m}{j} p_{ni}^{(j)}(x) v_i^{(m-j)}(x) &= \sum_{j=0}^{m-2} \binom{m}{j} \sum_{l=0}^{m-j-1} \binom{m-j-1}{l} \frac{k!}{(k-m+j+l+1)!} \\ &\quad \times \sum_{i=0}^n p_{ni}^{(j)}(x) (x-x_i)^{k-m+j+l+1} \\ &= 0, \end{aligned} \quad (4.26)$$

if $k-m+j+l+1 \geq j+1$, that is, $k \geq m$, since $l \geq 0$ —see also Lemma 4.1. Hence, $B(x) = 0$ in all cases. Now from (4.21) it follows that

$$\begin{aligned} R_{k,r}^{(m)}(x) &= \frac{(-1)^k}{k!} \sum_{i=0}^n p_{ni}^{(m)}(x) v_i(x) \\ &= \frac{(-1)^k}{k!} \sum_{i=0}^n p_{ni}^{(m)}(x) \int_{x_i}^x [f^{(k+1)}(t) - P_r(t)] (t-x_i)^k dt. \end{aligned} \quad (4.27)$$

On the other hand, we have

$$[f(x) - L_n(x)]^{(m)} = f^{(m)}(x) - L_n^{(m)}(x). \quad (4.28)$$

This completes the proof. $\square$

THEOREM 4.3. *Under the assumptions of Theorem 4.2,*

$$|f^{(m)}(x) - L_n^{(m)}(x)| \leq \frac{E_r(f^{(k+1)})}{(k+1)!} \sum_{i=0}^n |p_{ni}^{(m)}(x)| |x-x_i|^{k+1}, \quad (4.29)$$

where $E_r(\cdot)$ is defined by (3.6).

Proof. Let $P_r(t) = P_r^*(t)$, where $P_r^*(t)$ is defined by (3.6) for the function $g(t) = f^{(k+1)}(t)$. We have

$$\begin{aligned} |R_{k,r}^{(m)}(x)| &= \left| \frac{(-1)^k}{k!} \sum_{i=0}^n p_{ni}^{(m)}(x) \int_{x_i}^x [f^{(k+1)}(t) - P_r^*(t)] (t-x_i)^k dt \right| \\ &\leq \frac{\|f^{(k+1)}(t) - P_r^*(t)\|_\infty}{(k+1)!} \sum_{i=0}^n |p_{ni}^{(m)}(x)| |x-x_i|^{k+1} \\ &= \frac{E_r(f^{(k+1)})}{(k+1)!} \sum_{i=0}^n |p_{ni}^{(m)}(x)| |x-x_i|^{k+1}, \end{aligned} \quad (4.30)$$

since

$$\left| \int_{x_i}^x (t - x_i)^k dt \right| = \frac{|x - x_i|^{k+1}}{k+1}. \quad (4.31)$$

□

THEOREM 4.4. Under the assumptions of Theorem 3.3 and Lemma 4.1,

$$\begin{aligned} |f^{(m)}(x) - L_n^{(m)}(x)| &\leq \frac{\Gamma_{k+1} - \gamma_{k+1}}{2(k+1)!} \sum_{i=0}^n |p_{ni}^{(m)}(x)| |x - x_i|^{k+1}, \\ |f^{(m)}(x) - L_n^{(m)}(x)| &\leq \frac{1}{k!} \sum_{i=0}^n (S_{ki} - \gamma_{k+1}) |p_{ni}^{(m)}(x)| |x - x_i|^{k+1}, \\ |f^{(m)}(x) - L_n^{(m)}(x)| &\leq \frac{1}{k!} \sum_{i=0}^n (\Gamma_{k+1} - S_{ki}) |p_{ni}^{(m)}(x)| |x - x_i|^{k+1}. \end{aligned} \quad (4.32)$$

Proof. We choose $P_r(t) = \Gamma_{k+1} + \gamma_{k+1}/2$ in Theorem 4.2. Then we get

$$\begin{aligned} |f^{(m)}(x) - L_n^{(m)}(x)| &\leq \frac{1}{k!} \sum_{i=0}^n |p_{ni}^{(m)}(x)| \left| \int_{x_i}^x \left[f^{(k+1)}(t) - \frac{\Gamma_{k+1} + \gamma_{k+1}}{2} \right] (t - x_i)^k dt \right| \\ &\leq \frac{\Gamma_{k+1} - \gamma_{k+1}}{2(k!)} \sum_{i=0}^n |p_{ni}^{(m)}(x)| \left| \int_{x_i}^x (t - x_i)^k dt \right| \\ &= \frac{\Gamma_{k+1} - \gamma_{k+1}}{2(k+1)!} \sum_{i=0}^n |p_{ni}^{(m)}(x)| |x - x_i|^{k+1}. \end{aligned} \quad (4.33)$$

If we choose $P_r(t) = \gamma_{k+1}$ in Theorem 4.2, then we get

$$\begin{aligned} |f^{(m)}(x) - L_n^{(m)}(x)| &\leq \frac{1}{k!} \sum_{i=0}^n |p_{ni}^{(m)}(x)| \left| \int_{x_i}^x [f^{(k+1)}(t) - \gamma_{k+1}] (t - x_i)^k dt \right| \\ &\leq \frac{1}{k!} \sum_{i=0}^n (S_{ki} - \gamma_{k+1}) |p_{ni}^{(m)}(x)| |x - x_i|^{k+1}, \end{aligned} \quad (4.34)$$

since $\left| \int_{x_i}^x [f^{(k+1)}(t) - \gamma_{k+1}] dt \right| = |f^{(k)}(x) - f^{(k)}(x_i) - \gamma_{k+1}(x - x_i)|$.

In a similar way we prove that the third inequality holds. □

References

- [1] R. P. Agarwal and P. J. Y. Wong, *Error Inequalities in Polynomial Interpolation and Their Applications*, Mathematics and Its Applications, vol. 262, Kluwer Academic, Dordrecht, 1993.
- [2] P. Cerone and S. S. Dragomir, *Midpoint-type rules from an inequalities point of view*, Handbook of Analytic-Computational Methods in Applied Mathematics (G. Anastassiou, ed.), Chapman & Hall/CRC, Florida, 2000, pp. 135–200.
- [3] ———, *Trapezoidal-type rules from an inequalities point of view*, Handbook of Analytic-Computational Methods in Applied Mathematics (G. Anastassiou, ed.), Chapman & Hall/CRC, Florida, 2000, pp. 65–134.
- [4] X.-L. Cheng, *Improvement of some Ostrowski-Grüss type inequalities*, Comput. Math. Appl. **42** (2001), no. 1-2, 109–114.

- [5] S. S. Dragomir, R. P. Agarwal, and P. Cerone, *On Simpson's inequality and applications*, J. Inequal. Appl. **5** (2000), no. 6, 533–579.
- [6] S. S. Dragomir and S. Wang, *An inequality of Ostrowski-Grüss type and its applications to the estimation of error bounds for some special means and for some numerical quadrature rules*, Comput. Math. Appl. **33** (1997), no. 11, 15–20.
- [7] H. N. Mhaskar and D. V. Pai, *Fundamentals of Approximation Theory*, CRC Press, Florida; Narosa, New Delhi, 2000.
- [8] D. S. Mitrinović, J. E. Pečarić, and A. M. Fink, *Classical and New Inequalities in Analysis*, Mathematics and Its Applications (East European Series), vol. 61, Kluwer Academic, Dordrecht, 1993.
- [9] C. E. M. Pearce, J. E. Pečarić, N. Ujević, and S. Varošanec, *Generalizations of some inequalities of Ostrowski-Grüss type*, Math. Inequal. Appl. **3** (2000), no. 1, 25–34.
- [10] N. Ujević, *Error inequalities for a corrected interpolating polynomial*, New York J. Math. **10** (2004), 69–81.
- [11] ———, *Error inequalities for a perturbed interpolating polynomial*, Nonlinear Stud. **12** (2005), no. 3, 233–245.
- [12] N. Ujević and A. J. Roberts, *A corrected quadrature formula and applications*, ANZIAM J. **45** (2004), no. (E), E41–E56.

Nenad Ujević: Department of Mathematics, University of Split, Teslina 12/III, 21000 Split, Croatia
E-mail address: ujevic@pmfst.hr