FIXED POINT ITERATION FOR ASYMPTOTICALLY QUASI-NONEXPANSIVE MAPPINGS IN BANACH SPACES

SOMYOT PLUBTIENG AND RABIAN WANGKEEREE

Received 21 October 2004 and in revised form 20 April 2005

Suppose that C is a nonempty closed convex subset of a real uniformly convex Banach space X . Let $T : C \rightarrow C$ be an asymptotically quasi-nonexpansive mapping. In this paper, we introduce the three-step iterative scheme for such map with error members. Moreover, we prove that if T is uniformly L -Lipschitzian and completely continuous, then the iterative scheme converges strongly to some fixed point of T .

1. Introduction

Let C be a subset of normed space X , and let T be a self-mapping on C . T is said to be *nonexpansive* provided that $\|Tx - Ty\| \leq \|x - y\|$ for all $x, y \in C$; T is called *asymptotically nonexpansive* if there exists a sequence $\{k_n\}$ in $[0, \infty)$ with $\lim_{n \rightarrow \infty} k_n = 0$ such that $\|T^n x - T^n y\| \leq (1 + k_n)\|x - y\|$ for all $x, y \in C$ and $n \geq 1$. T is said to be an *asymptotically quasi-nonexpansive map* if there exists a sequence $\{k_n\}$ in $[0, \infty)$ with $\lim_{n \rightarrow \infty} k_n = 0$ such that $\|T^n x - p\| \leq (1 + k_n)\|x - p\|$ for all $x \in C$ and $p \in F(T)$, and $n \geq 1$ ($F(T)$ denotes the set of fixed points of T , that is, $F(T) = \{x \in C : Tx = x\}$).

From the above definitions, if $F(T) \neq \emptyset$, then asymptotically nonexpansive mapping must be asymptotically quasi-nonexpansive mapping.

The concept of asymptotically nonexpansiveness was introduced by Goebel and Kirk in 1972 [2]. In 2001, Noor [5, 6] introduced the three-step iterative scheme and he studied the approximate solutions of variational inclusions (inequalities) in Hilbert spaces. The three-step iterative approximation problems were studied extensively by Noor [5, 6], Glowinski and Le Tallec [1], and Haubruge et al. [3].

Recently, Xu and Noor [8] introduced the three-step iterative scheme for asymptotically nonexpansive mappings and they proved the following strong convergence theorem in Banach spaces.

THEOREM 1.1 (see [8, Theorem 2.1]). *Let X be a real uniformly convex Banach space, let C be a nonempty closed, bounded convex subset of X . Let T be a completely continuous and asymptotically nonexpansive self-mapping with sequence $\{k_n\}$ satisfying $k_n \geq 0$ and*

$\sum_{n=1}^{\infty} k_n < \infty$. Let $\{\alpha_n\}$, $\{\beta_n\}$, and $\{\gamma_n\}$ be real sequences in $[0, 1]$ satisfying

$$(i) \ 0 < \liminf_{n \rightarrow \infty} \alpha_n \leq \limsup_{n \rightarrow \infty} \alpha_n < 1,$$

$$(ii) \ 0 < \liminf_{n \rightarrow \infty} \beta_n \leq \limsup_{n \rightarrow \infty} \beta_n < 1.$$

For a given $x_0 \in D$, define

$$\begin{aligned} z_n &= \gamma_n T^n x_n + (1 - \gamma_n) x_n, \\ y_n &= \beta_n T^n z_n + (1 - \beta_n) x_n, \\ x_{n+1} &= \alpha_n T^n y_n + (1 - \alpha_n) x_n. \end{aligned} \quad (1.1)$$

Then $\{x_n\}$, $\{y_n\}$, and $\{z_n\}$ converge strongly to a fixed point of T .

In this paper, we will extend the iterative scheme (1.1) to the iterative scheme of asymptotically quasi-nonexpansive mappings with error members. Moreover, we will prove the strong convergence of iterative scheme to a fixed point of T (C need not to be a bounded set), requiring T to be uniformly L -Lipschitzian and completely continuous. The results presented in this paper generalize and extend the corresponding main results of Xu and Noor [8].

2. Preliminaries

For the sake of convenience, we first recall some definitions and conclusions.

Definition 2.1 (see [2]). A Banach space X is said to be *uniformly convex* if the modulus of convexity of X

$$\delta_X(\epsilon) = \inf \left\{ 1 - \frac{\|x+y\|}{2} : \|x\| = \|y\| = 1, \|x-y\| = \epsilon \right\} > 0 \quad (2.1)$$

for all $0 < \epsilon \leq 2$ (i.e., $\delta_X(\epsilon)$ is a function $(0, 2] \rightarrow (0, 1)$).

Definition 2.2. A mapping $T : C \rightarrow C$ is called *uniformly L -Lipschitzian* if there exists a constant $L > 0$ such that for all $x, y \in C$,

$$\|T^n x - T^n y\| \leq L \|x - y\|, \quad \forall n \geq 1. \quad (2.2)$$

In what follows, we will make use of the following lemmas.

LEMMA 2.3 (see [4]). Let the nonnegative number sequences $\{a_n\}$, $\{b_n\}$, and $\{d_n\}$ satisfy that

$$a_{n+1} \leq (1 + b_n) a_n + d_n, \quad \forall n = 1, 2, \dots, \sum_{n=1}^{\infty} b_n < \infty, \sum_{n=1}^{\infty} d_n < \infty. \quad (2.3)$$

Then,

- (1) $\lim_{n \rightarrow \infty} a_n$ exists;
- (2) if $\liminf_{n \rightarrow \infty} a_n = 0$, then $\lim_{n \rightarrow \infty} a_n = 0$.

LEMMA 2.4 ([7], J. Schu's Lemma). *Let X be a real uniformly convex Banach space, $0 < \alpha \leq t_n \leq \beta < 1$, $x_n, y_n \in X$, $\limsup_{n \rightarrow \infty} \|x_n\| \leq a$, $\limsup_{n \rightarrow \infty} \|y_n\| \leq a$, and $\lim_{n \rightarrow \infty} \|t_n x_n + (1 - t_n)y_n\| = a$, $a \geq 0$. Then, $\lim_{n \rightarrow \infty} \|x_n - y_n\| = 0$.*

3. Main results

In this section, we prove our main theorem. First of all, we will need the following lemmas.

LEMMA 3.1. *Let X be a real uniformly convex Banach space, C a nonempty closed convex subset of X . Let T be an asymptotically quasi-nonexpansive mapping with sequence $\{k_n\}_{n \geq 1}$ such that $\sum_{n=1}^{\infty} k_n < \infty$ and $F(T) \neq \emptyset$. Let $x_0 \in C$ and*

$$\begin{aligned} z_n &= \alpha_n'' T^n x_n + \beta_n'' x_n + \gamma_n'' u_n, \\ y_n &= \alpha_n' T^n z_n + \beta_n' x_n + \gamma_n' v_n, \\ x_{n+1} &= \alpha_n T^n y_n + \beta_n x_n + \gamma_n w_n, \end{aligned} \quad (3.1)$$

where $\{\alpha_n\}$, $\{\alpha_n'\}$, $\{\alpha_n''\}$, $\{\beta_n\}$, $\{\beta_n'\}$, $\{\beta_n''\}$, $\{\gamma_n\}$, $\{\gamma_n'\}$, and $\{\gamma_n''\}$ are real sequences in $[0, 1]$ and $\{u_n\}$, $\{v_n\}$, and $\{w_n\}$ are three bounded sequences in C such that

$$(i) \alpha_n + \beta_n + \gamma_n = \alpha_n' + \beta_n' + \gamma_n' = \alpha_n'' + \beta_n'' + \gamma_n'' = 1,$$

$$(ii) \sum_{n=1}^{\infty} \gamma_n < \infty, \sum_{n=1}^{\infty} \gamma_n' < \infty, \sum_{n=1}^{\infty} \gamma_n'' < \infty.$$

If $p \in F(T)$, then $\lim_{n \rightarrow \infty} \|x_n - p\|$ exists.

Proof. Let $p \in F(T)$. Since $\{u_n\}$, $\{v_n\}$, and $\{w_n\}$ are bounded sequences in C , put

$$M = \sup_{n \geq 1} \|u_n - p\| \vee \sup_{n \geq 1} \|v_n - p\| \vee \sup_{n \geq 1} \|w_n - p\|. \quad (3.2)$$

Then M is a finite number. So for each $n \geq 1$, we note that

$$\begin{aligned} \|x_{n+1} - p\| &= \|\alpha_n T^n y_n + \beta_n x_n + \gamma_n w_n - p\| \\ &\leq \alpha_n \|T^n y_n - p\| + \beta_n \|x_n - p\| + \gamma_n \|w_n - p\| \\ &\leq \alpha_n (1 + k_n) \|y_n - p\| + \beta_n \|x_n - p\| + \gamma_n \|w_n - p\|, \end{aligned} \quad (3.3)$$

$$\begin{aligned} \|y_n - p\| &= \|\alpha_n' T^n z_n + \beta_n' x_n + \gamma_n' v_n - p\| \\ &\leq \alpha_n' \|T^n z_n - p\| + \beta_n' \|x_n - p\| + \gamma_n' \|v_n - p\| \\ &\leq \alpha_n' (1 + k_n) \|z_n - p\| + \beta_n' \|x_n - p\| + \gamma_n' \|v_n - p\|, \end{aligned} \quad (3.4)$$

$$\|z_n - p\| \leq \alpha_n'' (1 + k_n) \|x_n - p\| + \beta_n'' \|x_n - p\| + \gamma_n'' \|u_n - p\|. \quad (3.5)$$

Substituting (3.5) into (3.4),

$$\begin{aligned}
\|y_n - p\| &\leq \alpha'_n \alpha''_n (1 + k_n)^2 \|x_n - p\| \\
&\quad + \alpha'_n \beta''_n (1 + k_n) \|x_n - p\| + \alpha'_n \gamma''_n (1 + k_n) \|u_n - p\| + \beta'_n \|x_n - p\| + \gamma'_n \|v_n - p\| \\
&\leq (1 - \beta'_n - \gamma'_n) \alpha''_n (1 + k_n)^2 \|x_n - p\| + \beta'_n \|x_n - p\| \\
&\quad + (1 - \beta'_n - \gamma'_n) \beta''_n \|x_n - p\| + m_n \\
&\leq \beta'_n (1 + k_n)^2 \|x_n - p\| + (1 - \beta'_n) \alpha''_n (1 + k_n)^2 \|x_n - p\| \\
&\quad + (1 - \beta'_n) \beta''_n (1 + k_n)^2 \|x_n - p\| + m_n \\
&= \beta'_n (1 + k_n)^2 \|x_n - p\| + (1 - \beta'_n) (\alpha''_n + \beta''_n) (1 + k_n)^2 \|x_n - p\| + m_n \\
&\leq \beta'_n (1 + k_n)^2 \|x_n - p\| + (1 - \beta'_n) (1 + k_n)^2 \|x_n - p\| + m_n \\
&= (1 + k_n)^2 \|x_n - p\| + m_n,
\end{aligned} \tag{3.6}$$

where $m_n = \gamma''_n (1 + k_n)M + \gamma'_n M$. Substituting (3.6) into (3.3) again, we have

$$\begin{aligned}
\|x_{n+1} - p\| &\leq \alpha_n (1 + k_n) ((1 + k_n)^2 \|x_n - p\| + m_n) + \beta_n \|x_n - p\| + \gamma_n \|w_n - p\| \\
&= \alpha_n (1 + k_n)^3 \|x_n - p\| + \alpha_n (1 + k_n) m_n + \beta_n \|x_n - p\| + \gamma_n \|w_n - p\| \\
&\leq (\alpha_n + \beta_n) (1 + k_n)^3 \|x_n - p\| + (1 + k_n) m_n + \gamma_n \|w_n - p\| \\
&\leq (1 + k_n)^3 \|x_n - p\| + (1 + k_n) m_n + \gamma_n \|w_n - p\| \\
&\leq (1 + k_n)^3 \|x_n - p\| + (1 + k_n) m_n + \gamma_n M \\
&= (1 + d_n) \|x_n - p\| + b_n,
\end{aligned} \tag{3.7}$$

where $d_n = 3k_n + 3k_n^2 + k_n^3$ and $b_n = (1 + k_n)m_n + \gamma_n M$. Since $\sum_{n=1}^{\infty} d_n < \infty$ and $\sum_{n=1}^{\infty} b_n < \infty$, by Lemma 2.3, we have that $\lim_{n \rightarrow \infty} \|x_n - p\|$ exists. This completes the proof. $\square$

LEMMA 3.2. *Let X be a real uniformly convex Banach space, C a nonempty closed convex subset of X . Let T be an asymptotically quasi-nonexpansive mapping with sequence $\{k_n\}_{n \geq 1}$ such that $\sum_{n=1}^{\infty} k_n < \infty$ and $F(T) \neq \emptyset$. Let $x_0 \in C$ and for each $n \geq 0$,*

$$\begin{aligned}
z_n &= \alpha''_n T^n x_n + \beta''_n x_n + \gamma''_n u_n, \\
y_n &= \alpha'_n T^n z_n + \beta'_n x_n + \gamma'_n v_n, \\
x_{n+1} &= \alpha_n T^n y_n + \beta_n x_n + \gamma_n w_n,
\end{aligned} \tag{3.8}$$

where $\{u_n\}$, $\{v_n\}$, and $\{w_n\}$ are three bounded sequences in C and $\{\alpha_n\}$, $\{\alpha'_n\}$, $\{\alpha''_n\}$, $\{\beta_n\}$, $\{\beta'_n\}$, $\{\beta''_n\}$, $\{\gamma_n\}$, $\{\gamma'_n\}$, and $\{\gamma''_n\}$ are real sequences in $[0, 1]$ which satisfy the same assumptions as Lemma 3.1 and the additional assumption that $0 \leq \alpha < \alpha_n$, $\beta_n, \alpha'_n, \beta'_n \leq \beta < 1$ for some α, β in $(0, 1)$. Then $\lim_{n \rightarrow \infty} \|T^n y_n - x_n\| = 0 = \lim_{n \rightarrow \infty} \|T^n z_n - x_n\|$.

Proof. For any $p \in F(T)$, it follows from Lemma 3.1, that $\lim_{n \rightarrow \infty} \|x_n - p\|$ exists. Let $\lim_{n \rightarrow \infty} \|x_n - p\| = a$ for some $a \geq 0$. From (3.6), we have

$$\|y_n - p\| \leq (1 + k_n)^2 \|x_n - p\| + m_n. \quad (3.9)$$

Taking $\limsup_{n \rightarrow \infty}$ in both sides, we obtain

$$\limsup_{n \rightarrow \infty} \|y_n - p\| \leq \limsup_{n \rightarrow \infty} \|x_n - p\| = \lim_{n \rightarrow \infty} \|x_n - p\| = a. \quad (3.10)$$

Note that

$$\begin{aligned} \limsup_{n \rightarrow \infty} \|T^n y_n - p\| &\leq \limsup_{n \rightarrow \infty} (1 + k_n) \|y_n - p\| = \limsup_{n \rightarrow \infty} \|y_n - p\| \leq a, \\ a = \lim_{n \rightarrow \infty} \|x_{n+1} - p\| &= \lim_{n \rightarrow \infty} \|\alpha_n T^n y_n + \beta_n x_n + \gamma_n w_n - p\| \\ &= \lim_{n \rightarrow \infty} \left\| \alpha_n \left[T^n y_n - p + \frac{\gamma_n}{2\alpha_n} (w_n - p) \right] + \beta_n \left[x_n - p + \frac{\gamma_n}{2\beta_n} (w_n - p) \right] \right\| \\ &= \lim_{n \rightarrow \infty} \left\| \alpha_n \left[T^n y_n - p + \frac{\gamma_n}{2\alpha_n} (w_n - p) \right] + (1 - \alpha_n) \left[x_n - p + \frac{\gamma_n}{2\beta_n} (w_n - p) \right] \right\|. \end{aligned} \quad (3.11)$$

By J. Schu's Lemma 2.4, we have

$$\lim_{n \rightarrow \infty} \left\| T^n y_n - x_n + \left(\frac{\gamma_n}{2\alpha_n} - \frac{\gamma_n}{2\beta_n} \right) (w_n - p) \right\| = 0. \quad (3.12)$$

Since $\lim_{n \rightarrow \infty} \|(\gamma_n/2\alpha_n - \gamma_n/2\beta_n)(w_n - p)\| = 0$, it follows that

$$\lim_{n \rightarrow \infty} \|T^n y_n - x_n\| = 0. \quad (3.13)$$

Finally, we will prove that $\lim_{n \rightarrow \infty} \|T^n z_n - x_n\| = 0$. To this end, we note that for each $n \geq 1$,

$$\|x_n - p\| \leq \|T^n y_n - x_n\| + \|T^n y_n - p\| \leq \|T^n y_n - x_n\| + (1 + k_n) \|y_n - p\|. \quad (3.14)$$

Since $\lim_{n \rightarrow \infty} \|T^n y_n - x_n\| = 0 = \lim_{n \rightarrow \infty} k_n$, we obtain that

$$a = \lim_{n \rightarrow \infty} \|x_n - p\| \leq \liminf_{n \rightarrow \infty} \|y_n - p\|. \quad (3.15)$$

It follows that

$$a \leq \liminf_{n \rightarrow \infty} \|y_n - p\| \leq \limsup_{n \rightarrow \infty} \|y_n - p\| \leq a. \quad (3.16)$$

This implies that

$$\lim_{n \rightarrow \infty} \|y_n - p\| = a. \quad (3.17)$$

On the other hand, we note that

$$\begin{aligned}
 \|z_n - p\| &= \|\alpha_n'' T^n x_n + \beta_n'' x_n + \gamma_n'' u_n - p\| \\
 &\leq \alpha_n'' (1 + k_n) \|x_n - p\| + \beta_n'' \|x_n - p\| + \gamma_n'' \|u_n - p\| \\
 &\leq \alpha_n'' (1 + k_n) \|x_n - p\| + (1 - \alpha_n'') (1 + k_n) \|x_n - p\| + \gamma_n'' \|u_n - p\| \\
 &\leq (1 + k_n) \|x_n - p\| + \gamma_n'' \|u_n - p\|.
 \end{aligned} \tag{3.18}$$

By boundedness of the sequence $\{u_n\}$ and $\lim_{n \rightarrow \infty} k_n = 0 = \lim_{n \rightarrow \infty} \gamma_n''$, we have

$$\limsup_{n \rightarrow \infty} \|z_n - p\| \leq \limsup_{n \rightarrow \infty} \|x_n - p\| = a, \tag{3.19}$$

and so

$$\begin{aligned}
 \limsup_{n \rightarrow \infty} \|T^n z_n - p\| &\leq \limsup_{n \rightarrow \infty} (1 + k_n) \|z_n - p\| \leq a, \\
 a = \lim_{n \rightarrow \infty} \|y_n - p\| &= \lim_{n \rightarrow \infty} \|\alpha_n' T^n z_n + \beta_n' x_n + \gamma_n' v_n - p\| \\
 &= \lim_{n \rightarrow \infty} \left\| \alpha_n' \left[T^n z_n - p + \frac{\gamma_n'}{2\alpha_n'} (v_n - p) \right] + \beta_n' \left[x_n - p + \frac{\gamma_n'}{2\beta_n'} (v_n - p) \right] \right\| \\
 &= \lim_{n \rightarrow \infty} \left\| \alpha_n' \left[T^n z_n - p + \frac{\gamma_n'}{2\alpha_n'} (v_n - p) \right] + (1 - \alpha_n') \left[x_n - p + \frac{\gamma_n'}{2\beta_n'} (v_n - p) \right] \right\|.
 \end{aligned} \tag{3.20}$$

By J. Schu's Lemma 2.4, we have

$$\lim_{n \rightarrow \infty} \left\| T^n z_n - x_n + \left(\frac{\gamma_n'}{2\alpha_n'} - \frac{\gamma_n'}{2\beta_n'} \right) (v_n - p) \right\| = 0. \tag{3.21}$$

Since $\lim_{n \rightarrow \infty} \|(\gamma_n'/2\alpha_n' - \gamma_n'/2\beta_n')(v_n - p)\| = 0$, it follows that

$$\lim_{n \rightarrow \infty} \|T^n z_n - x_n\| = 0. \tag{3.22}$$

This completes the proof. $\square$

THEOREM 3.3. *Let X be a real uniformly convex Banach space, C a nonempty closed convex subset of X . Let T be uniformly L -Lipschitzian, completely continuous, and an asymptotically quasi-nonexpansive mapping with sequence $\{k_n\}_{n \geq 1}$ such that $\sum_{n=1}^{\infty} k_n < \infty$ and $F(T) \neq \emptyset$. Let $x_0 \in C$ and for each $n \geq 0$,*

$$\begin{aligned}
 z_n &= \alpha_n'' T^n x_n + \beta_n'' x_n + \gamma_n'' u_n, \\
 y_n &= \alpha_n' T^n z_n + \beta_n' x_n + \gamma_n' v_n, \\
 x_{n+1} &= \alpha_n T^n y_n + \beta_n x_n + \gamma_n w_n,
 \end{aligned} \tag{3.23}$$

where $\{u_n\}$, $\{v_n\}$, and $\{w_n\}$ are three bounded sequences in C and $\{\alpha_n\}$, $\{\alpha_n'\}$, $\{\alpha_n''\}$, $\{\beta_n\}$, $\{\beta_n'\}$, $\{\beta_n''\}$, $\{\gamma_n\}$, $\{\gamma_n'\}$, and $\{\gamma_n''\}$ are real sequences in $[0, 1]$ which satisfy the same assumptions as Lemma 3.1 and the additional assumption that $0 \leq \alpha < \alpha_n$, $\beta_n, \alpha_n', \beta_n' \leq \beta < 1$ for some α, β in $(0, 1)$. Then $\{x_n\}$, $\{y_n\}$, and $\{z_n\}$ converge strongly to a fixed point of T .

Proof. It follows from Lemma 3.2 that

$$\lim_{n \rightarrow \infty} \|T^n y_n - x_n\| = 0 = \lim_{n \rightarrow \infty} \|T^n z_n - x_n\| \quad (3.24)$$

and this implies that

$$\|x_{n+1} - x_n\| \leq \alpha_n \|T^n y_n - x_n\| + \gamma_n \|w_n - x_n\| \rightarrow 0 \quad \text{as } n \rightarrow \infty. \quad (3.25)$$

We note that

$$\begin{aligned} \|T^n x_n - x_n\| &\leq \|T^n x_n - T^n y_n\| + \|T^n y_n - x_n\| \leq L \|x_n - y_n\| + \|T^n y_n - x_n\| \\ &\leq \alpha'_n L \|x_n - T^n z_n\| + \gamma'_n L \|v_n - x_n\| + \|T^n y_n - x_n\| \rightarrow 0 \quad \text{as } n \rightarrow \infty, \end{aligned} \quad (3.26)$$

$$\begin{aligned} \|x_n - Tx_n\| &\leq \|x_{n+1} - x_n\| + \|x_{n+1} - T^{n+1}x_{n+1}\| + \|T^{n+1}x_{n+1} - T^{n+1}x_n\| + \|T^{n+1}x_n - Tx_n\| \\ &\leq \|x_{n+1} - x_n\| + \|x_{n+1} - T^{n+1}x_{n+1}\| + (1 + k_{n+1})\|x_{n+1} - x_n\| + L\|T^n x_n - x_n\|. \end{aligned} \quad (3.27)$$

It follows from (3.25), (3.26), and the above inequality that

$$\lim_{n \rightarrow \infty} \|x_n - Tx_n\| = 0. \quad (3.28)$$

By Lemma 3.1, $\{x_n\}$ is bounded. It follows from our assumption that T is completely continuous and that there exists a subsequence $\{Tx_{n_k}\}$ of $\{Tx_n\}$ such that $Tx_{n_k} \rightarrow p \in C$ as $k \rightarrow \infty$. Moreover, by (3.28), we have $\|Tx_{n_k} - x_{n_k}\| \rightarrow 0$ which implies that $x_{n_k} \rightarrow p$ as $k \rightarrow \infty$. By (3.28) again, we have

$$\|p - Tp\| = \lim_{k \rightarrow \infty} \|x_{n_k} - Tx_{n_k}\| = 0. \quad (3.29)$$

This shows that $p \in F(T)$. Furthermore, since $\lim_{n \rightarrow \infty} \|x_n - p\|$ exists, we have $\lim_{n \rightarrow \infty} \|x_n - p\| = 0$, that is, $\{x_n\}$ converges to some fixed point of T . It follows that

$$\begin{aligned} \|y_n - x_n\| &\leq \alpha'_n \|T^n z_n - x_n\| + \gamma'_n \|v_n - x_n\| \rightarrow 0, \\ \|z_n - x_n\| &\leq \alpha''_n \|T^n x_n - x_n\| + \gamma''_n \|u_n - x_n\| \rightarrow 0. \end{aligned} \quad (3.30)$$

Therefore, $\lim_{n \rightarrow \infty} y_n = p = \lim_{n \rightarrow \infty} z_n$. This completes the proof. $\square$

References

- [1] R. Glowinski and P. Le Tallec, *Augmented Lagrangian and Operator-Splitting Methods in Non-linear Mechanics*, SIAM Studies in Applied Mathematics, vol. 9, SIAM, Pennsylvania, 1989.
- [2] K. Goebel and W. A. Kirk, *A fixed point theorem for asymptotically nonexpansive mappings*, Proc. Amer. Math. Soc. **35** (1972), 171–174.
- [3] S. Haubruge, V. H. Nguyen, and J. J. Strodiot, *Convergence analysis and applications of the Glowinski-Le Tallec splitting method for finding a zero of the sum of two maximal monotone operators*, J. Optim. Theory Appl. **97** (1998), no. 3, 645–673.

- [4] Q. Liu, *Iteration sequences for asymptotically quasi-nonexpansive mapping with an error member of uniform convex Banach space*, J. Math. Anal. Appl. **266** (2002), no. 2, 468–471.
- [5] M. A. Noor, *New approximation schemes for general variational inequalities*, J. Math. Anal. Appl. **251** (2000), no. 1, 217–229.
- [6] ———, *Three-step iterative algorithms for multivalued quasi variational inclusions*, J. Math. Anal. Appl. **255** (2001), no. 2, 589–604.
- [7] J. Schu, *Iterative construction of fixed points of strictly pseudocontractive mappings*, Appl. Anal. **40** (1991), no. 2-3, 67–72.
- [8] B. Xu and M. A. Noor, *Fixed-point iterations for asymptotically nonexpansive mappings in Banach spaces*, J. Math. Anal. Appl. **267** (2002), no. 2, 444–453.

Somyot Plubtieng: Department of Mathematics, Faculty of Science, Naresuan University, Phitsanulok 65000, Thailand

E-mail address: somyotp@nu.ac.th

Rabian Wangkeeree: Department of Mathematics, Faculty of Science, Naresuan University, Phitsanulok 65000, Thailand

E-mail address: rabianw@nu.ac.th