EXPANSION OF α -OPEN SETS AND DECOMPOSITION OF α -CONTINUOUS MAPPINGS

M. RAJAMANI AND K. BAGYALAKSHMI

Received 6 November 2004 and in revised form 22 June 2005

We introduce the notions of expansion $\mathcal{A}_\alpha$ of α -open sets and $\mathcal{A}_\alpha$ -expansion α -continuous mappings in topological spaces. The main result of this paper is that a map f is α -continuous if and only if it is $\mathcal{A}_\alpha$ -expansion α -continuous and $\mathcal{B}_\alpha$ -expansion α -continuous, where $\mathcal{A}_\alpha, \mathcal{B}_\alpha$ are two mutually dual expansions.

1. Introduction

In 1965, Njastad [2] introduced the notion of α -sets in topological space. In 1983, Mashhour et al. [1] introduced, with the help of α -sets, a weak form of continuity which they termed as α -continuity. Noiri [3] introduced the same concept, but under the name strong semicontinuity. Noiri [4] defined with the aid of α -sets a new weakened form of continuous mapping called weakly α -continuous mapping. Sen and Bhattacharyya [5] introduced another new weakened form of continuity called weak $^*\alpha$ -continuity and proved that a mapping is α -continuous if and only if it is weakly α -continuous and weak $^*\alpha$ -continuous.

In this paper, we give a general setting for such decompositions of α -continuity by using expansion of α -open sets, whereas in [6], Tong used expansion of open sets to give a general setting for the decomposition of continuous mapping into weakly continuous and weak * continuous mappings.

2. Preliminaries

Throughout this paper, $(X, \tau), (Y, \sigma)$, and so forth (or simply X, Y , etc.) will always denote topological spaces. The family of all α -open sets in X is denoted by τ_α .

We recall the definition of weakly α -continuous and weak $^*\alpha$ -continuous mappings.

Definition 2.1 [1]. A mapping $f : X \rightarrow Y$ is said to be α -continuous if for each open set V in Y , $f^{-1}(V) \in \tau_\alpha$.

Definition 2.2 [3]. A mapping $f : X \rightarrow Y$ is said to be weakly α -continuous if for each x in X and for each open set V in Y containing $f(x)$, there exists a set $U \in \tau_\alpha$ containing x such that $f(U) \subseteq \text{Cl } V$, where $\text{Cl } V$ means the closure of V .

PROPOSITION 2.3 [5]. A mapping $f : X \rightarrow Y$ is weakly α -continuous if and only if $f^{-1}(V) \subseteq \alpha\text{int}[f^{-1}(\text{Cl } V)]$, for every open set V in Y , where $\alpha\text{int}(A)$ means α -interior of A .

Definition 2.4 [5]. A mapping $f : X \rightarrow Y$ is said to be weak $\ast \alpha$ -continuous if and only if for every open set $V \subseteq Y$, $f^{-1}(\text{Fr } V)$ is α -closed in X , where $\text{Fr } V = \text{Cl } V \setminus V$ is the boundary operator for open sets.

3. Decompositions of α -continuity

Definition 3.1. Let (X, τ) be a topological space, let 2^X be the set of all subsets in X . A mapping $\mathcal{A}_\alpha : \tau_\alpha \rightarrow 2^X$ is said to be an expansion on X if $U \subseteq \mathcal{A}_\alpha U$ for each $U \in \tau_\alpha$.

Remark 3.2. If γ_α is the identity expansion, then γ_α is defined by $\gamma_\alpha U = U$. μ_α defined by $\mu_\alpha U = (\alpha\text{int } U \cap U^c)^c$ is an expansion. Cl_α defined by $\text{Cl}_\alpha U = \text{Cl } U$ and $\mathcal{F}_\alpha U$ defined by $\mathcal{F}_\alpha U = (\text{Fr } U)^c$ are expansions.

Definition 3.3 [6]. Let (X, τ) be a topological space. A pair of expansions $\mathcal{A}$ and $\mathcal{B}$ on X is said to be mutually dual if $\mathcal{A}U \cup \mathcal{B}U = U$ for each $U \in \tau$.

Remark 3.4. Let (X, τ) be a topological space. Then Cl_α and $\mathcal{F}_\alpha$ are mutually dual. This follows from [6, Proposition 2].

Example 3.5. Let $X = \{a, b, c\}$ with topologies $\tau = \{\phi, \{a\}, X\}$, $\tau_\alpha = \{\phi, \{a\}, \{a, b\}, \{a, c\}, X\}$. $\mathcal{A}_\alpha(\phi) = \phi$, $\mathcal{A}_\alpha\{a\} = \{a\}$, $\mathcal{A}_\alpha\{a, b\} = \{a, b\}$, $\mathcal{A}_\alpha\{a, c\} = X$, and $\mathcal{A}_\alpha(X) = X$. Then $\mathcal{A}_\alpha$ is an expansion. Let $\mathcal{B}_1(\phi) = X$, $\mathcal{B}_1\{a\} = \{a, c\}$, $\mathcal{B}_1\{a, b\} = X$, $\mathcal{B}_1\{a, c\} = \{a, c\}$, and $\mathcal{B}_1(X) = X$. $\mathcal{B}_2(\phi) = X$, $\mathcal{B}_2\{a\} = \{a, b\}$, $\mathcal{B}_2\{a, b\} = \{a, b\}$, $\mathcal{B}_2\{a, c\} = \{a, c\}$, and $\mathcal{B}_2(X) = X$. Then $\mathcal{B}_1, \mathcal{B}_2$ are both mutually dual to $\mathcal{A}_\alpha$.

PROPOSITION 3.6. Let (X, τ) be a topological space. Then γ_α and μ_α are mutually dual.

Proof.

$$\begin{aligned}
 (\gamma_\alpha U) \cap (\mu_\alpha U) &= U \cap (\alpha\text{int } U \cap U^c)^c \\
 &= (\alpha\text{int } U) \cap (\alpha\text{int } U \cap U^c)^c \\
 &= (\alpha\text{int } U) \cap ((\alpha\text{int } U)^c \cup U) \\
 &= ((\alpha\text{int } U) \cap (\alpha\text{int } U)^c) \cup (\alpha\text{int } U \cap U) \\
 &= \phi \cup U = U.
 \end{aligned} \tag{3.1}$$

□

Definition 3.7. Let (X, τ) and (Y, σ) be two topological spaces and let $\mathcal{A}_\alpha$ be an expansion on Y . Then the mapping $f : X \rightarrow Y$ is said to be $\mathcal{A}_\alpha$ -expansion α -continuous if $f^{-1}(V) \subseteq \alpha\text{int}[f^{-1}(\mathcal{A}_\alpha V)]$, for each $V \in \sigma$.

Remark 3.8. A weakly α -continuous mapping $f : X \rightarrow Y$ can be renamed as Cl_α -expansion α -continuous mapping.

THEOREM 3.9. Let (X, τ) and (Y, σ) be two topological spaces and $\mathcal{A}_\alpha, \mathcal{B}_\alpha$ are two mutually dual expansions on Y . Then the mapping $f : X \rightarrow Y$ is α -continuous if and only if f is $\mathcal{A}_\alpha$ -expansion α -continuous and $\mathcal{B}_\alpha$ -expansion α -continuous.

Proof. Necessity. Suppose that f is α -continuous. Since $\mathcal{A}_\alpha, \mathcal{B}_\alpha$ are mutually dual on Y , $\mathcal{A}_\alpha V \cap \mathcal{B}_\alpha V = V$ for each $V \in \sigma$.

Then

$$\begin{aligned} f^{-1}(V) &= f^{-1}(\mathcal{A}_\alpha V \cap \mathcal{B}_\alpha V) \\ &= f^{-1}(\mathcal{A}_\alpha V) \cap f^{-1}(\mathcal{B}_\alpha V). \end{aligned} \quad (3.2)$$

Since f is α -continuous, $f^{-1}(V) = \alpha \text{ int } f^{-1}(V)$.

Therefore,

$$\begin{aligned} f^{-1}(V) &= \alpha \text{ int } f^{-1}(V) = \alpha \text{ int } (f^{-1})(\mathcal{A}_\alpha V \cap \mathcal{B}_\alpha V) \\ &= \alpha \text{ int } f^{-1}(\mathcal{A}_\alpha V) \cap \alpha \text{ int } f^{-1}(\mathcal{B}_\alpha V). \end{aligned} \quad (3.3)$$

This implies that $f^{-1}(V) \subseteq \alpha \text{ int } f^{-1}(\mathcal{A}_\alpha V)$ and $f^{-1}(V) \subseteq \alpha \text{ int } f^{-1}(\mathcal{B}_\alpha V)$. This shows that f is $\mathcal{A}_\alpha$ -expansion α -continuous and $\mathcal{B}_\alpha$ -expansion α -continuous.

Sufficiency. Since f is $\mathcal{A}_\alpha$ -expansion α -continuous, $f^{-1}(V) \subseteq \alpha \text{ int } f^{-1}(\mathcal{A}_\alpha V)$ for each $V \in \sigma$. Since f is $\mathcal{B}_\alpha$ -expansion α -continuous, $f^{-1}(V) \subseteq \alpha \text{ int } f^{-1}(\mathcal{B}_\alpha V)$ for each $V \in \sigma$. As $\mathcal{A}_\alpha$ and $\mathcal{B}_\alpha$ are two mutually dual expansions on Y , $\mathcal{A}_\alpha V \cap \mathcal{B}_\alpha V = V$,

$$\begin{aligned} f^{-1}(V) &= f^{-1}(\mathcal{A}_\alpha V \cap \mathcal{B}_\alpha V) = f^{-1}(\mathcal{A}_\alpha V) \cap f^{-1}(\mathcal{B}_\alpha V), \\ \alpha \text{ int } f^{-1}(V) &= \alpha \text{ int } f^{-1}(\mathcal{A}_\alpha V) \cap \alpha \text{ int } f^{-1}(\mathcal{B}_\alpha V) \supseteq f^{-1}(V) \cap f^{-1}(V) = f^{-1}(V). \end{aligned} \quad (3.4)$$

This implies that $f^{-1}(V) \subseteq \alpha \text{ int } f^{-1}(V)$. Always, $\alpha \text{ int } f^{-1}(V) \subseteq f^{-1}(V)$. So $f^{-1}(V) = \alpha \text{ int } f^{-1}(V)$. Therefore, $f^{-1}(V)$ is an α -open set in X for each $V \in \sigma$. Hence f is α -continuous. $\square$

Definition 3.10. Let (X, τ) and (Y, σ) be two topological spaces, $\mathcal{B}_\alpha$ an expansion on Y . Then a mapping $f : X \rightarrow Y$ is said to be α -closed $\mathcal{B}_\alpha$ -continuous if $f^{-1}((\mathcal{B}_\alpha V)^c)$ is an α -closed set in X for each $V \in \sigma$.

Remark 3.11. A weak $\ast$ - α -continuous mapping can be renamed as α -closed $\mathcal{F}_\alpha$ -continuous mapping since $(\mathcal{F}_\alpha V)^c = (\text{Fr } V)^c = \text{Fr } V$.

PROPOSITION 3.12. An α -closed $\mathcal{B}_\alpha$ -continuous mapping is $\mathcal{B}_\alpha$ -expansion α -continuous.

Proof. First, we prove that $(f^{-1}((\mathcal{B}_\alpha V)^c))^c = f^{-1}(\mathcal{B}_\alpha V)$.

Let $x \in (f^{-1}((\mathcal{B}_\alpha V)^c))^c$. Then $x \notin f^{-1}(\mathcal{B}_\alpha V)^c$. Hence $f(x) \notin (\mathcal{B}_\alpha V)^c$, $f(x) \in \mathcal{B}_\alpha V$, and $x \in f^{-1}(\mathcal{B}_\alpha V)$.

Conversely, if $x \in f^{-1}(\mathcal{B}_\alpha V)$, then $f(x) \in \mathcal{B}_\alpha V$. Hence $f(x) \notin (\mathcal{B}_\alpha V)^c$, $x \notin f^{-1}(\mathcal{B}_\alpha V)^c$, $x \in (f^{-1}((\mathcal{B}_\alpha V)^c))^c$. Therefore, $(f^{-1}((\mathcal{B}_\alpha V)^c))^c = f^{-1}(\mathcal{B}_\alpha V)$.

Since $f^{-1}((\mathcal{B}_\alpha V)^c)$ is an α -closed set in X , $(f^{-1}((\mathcal{B}_\alpha V)^c))^c$ is an α -open set in X . Hence $f^{-1}(\mathcal{B}_\alpha V)$ is an α -open in X and $f^{-1}(\mathcal{B}_\alpha V) = \alpha \text{ int } f^{-1}(\mathcal{B}_\alpha V)$.

Since $\mathcal{B}_\alpha$ is an expansion on Y , $V \subseteq \mathcal{B}_\alpha V$, we have $f^{-1}(V) \subseteq f^{-1}(\mathcal{B}_\alpha V) = \alpha \text{ int } f^{-1}(\mathcal{B}_\alpha V)$. Therefore, f is $\mathcal{B}_\alpha$ -expansion α -continuous. $\square$

By Theorem 3.9 and Proposition 3.12, we have the following corollary.

COROLLARY 3.13. *Let (X, τ) and (Y, σ) be two topological spaces and $\mathcal{A}_\alpha, \mathcal{B}_\alpha$ are two mutually dual expansions on Y . Then a mapping $f : X \rightarrow Y$ is α -continuous if and only if f is $\mathcal{A}_\alpha$ -expansion α -continuous, and α -closed $\mathcal{B}_\alpha$ -continuous.*

By Remarks 3.8, 3.11, and by the above corollary, we have the following corollary.

COROLLARY 3.14 [5]. *A mapping is α -continuous if and only if it is weakly α -continuous and weak $*$ - α -continuous.*

References

- [1] A. S. Mashhour, I. A. Hasanein, and S. N. El-Deeb, α -continuous and α -open mappings, *Acta Math. Hungar.* **41** (1983), no. 3-4, 213–218.
- [2] O. Njåstad, *On some classes of nearly open sets*, *Pacific J. Math.* **15** (1965), no. 3, 961–970.
- [3] T. Noiri, *A function which preserves connected spaces*, *Časopis Pěst. Mat.* **107** (1982), no. 4, 393–396, 428.
- [4] ———, *Weakly α -continuous functions*, *Int. J. Math. Math. Sci.* **10** (1987), no. 3, 483–490.
- [5] A. K. Sen and P. Bhattacharyya, *On weakly α -continuous functions*, *Tamkang J. Math.* **24** (1993), no. 4, 445–460.
- [6] J. C. Tong, *Expansion of open sets and decomposition of continuous mappings*, *Rend. Circ. Mat. Palermo (2)* **43** (1994), no. 2, 303–308.

M. Rajamani: Post Graduate and Research Department of Mathematics, N.G.M. College (Autonomous), Bharathiar University, Coimbatore, Pollachi-642 001, Tamil Nadu, India
E-mail address: rajkarthy@yahoo.com

K. Bagyalakshmi: Post Graduate and Research Department of Mathematics, N.G.M. College (Autonomous), Bharathiar University, Coimbatore, Pollachi-642 001, Tamil Nadu, India
E-mail address: rajkarthy@yahoo.com