LERAY-SCHAUDER RESULTS FOR MULTIVALUED NONLINEAR CONTRACTIONS DEFINED ON CLOSED SUBSETS OF A FRÉCHET SPACE

RAVI P. AGARWAL AND DONAL O'REGAN

Received 9 December 2005; Revised 21 February 2006; Accepted 25 April 2006

New Leray-Schauder results are presented for multivalued contractions defined on subsets of a Fréchet space E . The proof relies on fixed point results in Banach spaces and on viewing E as the projective limit of a sequence of Banach spaces.

Copyright © 2006 Hindawi Publishing Corporation. All rights reserved.

1. Introduction

In this paper, we present new fixed point results for nonlinear contractions (both single and multivalued) defined on subsets X (which may have empty interior) of a Fréchet space E . Some results for single-valued maps were presented in [2, 3] and the approach in these papers was based on constructing a specific map F_n (for each $n \in \mathbb{N} = \{1, 2, \dots\}$) whose fixed points converge to a fixed point of the original operator F . In the approach in this paper, the maps $\{F_n\}_{n \in \mathbb{N}}$ only need to satisfy a closure property and are specified in a completely different way. The advantage of this approach is that multivalued maps can also be discussed. Our theory is based on results in Banach spaces and on viewing a Fréchet space E as a projective limit of a sequence of Banach spaces $\{E_n\}_{n \in \mathbb{N}}$.

For the remainder of this section, we present some definitions and some known facts. Let (X, d) be a metric space and S a nonempty subset of X . For $x \in X$, let $d(x, S) = \inf_{y \in S} d(x, y)$. Also $\text{diam } S = \sup\{d(x, y) : x, y \in S\}$. We let $B(x, r)$ denote the open ball in X centered at x of radius r and by $B(S, r)$ we denote $\bigcup_{x \in S} B(x, r)$. For two nonempty subsets S_1 and S_2 of X , we define the generalized Hausdorff distance H to be

$$H(S_1, S_2) = \inf \{ \epsilon > 0 : S_1 \subseteq B(S_2, \epsilon), S_2 \subseteq B(S_1, \epsilon) \}. \quad (1.1)$$

Now suppose $G : S \rightarrow 2^X$; here 2^X denotes the family of nonempty subsets of X . Then G is said to be hemicompact if each sequence $\{x_n\}_{n \in \mathbb{N}}$ in S has a convergent subsequence whenever $d(x_n, G(x_n)) \rightarrow 0$ as $n \rightarrow \infty$.

We now recall a result from the literature.

2 Multivalued nonlinear contractions

THEOREM 1.1. *Let (X, d) be a complete metric space, $\mathbb{C} \subseteq X$ closed, and $F : \mathbb{C} \rightarrow X$ with $F(\mathbb{C})$ bounded (i.e., there exists $M > 0$ with $d(z, w) \leq M$ for $z, w \in F(\mathbb{C})$). Suppose the following condition is satisfied:*

$$\begin{aligned} & \text{there exists a continuous nondecreasing function} \\ & \phi : [0, \infty) \longrightarrow [0, \infty) \text{ satisfying } \phi(z) < z \text{ for } z > 0 \\ & \text{such that } d(Fx, Fy) \leq \phi(d(x, y)) \text{ for } x, y \in \mathbb{C}. \end{aligned} \quad (1.2)$$

Then F is hemicompact.

Now let I be a directed set with order $\leq$ and let $\{E_\alpha\}_{\alpha \in I}$ be a family of locally convex spaces. For each $\alpha \in I$, $\beta \in I$ for which $\alpha \leq \beta$, let $\pi_{\alpha, \beta} : E_\beta \rightarrow E_\alpha$ be a continuous map. Then the set

$$\left\{ x = (x_\alpha) \in \prod_{\alpha \in I} E_\alpha : x_\alpha = \pi_{\alpha, \beta}(x_\beta) \quad \forall \alpha, \beta \in I, \alpha \leq \beta \right\} \quad (1.3)$$

is a closed subset of $\prod_{\alpha \in I} E_\alpha$ and is called the projective limit of $\{E_\alpha\}_{\alpha \in I}$ and is denoted by $\lim_{\leftarrow} E_\alpha$ (or $\lim_{\leftarrow} \{E_\alpha, \pi_{\alpha, \beta}\}$ or the generalized intersection [5, page 439] $\bigcap_{\alpha \in I} E_\alpha$).

Existence in Section 2 is based on the following fixed point results in the literature [1, 6].

THEOREM 1.2 [6, Theorem 3.9]. *Let U be an open subset in a Banach space $(X, \|\cdot\|)$ and $F : \overline{U} \rightarrow X$. Assume $0 \in U$ and suppose there exists a continuous nondecreasing function $\phi : [0, \infty) \rightarrow [0, \infty)$ satisfying $\phi(z) < z$ for $z > 0$ such that $\|Fx - Fy\| \leq \phi(\|x - y\|)$ for all $x, y \in \overline{U}$. In addition, assume $F(\overline{U})$ is bounded and $x \neq \lambda Fx$ for $x \in \partial U$ and $\lambda \in (0, 1)$. Then F has a fixed point in $\overline{U}$.*

THEOREM 1.3 [1, Theorem 2.3 (and Remark 2.1)]. *Let U be an open subset in a Banach space $(X, \|\cdot\|)$ and $F : \overline{U} \rightarrow \mathbb{C}(X)$ a closed map (i.e., has closed graph); here $\mathbb{C}(X)$ denotes the family of nonempty closed subsets of X . Assume $0 \in U$ and suppose there exists a continuous strictly increasing function $\phi : [0, \infty) \rightarrow [0, \infty)$ satisfying $\phi(z) < z$ for $z > 0$ such that $H(Fx, Fy) \leq \phi(\|x - y\|)$ for all $x, y \in \overline{U}$. In addition, assume the following conditions hold:*

$$\Phi : [0, \infty) \longrightarrow [0, \infty), \text{ given by } \Phi(x) = x - \phi(x), \text{ is strictly increasing,} \quad (1.4)$$

$$\Phi^{-1}(a) + \Phi^{-1}(b) \leq \Phi^{-1}(a + b) \quad \text{for } a, b \geq 0, \quad (1.5)$$

$$\sum_{i=0}^{\infty} \phi^i(t) < \infty \quad \text{for } t > 0, \quad (1.6)$$

$$\sum_{i=1}^{\infty} \phi^i(x - \phi(x)) \leq \phi(x) \quad \text{for } x > 0, \quad (1.7)$$

$$F(\overline{U}) \text{ is bounded,} \quad (1.8)$$

$$x \notin \lambda Fx \quad \text{for } x \in \partial U, \lambda \in (0, 1). \quad (1.9)$$

Then F has a fixed point in $\overline{U}$.

Remark 1.4. In fact, the assumption that F is closed can be removed in Theorem 1.3. In [1, Theorem 2.3], we assume a more general contractive condition and the map $G: \overline{U} \times [0, 1] \rightarrow \mathbb{C}(X)$ (given by $G(x, \lambda) = \lambda Fx$ in our situation) was assumed to be closed in order to guarantee that if $\{x_n\}_1^\infty \subseteq \overline{U}$, $\{\lambda\}_1^\infty \subseteq [0, 1]$ with $x_n \in G(x_n, \lambda_n)$ and $(x_n, \lambda_n) \rightarrow (x, \lambda)$, then $x \in G(x, \lambda)$. However, this is automatically true in Theorem 1.3 since the contractive condition and (1.8) guarantee that G is continuous in the Hausdorff metric and as a result,

$$\text{dist}(x, G(x, \lambda)) \leq d(x, x_n) + H(G(x_n, \lambda_n), G(x, \lambda)). \quad (1.10)$$

Remark 1.5. If $\phi(t) = kt$, $0 \leq k < 1$, then trivially (1.2)–(1.7) hold.

2. Fixed point theory in Fréchet spaces

Let $E = (E, \{\|\cdot\|_n\}_{n \in \mathbb{N}})$ be a Fréchet space with the topology generated by a family of seminorms $\{\|\cdot\|_n : n \in \mathbb{N}\}$. We assume that the family of seminorms satisfies

$$\|x\|_1 \leq \|x\|_2 \leq \|x\|_3 \leq \cdots \quad \text{for every } x \in E. \quad (2.1)$$

A subset X of E is bounded if for every $n \in \mathbb{N}$ there exists $r_n > 0$ such that $\|x\|_n \leq r_n$ for all $x \in X$. To E we associate a sequence of Banach spaces $\{(\mathbf{E}_n, \|\cdot\|_n)\}$ described as follows. For every $n \in \mathbb{N}$, we consider the equivalence relation $\sim_n$ defined by

$$x \sim_n y \quad \text{iff } \|x - y\|_n = 0. \quad (2.2)$$

We denote by $\mathbf{E}^n = (E/\sim_n, \|\cdot\|_n)$ the quotient space, and by $(\mathbf{E}_n, \|\cdot\|_n)$ the completion of $\mathbf{E}^n$ with respect to $\|\cdot\|_n$ (the norm on $\mathbf{E}^n$ induced by $\|\cdot\|_n$ and its extension to $\mathbf{E}_n$ are still denoted by $\|\cdot\|_n$). This construction defines a continuous map $\mu_n : E \rightarrow \mathbf{E}_n$. Now since (2.1) is satisfied, the seminorm $\|\cdot\|_n$ induces a seminorm on $\mathbf{E}_m$ for every $m \geq n$ (again this seminorm is denoted by $\|\cdot\|_n$). Also (2.2) defines an equivalence relation on $\mathbf{E}_m$ from which we obtain a continuous map $\mu_{n,m} : \mathbf{E}_m \rightarrow \mathbf{E}_n$ since $\mathbf{E}_m/\sim_n$ can be regarded as a subset of $\mathbf{E}_n$. We now assume the following condition holds: for each $n \in \mathbb{N}$, there exists a Banach space $(E_n, \|\cdot\|_n)$ and an isomorphism (between normed spaces) $j_n : \mathbf{E}_n \rightarrow E_n$.

Remark 2.1. (i) For convenience, the norm on E_n is denoted by $\|\cdot\|_n$.

(ii) Usually in applications, $\mathbf{E}_n = \mathbf{E}^n$ for each $n \in \mathbb{N}$.

(iii) Note that if $x \in \mathbf{E}_n$ (or $\mathbf{E}^n$), then $x \in E$. However, if $x \in E_n$, then x is not necessarily in E and in fact, E_n is easier to use in applications (even though E_n is isomorphic to $\mathbf{E}_n$). For example, if $E = \mathbb{C}[0, \infty)$, then $\mathbf{E}^n$ consists of the class of functions in E which coincide on the interval $[0, n]$ and $E_n = \mathbb{C}[0, n]$.

Finally, we assume

$$E_1 \supseteq E_2 \supseteq \cdots \quad \text{and for each } n \in \mathbb{N}, \quad \|x\|_n \leq \|x\|_{n+1} \quad \forall x \in E_{n+1}. \quad (2.3)$$

Let $\lim_- E_n$ (or $\bigcap_1^\infty E_n$ where $\bigcap_1^\infty$ is the generalized intersection [5]) denote the projective limit of $\{E_n\}_{n \in \mathbb{N}}$ (note that $\pi_{n,m} = j_n \mu_{n,m} j_m^{-1} : E_m \rightarrow E_n$ for $m \geq n$) and note that $\lim_- E_n \cong E$, so for convenience, we write $E = \lim_- E_n$.

4 Multivalued nonlinear contractions

For each $X \subseteq E$ and each $n \in \mathbb{N}$, we set $X_n = j_n \mu_n(X)$ and we let $\overline{X_n}$ and ∂X_n denote, respectively, the closure and the boundary of X_n with respect to $|\cdot|_n$ in E_n . Also the pseudo-interior of X is defined by [4]

$$\text{pseudo-intt}(X) = \{x \in X : j_n \mu_n(x) \in \overline{X_n} \setminus \partial X_n \text{ for every } n \in \mathbb{N}\}. \quad (2.4)$$

Also, here H_n and diam_n denote the Hausdorff metric and the diameter induced by $|\cdot|_n$ on E_n .

We begin with single-valued maps and present two results. The first was motivated by Volterra type operators.

THEOREM 2.2. *Let E and E_n be as described above and let $F : X \rightarrow E$ with $X \subseteq E$ and for each $n \in \mathbb{N}$ assume that $F : \overline{X_n} \rightarrow E_n$. Suppose the following conditions are satisfied:*

- (a) $0 \in \text{pseudo-intt}(X)$,
- (b) for each $n \in \mathbb{N}$, $F(\overline{X_n})$ is bounded,
- (c) for each $n \in \mathbb{N}$, $F : \overline{X_n} \rightarrow E_n$ and there exists a continuous nondecreasing function $\phi_n : [0, \infty) \rightarrow [0, \infty)$ satisfying $\phi_n(z) < z$ for $z > 0$ such that $|Fx - Fy|_n \leq \phi_n(|x - y|_n)$ for all $x, y \in \overline{X_n}$ for each $n \in \mathbb{N}$, $y \neq \lambda Fy$, in E_n for all $\lambda \in (0, 1)$, $y \in \partial X_n$,
- (d) for each $n \in \{2, 3, \dots\}$, if $y \in \overline{X_n}$ solves $y = Fy$ in E_n , then $y \in \overline{X_k}$ for $k \in \{1, \dots, n-1\}$.

Then F has a fixed point in E .

Remark 2.3. If $F(X)$ is bounded, then clearly Theorem 2.2(b) holds.

Proof. Fix $n \in \mathbb{N}$. From Theorem 1.2, there exists $y_n \in \overline{X_n}$ with $y_n = Fy_n$ (note that $0 \in \overline{X_n} \setminus \partial X_n$ and $F(\overline{X_n})$ is bounded). Let us look at $\{y_n\}_{n \in \mathbb{N}}$. Notice that $y_1 \in \overline{X_1}$ and $y_k \in \overline{X_1}$ for $k \in \mathbb{N} \setminus \{1\}$ from Theorem 2.2(d). As a result, $y_n \in \overline{X_1}$ for $n \in \mathbb{N}$, $y_n = Fy_n$ in E_n together with Theorem 1.1 implies there is a subsequence $\mathbb{N}_1^*$ of $\mathbb{N}$ and a $z_1 \in \overline{X_1}$ with $y_n \rightarrow z_1$ in E_1 as $n \rightarrow \infty$ in $\mathbb{N}_1^*$. Let $\mathbb{N}_1 = \mathbb{N}_1^* \setminus \{1\}$. Now $y_n \in \overline{X_2}$ for $n \in \mathbb{N}_1$ together with Theorem 1.1 guarantees that there exists a subsequence $\mathbb{N}_2^*$ of $\mathbb{N}_1$ and a $z_2 \in \overline{X_2}$ with $y_n \rightarrow z_2$ in E_2 as $n \rightarrow \infty$ in $\mathbb{N}_2^*$. Note from (2.3) that $z_2 = z_1$ in E_1 since $\mathbb{N}_2^* \subseteq \mathbb{N}_1$. Let $\mathbb{N}_2 = \mathbb{N}_2^* \setminus \{2\}$. Proceed inductively to obtain subsequences of integers

$$\begin{aligned} \mathbb{N}_1^* \supseteq \mathbb{N}_2^* \supseteq \dots, \\ \mathbb{N}_k^* \subseteq \{k, k+1, \dots\}, \end{aligned} \quad (2.5)$$

and $z_k \in \overline{X_k}$ with $y_n \rightarrow z_k$ in E_k as $n \rightarrow \infty$ in $\mathbb{N}_k^*$. Note that $z_{k+1} = z_k$ in E_k for $k \in \{1, 2, \dots\}$. Also let $\mathbb{N}_k = \mathbb{N}_k^* \setminus \{k\}$.

Fix $k \in \mathbb{N}$. Let $y = z_k$ in E_k . Notice that y is well defined and $y \in \lim_{-} E_n = E$. Now $y_n = Fy_n$ in E_n for $n \in \mathbb{N}_k$ and $y_n \rightarrow y$ in E_k as $n \rightarrow \infty$ in $\mathbb{N}_k$ (since $y = z_k$ in E_k) together with the fact that $F : \overline{X_k} \rightarrow E_k$ is continuous (note that $y_n \in \overline{X_k}$ for $n \in \mathbb{N}_k$) implies $y = Fy$ in E_k . We can do this for each $k \in \mathbb{N}$, so $y = Fy$ in E . $\square$

Our next result was motivated by contractions considered in [3]. In this case, the map F_n will be related to F by the closure property Theorem 2.4(f).

THEOREM 2.4. *Let E and E_n be as described in the beginning of Section 2 and let $F : X \rightarrow E$ with $X \subseteq E$. Also for each $n \in \mathbb{N}$ assume there exists $F_n : \overline{X_n} \rightarrow E_n$. Suppose the following conditions are satisfied:*

- (a) $0 \in \text{pseudo-intt}(X)$,
- (b) $\overline{X_1} \supseteq \overline{X_2} \supseteq \dots$,
- (c) *for each $n \in \mathbb{N}$, $F_n(\overline{X_n})$ is bounded, for each $n \in \mathbb{N}$, $F_n : \overline{X_n} \rightarrow E_n$ and there exists a continuous nondecreasing function $\phi_n : [0, \infty) \rightarrow [0, \infty)$ satisfying $\phi_n(z) < z$ for $z > 0$ such that $|F_n x - F_n y|_n \leq \phi_n(|x - y|_n)$ for all $x, y \in \overline{X_n}$ for each $n \in \mathbb{N}$, $y \neq \lambda F_n y$ in E_n for all $\lambda \in (0, 1)$, $y \in \partial X_n$,*
- (d) *for each $n \in \mathbb{N}$, the map $\mathcal{H}_n : \overline{X_n} \rightarrow 2^{E_n}$ given by $\mathcal{H}_n(y) = \bigcup_{m=n}^{\infty} F_m(y)$ (see Remark 2.5) satisfies $H_n(\mathcal{H}_n(x), \mathcal{H}_n(y)) \leq \psi_n(|x - y|_n)$ for all $x, y \in \overline{X_n}$; here $\psi_n : [0, \infty) \rightarrow [0, \infty)$ is continuous, $\psi_n(z) < z$ for $z > 0$ with the map $\Psi_n : [0, \infty) \rightarrow [0, \infty)$, defined by $\Psi_n(x) = x - \psi_n(x)$, strictly increasing,*
- (e) *for each $k \in \mathbb{N}$, for every $\epsilon > 0$, and sequence $\{x_n\}_{n \in S}$, $S = \{k, k+1, k+2, \dots\}$, with $x_n \in \overline{X_n}$ and $x_n \in \mathcal{H}_n x_n$ in E_n , there exists $n_k \in S$ such that $\text{diam}_k(\mathcal{H}_k x_n) < \epsilon$ for each $n \in S$ with $n \geq n_k$,*
- (f) *if there exists $w \in E$ and a sequence $\{y_n\}_{n \in \mathbb{N}}$ with $y_n \in \overline{X_n}$ and $y_n = F_n y_n$ in E_n such that for every $k \in \mathbb{N}$ with $y_n \rightarrow w$ in E_k as $n \rightarrow \infty$ in $S = \{k+1, k+2, \dots\}$, then $w = Fw$ in E .*

Then F has a fixed point in E .

Remark 2.5. The definition of $\mathcal{H}_n$ in Theorem 2.4(d) is as follows. If $y \in \overline{X_n}$ and $y \notin \overline{X_{n+1}}$, then $\mathcal{H}_n(y) = F_n(y)$, whereas if $y \in \overline{X_{n+1}}$ and $y \notin \overline{X_{n+2}}$, then $\mathcal{H}_n(y) = F_n(y) \cup F_{n+1}(y)$, and so on.

Proof. Fix $n \in \mathbb{N}$. From Theorem 1.2 there exists $y_n \in \overline{X_n}$ with $y_n = F_n y_n$ in E_n . Let us look at $\{y_n\}_{n \in \mathbb{N}}$. From Theorem 2.4(b) we know that $y_n \in \overline{X_1}$ for $n \in \mathbb{N}$. Note as well that $y_n \in \mathcal{H}_1 y_n$ for $n \in \mathbb{N}$ since $|x|_1 \leq |x|_n$ for all $x \in E_n$ and $y_n = F_n y_n$ in E_n . We claim

$$\exists z_1 \in E_1 \quad \text{with } y_n \longrightarrow z_1 \text{ in } E_1, \quad n \longrightarrow \infty \text{ in } \mathbb{N}. \quad (2.6)$$

To see this, let $\epsilon > 0$ be given. Let $m, n \in \mathbb{N}$. It is easy to see, since $y_n \in \mathcal{H}_1 y_n$ and $y_m \in \mathcal{H}_1 y_m$, that

$$|y_n - y_m|_1 \leq H_1(\mathcal{H}_1 y_n, \mathcal{H}_1 y_m) + \text{diam}_1(\mathcal{H}_1 y_n) + \text{diam}_1(\mathcal{H}_1 y_m), \quad (2.7)$$

so Theorem 2.4(d) yields

$$|y_n - y_m|_1 \leq \Psi_1^{-1}(\text{diam}_1(\mathcal{H}_1 y_n) + \text{diam}_1(\mathcal{H}_1 y_m)). \quad (2.8)$$

Now Theorem 2.4(e) guarantees that there exists $n_1 \in \mathbb{N}$ such that

$$|y_n - y_m|_1 \leq \Psi_1^{-1}(2\epsilon) \quad \text{for } m, n \geq n_1. \quad (2.9)$$

Consequently, $\{y_n\}_{n \in \mathbb{N}}$ is Cauchy, so (2.6) holds. Let $\mathbb{N}_1 = \mathbb{N} \setminus \{1\}$.

6 Multivalued nonlinear contractions

Now $y_n \in \mathcal{H}_2 y_n$ for $n \in \mathbb{N}_1$. Let $m, n \in \mathbb{N}_1$ and since $y_n \in \mathcal{H}_2 y_n$ and $y_m \in \mathcal{H}_2 y_m$ we have

$$\|y_n - y_m\|_2 \leq \Psi_2^{-1}(\text{diam}_2(\mathcal{H}_2 y_n) + \text{diam}_2(\mathcal{H}_2 y_m)). \quad (2.10)$$

This together with Theorem 2.4(e) guarantees that $\{y_n\}_{n \in \mathbb{N}_1}$ is Cauchy, so there exists a $z_2 \in E_2$ with $y_n \rightarrow z_2$ in E_2 as $n \rightarrow \infty$ in $\mathbb{N}_1$. Note that $z_2 = z_1$ in E_1 since $\mathbb{N}_1 \subseteq \mathbb{N}$. Let $\mathbb{N}_2 = \mathbb{N}_1 \setminus \{2\}$. Proceed inductively to obtain $z_k \in E_k$ with $y_n \rightarrow z_k$ in E_k as $n \rightarrow \infty$ in $\mathbb{N}_{k-1} = \{k, k+1, \dots\}$. Note that $z_{k+1} = z_k$ in E_k for $k \in \mathbb{N}$. Also let $\mathbb{N}_k = \mathbb{N}_{k-1} \setminus \{k\}$.

Fix $k \in \mathbb{N}$. Let $y = z_k$ in E_k . Notice that y is well defined and $y \in \lim_{\leftarrow} E_n = E$. Now $y_n = F_n y_n$ in E_n for $n \in \mathbb{N}_k$ and $y_n \rightarrow y$ in E_k as $n \rightarrow \infty$ in $\mathbb{N}_k$ (since $y = z_k$ in E_k) together with Theorem 2.4(f) implies $y = Fy$ in E . $\square$

Our next two results are for multivalued maps.

THEOREM 2.6. *Let E and E_n be as described above and let $F : X \rightarrow 2^E$ with $X \subseteq E$ and for each $n \in \mathbb{N}$, assume $F : \overline{X_n} \rightarrow \mathbb{C}(E_n)$. Suppose the following conditions are satisfied:*

- (a) $0 \in \text{pseudo-intt}(X)$,
- (b) for each $n \in \mathbb{N}$, $F(\overline{X_n})$ is bounded,
- (c) for each $n \in \mathbb{N}$, $F : \overline{X_n} \rightarrow \mathbb{C}(E_n)$, and there exists a continuous strictly increasing function $\phi_n : [0, \infty) \rightarrow [0, \infty)$ satisfying $\phi_n(z) < z$ for $z > 0$ such that $H_n(Fx, Fy) \leq \phi_n(\|x - y\|_n)$ for all $x, y \in \overline{X_n}$,
- (d) for each $n \in \mathbb{N}$, the map $\Phi_n : [0, \infty) \rightarrow [0, \infty)$ given by $\Phi_n(x) = x - \phi_n(x)$ is strictly increasing, $\Phi_n^{-1}(a) + \Phi_n^{-1}(b) \leq \Phi_n^{-1}(a + b)$ for $a, b \geq 0$, with $\sum_{i=0}^{\infty} \phi_n^i(t) < \infty$ for $t > 0$ and $\sum_{i=1}^{\infty} \phi_n^i(x - \phi(x)) \leq \phi_n(x)$ for $x > 0$,
- (e) for each $n \in \mathbb{N}$, $y \notin \lambda Fy$ in E_n for all $\lambda \in (0, 1)$, $y \in \partial X_n$,
- (f) for each $n \in \{2, 3, \dots\}$, if $y \in \overline{X_n}$ solves $y \in Fy$ in E_n , then $y \in \overline{X_k}$ for $k \in \{1, \dots, n-1\}$,
- (g) for each $k \in \mathbb{N}$, for every $\epsilon > 0$ and sequence $\{x_n\}_{n \in \mathbb{N}}$, $S = \{k, k+1, k+2, \dots\}$, with $x_n \in \overline{X_n}$ and $x_n \in Fx_n$ in E_n there exists $n_k \in S$ such that $\text{diam}_k(Fx_n) < \epsilon$ for each $n \in S$ with $n \geq n_k$.

Then F has a fixed point in E .

Proof. Fix $n \in \mathbb{N}$. From Theorem 1.3 (and Remark 1.4) there exists $y_n \in \overline{X_n}$ with $y_n \in Fy_n$ in E_n . Let us look at $\{y_n\}_{n \in \mathbb{N}}$. Notice that $y_n \in \overline{X_1}$ for $n \in \mathbb{N}$ from Theorem 2.6(f). Let $\epsilon > 0$ be given and $m, n \in \mathbb{N}$. Now since $y_n \in Fy_n$ and $y_m \in Fy_m$, we have

$$\|y_n - y_m\|_1 \leq H_1(Fy_n, Fy_m) + \text{diam}_1(Fy_n) + \text{diam}_1(Fy_m) \quad (2.11)$$

so

$$\|y_n - y_m\|_1 \leq \Phi_1^{-1}(\text{diam}_1(Fy_n) + \text{diam}_1(Fy_m)). \quad (2.12)$$

This, together with Theorem 2.6(g), guarantees that $\{y_n\}_{n \in \mathbb{N}}$ is Cauchy, so there exists a $z_1 \in E_1$ with $y_n \rightarrow z_1$ in E_1 as $n \rightarrow \infty$ in $\mathbb{N}$. Let $\mathbb{N}_1 = \mathbb{N} \setminus \{1\}$. Proceed inductively to obtain $z_k \in E_k$ with $y_n \rightarrow z_k$ in E_k as $n \rightarrow \infty$ in $\mathbb{N}_{k-1} = \{k, k+1, \dots\}$. Note that $z_{k+1} = z_k$ in E_k for $k \in \mathbb{N}$. Also let $\mathbb{N}_k = \mathbb{N}_{k-1} \setminus \{k\}$.

Fix $k \in \mathbb{N}$. Let $y = z_k$ in E_k . Notice that $y_n \in Fy_n$ in E_n for $n \in \mathbb{N}_k$ and $y_n \rightarrow y$ in E_k as $n \rightarrow \infty$ in $\mathbb{N}_k$ together with Remark 1.4 (note that $F : \overline{X_k} \rightarrow \mathbb{C}(E_k)$) implies $y \in Fy$ in E_k . We can do this for each $k \in \mathbb{N}$, so $y \in Fy$ in E . $\square$

THEOREM 2.7. *Let E and E_n be as described in the beginning of Section 2 and let $F : X \rightarrow 2^E$ with $X \subseteq E$. Also for each $n \in \mathbb{N}$ assume there exists $F_n : \overline{X_n} \rightarrow \mathbb{C}(E_n)$. Suppose the following conditions are satisfied:*

- (a) $0 \in \text{pseudo-intt}(X)$,
- (b) $\overline{X_1} \supseteq \overline{X_2} \supseteq \dots$,
- (c) for each $n \in \mathbb{N}$, $F_n(\overline{X_n})$ is bounded,
- (d) for each $n \in \mathbb{N}$, $F_n : \overline{X_n} \rightarrow \mathbb{C}(E_n)$ and there exists a continuous strictly increasing function $\phi_n : [0, \infty) \rightarrow [0, \infty)$ satisfying $\phi_n(z) < z$ for $z > 0$ such that $H_n(F_n x, F_n y) \leq \phi_n(|x - y|_n)$ for all $x, y \in \overline{X_n}$,
- (e) for each $n \in \mathbb{N}$, the map $\Phi_n : [0, \infty) \rightarrow [0, \infty)$ given by $\Phi_n(x) = x - \phi_n(x)$ is strictly increasing, $\Phi_n^{-1}(a) + \Phi_n^{-1}(b) \leq \Phi_n^{-1}(a + b)$ for $a, b \geq 0$, with $\sum_{i=0}^{\infty} \phi_n^i(t) < \infty$ for $t > 0$ and $\sum_{i=1}^{\infty} \phi_n^i(x - \phi(x)) \leq \phi_n(x)$ for $x > 0$,
- (f) for each $n \in \mathbb{N}$, $y \notin \lambda F_n y$ in E_n for all $\lambda \in (0, 1)$ and $y \in \partial X_n$,
- (g) for each $n \in \mathbb{N}$, the map $\mathcal{H}_n : \overline{X_n} \rightarrow 2^{E_n}$ given by $\mathcal{H}_n(y) = \bigcup_{m=n}^{\infty} F_m(y)$ satisfies $H_n(\mathcal{H}_n(x), \mathcal{H}_n(y)) \leq \psi_n(|x - y|_n)$ for all $x, y \in \overline{X_n}$; here $\psi_n : [0, \infty) \rightarrow [0, \infty)$ is continuous, $\psi_n(z) < z$ for $z > 0$ with the map $\Psi_n : [0, \infty) \rightarrow [0, \infty)$ defined by $\Psi_n(x) = x - \psi_n(x)$ is strictly increasing,
- (h) for each $k \in \mathbb{N}$, for every $\epsilon > 0$ and sequence $\{x_n\}_{n \in S}$, $S = \{k, k+1, k+2, \dots\}$, with $x_n \in \overline{X_n}$ and $x_n \in \mathcal{H}_n x_n$ in E_n there exists $n_k \in S$ such that $\text{diam}_k(\mathcal{H}_k x_n) < \epsilon$ for each $n \in S$ with $n \geq n_k$,
- (i) if there exists a $w \in E$ and a sequence $\{y_n\}_{n \in \mathbb{N}}$ with $y_n \in \overline{X_n}$ and $y_n \in F_n y_n$ in E_n such that for every $k \in \mathbb{N}$ with $y_n \rightarrow w$ in E_k as $n \rightarrow \infty$ in $S = \{k+1, k+2, \dots\}$, then $w \in Fw$ in E .

Then F has a fixed point in E .

Proof. The proof is essentially the same as in Theorem 2.4 (except that here we use Theorem 1.3 (and Remark 1.4) instead of Theorem 1.2). $\square$

References

- [1] R. P. Agarwal, J. Dshalalow, and D. O'Regan, *Fixed point and homotopy results for generalized contractive maps of Reich type*, *Applicable Analysis* **82** (2003), no. 4, 329–350.
- [2] R. P. Agarwal and D. O'Regan, *Various homotopy results for generalized single valued contractions on Fréchet spaces*, *Journal of Concrete and Applicable Mathematics* **1** (2003), no. 2, 101–111.
- [3] M. Frigon, *Fixed point results for generalized contractions in gauge spaces and applications*, *Proceedings of the American Mathematical Society* **128** (2000), no. 10, 2957–2965.
- [4] ———, *Fixed point results for compact maps on closed subsets of Fréchet spaces and applications to differential and integral equations*, *Bulletin of the Belgian Mathematical Society. Simon Stevin* **9** (2002), no. 1, 23–37.
- [5] L. V. Kantorovich and G. P. Akilov, *Functional Analysis in Normed Spaces*, Pergamon Press, Oxford, 1964.

8 Multivalued nonlinear contractions

- [6] D. O'Regan, *Fixed point theorems for nonlinear operators*, Journal of Mathematical Analysis and Applications **202** (1996), no. 2, 413–432.

Ravi P. Agarwal: Department of Mathematical Science, Florida Institute of Technology, Melbourne, FL 32901, USA

E-mail address: agarwal@fit.edu

Donal O'Regan: Department of Mathematics, National University of Ireland, Galway, Ireland

E-mail address: donal.oregan@nuigalway.ie