PARABOLIC INEQUALITIES IN L^1 AS LIMITS OF RENORMALIZED EQUATIONS

K. AZELMAT, M. KBIRI ALAOUI, D. MESKINE, AND A. SOUISSI

Received 21 April 2006; Revised 10 August 2006; Accepted 21 August 2006

The paper deals with the existence of solutions of some parabolic bilateral problems approximated by the renormalized solutions of some parabolic equations.

Copyright © 2006 Hindawi Publishing Corporation. All rights reserved.

1. Introduction

Let Ω be a bounded domain in $\mathbb{R}^N$ and $T > 0$. We denote by Q the cylinder $\Omega \times (0, T)$ and $\Gamma = \partial Q$.

Let

$$A(u) = -\operatorname{div}(a(x, t, u, \nabla u)) \quad (1.1)$$

be a Leray-Lions operator acting on $L^p(0, T; W_0^{1,p}(\Omega))$, $1 < p < \infty$, into its dual $L^{p'}(0, T; W^{-1,p'}(\Omega))$ ($1/p + 1/p' = 1$). Consider the following parabolic problem:

$$u \in \mathcal{K} = \{v \in L^p(0, T; W_0^{1,p}(\Omega)) : v(t) \in K \text{ a.e.}\},$$

$$\int_0^T \left\langle \frac{\partial v}{\partial t}, u - v \right\rangle dt + \int_Q a(x, t, u, \nabla u)(\nabla u - \nabla v) dx dt \leq \int_0^T \langle f, u - v \rangle dt, \quad (P)$$

$$\forall v \in \mathcal{K} \cap \left\{ v \in L^p(0, T; W_0^{1,p}(\Omega)) : \frac{\partial v}{\partial t} \in L^{p'}(0, T; W^{-1,p'}(\Omega)); v(0) = 0 \right\},$$

where K is a given convex in $W_0^{1,p}(\Omega)$ and $f \in L^{p'}(0, T; W^{-1,p'}(\Omega))$.

It is well known that (P) admits at least one solution via a classical penalty method (see Lions [5] for $p \geq 2$ and Landes-Mustonen [4] for $1 < p < 2$). Recently in [6], the authors

2 Parabolic inequalities in L^1

approximated (P) by the following sequence of parabolic equations:

$$\begin{aligned} \frac{\partial u_n}{\partial t} + A(u_n) + |h(x, u_n)|^{n-1} h(x, u_n) |G(x, t, u_n, \nabla u_n)| &= f \quad \text{in } Q, \\ u_n(x, t) &= 0 \quad \text{on } \partial Q, \\ u_n(x, 0) &= 0 \quad \text{in } \Omega, \end{aligned} \tag{P_n}$$

where h and G are two Carathéodory functions satisfying some natural growth conditions. The obtained convex K depends on two obstacles constructed from h .

In the L^1 case, that is, $f \in L^1(\Omega \times]0, T[)$, the formulations (P) and (P_n) are not appropriate. So, we introduce the renormalized problem (R_n) associated to (P_n) (see the definition below). The study of the asymptotic behavior of (R_n) as $n \rightarrow \infty$ leads to some bilateral parabolic problem. Our approach allows us also to prove the existence of solutions for general parabolic inequalities of type

$$T_k(u) \in \mathcal{H},$$

$$\begin{aligned} \int_0^T \left\langle \frac{\partial v}{\partial t}, T_k(u - v) \right\rangle dt + \int_Q a(x, t, u, \nabla u) \nabla T_k(u - v) dx dt \\ + \int_Q H(x, t, u, \nabla u) T_k(u - v) dx dt \leq \int_Q f T_k(u - v) dx dt, \quad \forall v \in \mathcal{H} \cap D \cap L^\infty(Q), \end{aligned} \tag{1.2}$$

where $D = \{v \in L^p(0, T; W_0^{1,p}(\Omega)), \partial v / \partial t \in L^{p'}(0, T, W_0^{-1,p'}(\Omega)) + L^1(Q), v(0) = 0\}$ and where H is a given Carathéodory function satisfying some natural growth assumption.

For some recent and classical results for some parabolic inequalities problems, the reader can refer to [2, 7, 9, 10].

2. Main result

Let Ω be an open bounded subset of $\mathbb{R}^N$, $N \geq 2$ and $1 < p < +\infty$.

We denote by Q the cylinder $\Omega \times (0, T)$ and $\Gamma = \partial Q$.

Let $A(u) = -\operatorname{div}(a(x, t, \nabla u))$ be a Leray-Lions operator defined on $L^p(0, T; W_0^{1,p}(\Omega))$ into its dual $L^{p'}(0, T; W^{-1,p'}(\Omega))$, where $a : \Omega \times \mathbb{R} \times \mathbb{R}^N \rightarrow \mathbb{R}^N$ is a Carathéodory function satisfying for a.e. $x \in \Omega$, for all $t \in \mathbb{R}$ and for all $\zeta, \zeta' \in \mathbb{R}^N$, ($\zeta \neq \zeta'$) the following hold:

$$\begin{aligned} |a(x, t, \zeta)| &\leq \beta(k(x, t) + |\zeta|^{p-1}), \\ (a(x, t, \zeta) - a(x, t, \zeta'))(\zeta - \zeta') &> 0, \\ a(x, t, \zeta)\zeta &\geq \alpha|\zeta|^p, \end{aligned} \tag{2.1}$$

with $\alpha > 0, \beta > 0, k \in L^{p'}(Q)$.

Furthermore, let $h : \Omega \times \mathbb{R} \rightarrow \mathbb{R}$ be a Carathéodory function such that

$$h(x, 0) = 0, \quad h(x, s) \text{ is nondecreasing with respect to } s. \tag{2.2}$$

G is a Carathéodory function satisfying the following assumptions:

$$|G(x, t, s, \xi)| \leq b(|s|)(c(x, t) + |\xi|^p), \quad G(x, t, s, 0) = 0, \quad (2.3)$$

$$\begin{aligned} & \{v \in L^p(0, T; W_0^{1,p}(\Omega)) : G(x, t, v, \nabla v) = 0 \text{ a.e. in } Q\} \\ & \subset \{v \in L^p(0, T; W_0^{1,p}(\Omega)) : |h(x, v)| \leq 1 \text{ a.e. in } Q\}. \end{aligned} \quad (2.4)$$

Let us suppose

$$\begin{aligned} & \text{for almost } x \in \Omega \setminus \Omega_+^\infty \text{ there exists } \epsilon = \epsilon(x) > 0 \text{ such that} \\ & \quad h(x, s) > 1, \quad \forall s \in]q_+(x), q_+(x) + \epsilon[, \\ & \text{for almost } x \in \Omega \setminus \Omega_-^\infty \text{ there exists } \epsilon = \epsilon(x) > 0 \text{ such that} \\ & \quad h(x, s) < -1, \quad \forall s \in]q_-(x) - \epsilon, q_-(x)[, \end{aligned} \quad (2.5)$$

where b is a continuous nondecreasing function and $c(x, t) \in L^1(Q)$, $c \geq 0$, and

$$\begin{aligned} q_+(x) &= \inf \{s > 0, h(x, s) \geq 1\}, \\ q_-(x) &= \sup \{s > 0, h(x, s) \leq -1\}, \\ \Omega_+^\infty &= \{x \in \Omega : q_+(x) = +\infty\}, \\ \Omega_-^\infty &= \{x \in \Omega : q_-(x) = -\infty\}. \end{aligned} \quad (2.6)$$

We define for all s and k in $\mathbb{R}$, $k \geq 0$, $T_k(s) = \max(-k, \min(k, s))$.

We will say that u_n is a renormalized solution of (P_n) if

$$\begin{aligned} & T_k(u_n) \in L^p(0, T; W_0^{1,p}(\Omega)), \quad \forall k > 0, \\ & \lim_{h \rightarrow \infty} \int_{\{h \leq |u_n| \leq h+1\}} a(x, t, \nabla u_n) \nabla u_n dx dt = 0, \\ & u_n \text{ satisfies in the distributional sense} \\ & (A(u_n))_t - \operatorname{div}(a(x, t, \nabla u_n) A'(u_n)) + a(x, t, \nabla u_n) \nabla u_n A''(u_n) \\ & \quad + |h(x, u_n)|^{n-1} h(x, u_n) |G(x, t, u_n, \nabla u_n)| A'(u_n) = f A'(u_n), \\ & \forall A \in C^1(\mathbb{R}), A', A'' \in L^\infty(\Omega), \quad A' \text{ has a compact support and } u_n \text{ satisfies} \\ & \text{the initial condition in the sense that } A(u_n) \in C([0, T], L^1(\Omega)). \end{aligned} \quad (R_n)$$

Thanks to [8, Theorem 3.2, page 164], there exists at least one solution u_n of (R_n) .

THEOREM 2.1. *Under the hypotheses (2.1)–(2.5), $f \in L^1(Q)$, the problem (P_n) has at least one renormalized solution (u_n) such that*

$$T_k(u_n) \longrightarrow T_k(u) \quad \text{strongly in } L^p(0, T; W_0^{1,p}(\Omega)), \quad (2.7)$$

4 Parabolic inequalities in L^1

where u is a solution of the following obstacle problem:

$$\begin{aligned} q_-(x) &\leq u(x, t) \leq q_+(x) \quad \text{a.e. } (x, t) \in Q, \\ T_k(u) &\in L^p(0, T; W_0^{1,p}(\Omega)), \\ \int_0^T \left\langle \frac{\partial v}{\partial t}, T_k(u - v) \right\rangle dt + \int_Q a(x, t, \nabla u) \nabla T_k(u - v) dx dt \\ &\leq \int_Q f T_k(u - v) dx dt, \quad \forall v \in \mathcal{H} \cap D \cap L^\infty(Q), \end{aligned} \tag{R}$$

where

$$\begin{aligned} D &= \left\{ v \in L^p(0, T; W_0^{1,p}(\Omega)), \frac{\partial v}{\partial t} \in L^{p'}(0, T; W^{-1,p'}(\Omega)) + L^1(Q), v(0) = 0 \right\}, \\ \mathcal{H} &= \{v \in L^p(0, T; W_0^{1,p}(\Omega)), v(t) \in K\}, \quad K = \{v \in W_0^{1,p}(\Omega), q_- \leq v \leq q_+\}. \end{aligned} \tag{2.8}$$

Moreover, if $q_-, q_+ \in L^\infty(\Omega)$, then $u \in L^p(0, T; W_0^{1,p}(\Omega)) \cap L^\infty(Q)$.

Remark 2.2. The same result can be obtained when dealing with general operator of Leray-Lions type depending also on u , that is, $A(u) = -\operatorname{div}(a(x, t, u, \nabla u))$.

Proof of Theorem 2.1.

Step 1. Let $A(t) = H_m(t)$, $H_m(t) = \int_0^t h_m(s) ds$, where

$$h_m(s) = \begin{cases} 1 & \text{if } |s| \leq m, \\ \text{affine} & \text{if } m \leq |s| \leq m+1, \\ 0 & \text{if } m+1 \leq |s|. \end{cases} \tag{2.9}$$

Taking now $T_k(H_m(u_n))$ as test function in (R_n) , we obtain

$$\begin{aligned} \int_0^T \left\langle \frac{\partial H_m(u_n)}{\partial t}, T_k(H_m(u_n)) \right\rangle dt + \int_{|H_m(u_n)| < k} a(x, t, \nabla u_n) \nabla u_n h_m^2(u_n) dx dt \\ + \int_Q |h(x, u_n)|^{n-1} h(x, u_n) |G(x, t, u_n, \nabla u_n)| h_m(u_n) T_k(H_m(u_n)) dx dt \\ + \int_Q a(\cdot, t, \nabla u_n) \nabla u_n h'_m(u_n) T_k(H_m(u_n)) dx dt = \int_Q f h_m(u_n) T_k(H_m(u_n)) dx dt. \end{aligned} \tag{2.10}$$

Since

$$\begin{aligned} \int_0^T \left\langle \frac{\partial H_m(u_n)}{\partial t}, T_k(H_m(u_n)) \right\rangle dt \\ = \int_\Omega \left(\int_0^{H_m(u_n(x, T))} T_k(s) ds \right) dx - \int_\Omega \left(\int_0^{H_m(u_n(x, 0))} T_k(s) ds \right) dx \end{aligned} \tag{2.11}$$

and by using the fact that $\int_{\Omega} (\int_0^{H_m(u_n(x,T))} T_k(s) ds) \geq 0$, we obtain

$$\begin{aligned} \int_{\{|H_m(u_n)| < k\}} a(x, t, \nabla u_n) \nabla u_n h_m^2(u_n) dx dt &\leq Ck + \int_{\{m \leq |u_n| \leq m+1\}} a(x, t, \nabla u_n) \nabla u_n dx dt, \\ \int_Q |h(x, u_n)|^{n-1} h(x, u_n) |G(x, t, u_n, \nabla u_n)| h_m(u_n) T_k(H_m(u_n)) dx dt \\ &\leq Ck + \int_{\{m \leq |u_n| \leq m+1\}} a(x, t, \nabla u_n) \nabla u_n dx dt. \end{aligned} \quad (2.12)$$

We have $H_m(s)$ (resp., $h_m(s)$) tends to s (resp., to 1) as m goes to $+\infty$.

Using Fatou's lemma and the definition of the renormalized solution leads to

$$\int_Q |\nabla T_k(u_n)|^p dx dt \leq Ck, \quad (2.13)$$

$$\int_Q |h(x, u_n)|^{n-1} h(x, u_n) |G(x, t, u_n, \nabla u_n)| T_k(u_n) dx dt \leq Ck, \quad (2.14)$$

which gives

$$\int_Q |h(x, u_n)|^n |G(x, t, u_n, \nabla u_n)| \frac{|T_k(u_n)|}{k} dx dt \leq C, \quad (2.15)$$

and as $k \rightarrow 0$ we obtain

$$\int_Q |h(x, u_n)|^n |G(x, t, u_n, \nabla u_n)| dx dt \leq C. \quad (2.16)$$

Choosing now a C^2 function ρ_k , such that $\rho_k(s) = s$ for $|s| \leq k$ and $2k \text{sign}(s)$ for $|s| > 2k$, we get

$$\begin{aligned} (\rho_k(u_n))_t - \text{div}(a(x, t, \nabla u_n) \rho'_k(u_n)) + a(x, t, \nabla u_n) \nabla u_n \rho''_k(u_n) \\ + |h(x, u_n)|^{n-1} h(x, u_n) |G(x, t, u_n, \nabla u_n)| \rho'_k(u_n) = f \rho_k(u_n). \end{aligned} \quad (2.17)$$

We deduce that $(\rho_k(u_n))_t$ is bounded in $L^1(Q) + L^{p'}(0, T; W^{-1, p'}(\Omega))$.

Now thanks to the following result.

LEMMA 2.3 [11]. *Let $p > 1$. If (u_n) is a bounded sequence of $L^p(0, T; W_0^{1, p}(\Omega))$ such that $\partial u_n / \partial t$ is bounded in $L^1 + L^{p'}(0, T; W^{-1, p'}(\Omega))$, then u_n is relatively compact in $L^p(Q)$.*

6 Parabolic inequalities in L^1

We deduce that $\rho_k(u_n)$ is relatively compact in $L^p(Q)$ and so there exists a measurable function u such that $u_n \rightarrow u$ a.e. in Q .

Finally, we deduce from (2.13) that $T_k(u_n) \rightharpoonup T_k(u)$ weakly in $L^p(0, T; W_0^{1,p}(\Omega))$, and strongly in $L^p(Q)$.

Step 2. We are dealing now with the almost convergence of the gradient.

We have to prove that, for $0 < \theta < 1$,

$$\lim_{n \rightarrow \infty} \int_Q ([a(x, t, \nabla T_k(u_n)) - a(x, t, \nabla T_k(u))] [\nabla T_k(u_n) - \nabla T_k(u)])^\theta dx dt = 0. \quad (2.18)$$

Let $\omega \in L^p(0, T; W_0^{1,p}(\Omega))$, we define for any $\mu > 0$, ω_μ the time regularization of ω ,

$$\omega_\mu(x, t) = \mu \int_{-\infty}^t \bar{\omega}(x, s) \exp(\mu(s - t)) ds, \quad (2.19)$$

where $\bar{\omega}$ is the zero extension of ω for $s > T$. Furthermore, ω_μ satisfies the following properties (see [3]):

$$\begin{aligned} \omega_\mu &\longrightarrow \omega \quad \text{strongly in } L^p(0, T; W_0^{1,p}(\Omega)), \\ \frac{\partial \omega_\mu}{\partial t} &= \mu(\omega - \omega_\mu) \quad \text{in the distributional sense.} \end{aligned} \quad (2.20)$$

Letting $\eta > 0$, we obtain

$$\begin{aligned} &\int_Q ([a(x, t, \nabla T_k(u_n)) - a(x, t, \nabla T_k(u))] [\nabla T_k(u_n) - \nabla T_k(u)])^\theta \\ &\leq C \text{meas} \{ |T_k(u_n) - T_k(u)_\mu| \geq \eta \}^{1-\theta} \\ &\quad + C \left(\int_{\{|T_k(u_n) - T_k(u)_\mu| < \eta\}} [a(x, t, \nabla T_k(u_n)) \right. \\ &\quad \left. - a(x, t, \nabla T_k(u))] [\nabla T_k(u_n) - \nabla T_k(u)] \right)^\theta. \end{aligned} \quad (2.21)$$

On the other hand, we have

$$\begin{aligned}
& \int_{\{|T_k(u_n) - T_k(u)_\mu| < \eta\}} [a(x, t, \nabla T_k(u_n)) - a(x, t, \nabla T_k(u))] [\nabla T_k(u_n) - \nabla T_k(u)] dx dt \\
& \leq \int_{\{|T_k(u_n) - T_k(u)_\mu| < \eta\}} [a(x, t, \nabla T_k(u_n)) \\
& \quad - a(x, t, \nabla T_k(u)_\mu)] [\nabla T_k(u_n) - \nabla T_k(u)_\mu] dx dt \\
& \quad + \int_{\{|T_k(u_n) - T_k(u)_\mu| < \eta\}} a(x, t, \nabla T_k(u_n)) (\nabla T_k(u)_\mu - \nabla T_k(u)) dx dt \\
& \quad + \int_{\{|T_k(u_n) - T_k(u)_\mu| < \eta\}} [a(x, t, \nabla T_k(u)_\mu) - a(x, t, \nabla T_k(u))] \nabla T_k(u_n) dx dt \\
& \quad - \int_{\{|T_k(u_n) - T_k(u)_\mu| < \eta\}} a(x, t, \nabla T_k(u)_\mu) \nabla T_k(u)_\mu dx dt \\
& \quad + \int_{\{|T_k(u_n) - T_k(u)_\mu| < \eta\}} a(x, t, \nabla T_k(u)) \nabla T_k(u) dx dt \\
& \leq I_1 + I_2 + I_3 + I_4 + I_5.
\end{aligned} \tag{2.22}$$

Take $T_\eta(H_m(u_n) - T_k(u)_\mu)$ as test function in (R_n) with $A(t) = H_m(t)$. We obtain

$$\begin{aligned}
& \int_0^T \left\langle \frac{\partial H_m(u_n)}{\partial t}, T_\eta(H_m(u_n) - T_k(u)_\mu) \right\rangle dt \\
& \quad + \int_{\{|H_m(u_n) - T_k(u)_\mu| < \eta\}} a(x, t, \nabla u_n) \nabla u_n h_m^2(u_n) - \nabla T_k(u)_\mu h_m(u_n) dx dt \\
& \quad + \int_Q |h(x, u_n)|^{n-1} h(x, u_n) |G(x, t, u_n, \nabla u_n)| h_m(u_n) T_\eta(H_m(u_n) - T_k(u)_\mu) dx dt \\
& \quad + \int_Q a(x, t, \nabla u_n) \nabla u_n h'_m(u_n) T_\eta(H_m(u_n) - T_k(u)_\mu) dx dt \\
& = \int_Q f h_m(u_n) T_\eta(H_m(u_n) - T_k(u)_\mu) dx dt.
\end{aligned} \tag{2.23}$$

We have

$$\begin{aligned}
& \int_0^T \left\langle \frac{\partial H_m(u_n)}{\partial t}, T_\eta(H_m(u_n) - T_k(u)_\mu) \right\rangle dt \\
& = \int_0^T \left\langle \frac{\partial H_m(u_n) - T_k(u)_\mu}{\partial t}, T_k(H_m(u_n) - T_k(u)_\mu) \right\rangle dt \\
& \quad + \int_0^T \left\langle \frac{\partial T_k(u)_\mu}{\partial t}, T_k(H_m(u_n) - T_k(u)_\mu) \right\rangle dt.
\end{aligned} \tag{2.24}$$

8 Parabolic inequalities in L^1

Using the fact that

$$\begin{aligned}
 & \int_0^T \left\langle \frac{\partial H_m(u_n) - T_k(u)_\mu}{\partial t}, T_\eta(H_m(u_n) - T_k(u)_\mu) \right\rangle dt \geq 0, \\
 & \int_0^T \left\langle \frac{\partial T_k(u)_\mu}{\partial t}, T_k(H_m(u_n) - T_k(u)_\mu) \right\rangle dt \\
 & = \mu \int_Q (T_k(u) - T_k(u)_\mu) T_\eta(H_m(u_n) - T_k(u)_\mu) dx dt,
 \end{aligned} \tag{2.25}$$

consequently,

$$\begin{aligned}
 & \limsup_{n \rightarrow \infty} \limsup_{m \rightarrow \infty} \int_0^T \left\langle \frac{\partial H_m(u_n)}{\partial t}, T_\eta(H_m(u_n) - T_k(u)_\mu) \right\rangle dt \\
 & \geq \mu \int_Q (T_k(u) - T_k(u)_\mu) T_\eta(u - T_k(u)_\mu) dx dt = \epsilon(m, n) \geq 0.
 \end{aligned} \tag{2.26}$$

This implies that

$$\begin{aligned}
 & \int_{\{|H_m(u_n) - T_k(u)_\mu| < \eta\}} a(x, t, \nabla u_n) \nabla u_n h_m^2(u_n) - \nabla T_k(u)_\mu h_m(u_n) dx dt \\
 & + \int_Q |h(x, u_n)|^{n-1} h(x, u_n) |G(x, t, u_n, \nabla u_n)| h_m(u_n) T_\eta(H_m(u_n) - T_k(u)_\mu) dx dt \\
 & + \int_Q a(x, t, \nabla u_n) \nabla u_n h'_m(u_n) T_\eta(H_m(u_n) - T_k(u)_\mu) dx dt \\
 & \leq \int_Q f h_m(u_n) T_\eta(H_m(u_n) - T_k(u)_\mu) dx dt + \epsilon(m, n),
 \end{aligned} \tag{2.27}$$

which gives by using the fact that

$$\begin{aligned}
 & \int_Q |h(x, u_n)|^{n-1} h(x, u_n) |G(x, t, u_n, \nabla u_n)| h_m(u_n) T_\eta(H_m(u_n) - T_k(u)_\mu) dx dt \leq C\eta, \\
 & \int_{\{|H_m(u_n) - T_k(u)_\mu| < \eta\}} a(x, t, \nabla u_n) \nabla u_n h_m^2(u_n) - \nabla T_k(u)_\mu h_m(u_n) dx dt \\
 & \leq C\eta + \epsilon(m, n) + \eta \int_{\{m \leq |u_n| \leq m+1\}} a(x, t, \nabla u_n) \nabla u_n dx dt,
 \end{aligned} \tag{2.28}$$

which gives as $m \rightarrow \infty$,

$$\int_{\{|u_n - T_k(u)_\mu| < \eta\}} a(x, t, \nabla u_n) \nabla u_n - \nabla T_k(u)_\mu dx dt \leq C\eta + \epsilon(n). \tag{2.29}$$

Finally from (2.22),

$$|I_1| \leq C\eta + \epsilon(n) - \int_{\{|T_k(u_n) - T_k(u)_\mu| < \eta\}} a(x, t, \nabla T_k(u)_\mu) (\nabla T_k(u_n) - \nabla T_k(u)). \quad (2.30)$$

Since $a(x, t, \nabla T_k(u)_\mu) \chi_{\{|T_k(u_n) - T_k(u)_\mu| < \eta\}} \rightarrow a(x, t, \nabla T_k(u)_\mu) \chi_{\{|T_k(u) - T_k(u)_\mu| < \eta\}}$ in $L^{p'}(Q)$ and $T_k(u_n) \rightarrow T_k(u)$ weakly in $L^p(0, T; W_0^{1,p}(\Omega))$, then

$$\begin{aligned} & - \int_{\{|T_k(u_n) - T_k(u)_\mu| < \eta\}} a(x, t, \nabla T_k(u)_\mu) (\nabla T_k(u_n) - \nabla T_k(u)) dx dt \\ & = - \int_{\{|T_k(u) - T_k(u)_\mu| < \eta\}} a(x, t, \nabla T_k(u)_\mu) (\nabla T_k(u) - \nabla T_k(u)) dx dt + \epsilon(n). \end{aligned} \quad (2.31)$$

So

$$|I_1| \leq C\eta + \epsilon(n). \quad (2.32)$$

For what concerns the term I_2 , one has

$$I_2 = \epsilon(n, \mu), \quad (2.33)$$

since

$$\begin{aligned} a(x, t, \nabla T_k(u_n)) \chi_{\{|T_k(u_n) - T_k(u)_\mu| < \eta\}} & \longrightarrow a(x, t, \nabla T_k(u)) \chi_{\{|T_k(u) - T_k(u)_\mu| < \eta\}} \quad \text{in } (L^{p'}(Q))^N, \\ (\nabla T_k(u)_\mu - \nabla T_k(u)) \chi_{\{|T_k(u_n) - T_k(u)_\mu| < \eta\}} & \longrightarrow (\nabla T_k(u)_\mu - \nabla T_k(u)) \chi_{\{|T_k(u) - T_k(u)_\mu| < \eta\}}. \end{aligned} \quad (2.34)$$

In the same way, we show that

$$I_3 = \epsilon(n, \mu), \quad I_4 = \epsilon(n, \mu), \quad I_5 = \epsilon(n, \mu). \quad (2.35)$$

Combining the above estimates, we get

$$\lim_{n \rightarrow \infty} \int_Q ([a(x, t, \nabla T_k(u_n)) - a(x, t, \nabla T_k(u))] [\nabla T_k(u_n) - \nabla T_k(u)])^\theta dx dt = 0. \quad (2.36)$$

Then there exists a subsequence also denoted by (u_n) such that

$$\nabla u_n \longrightarrow \nabla u \quad \text{a.e. in } Q. \quad (2.37)$$

Step 3. From (2.16), we deduce that

$$\int_Q |h(x, u_n)|^n |G(x, t, u_n, \nabla u_n)| dx dt \leq C, \quad (2.38)$$

which gives for every $\beta > 0$,

$$\int_{|h(x, T_\beta(u_n))| > k} |G(x, t, T_\beta(u_n), \nabla T_\beta(u_n))| dx dt \leq \frac{C}{k^n}, \quad (2.39)$$

where $k > 1$. Letting $n \rightarrow +\infty$ for k fixed, we deduce by using Fatou's lemma

$$\int_{|h(x, T_\beta(u))| > k} |G(x, t, T_\beta(u), \nabla T_\beta(u))| dx dt = 0, \quad (2.40)$$

and so, by (2.4)

$$|h(x, T_\beta(u))| \leq 1 \quad \text{a.e. in } Q. \quad (2.41)$$

So

$$q_-(x) \leq T_\beta(u(x)) \leq q_+(x) \quad \text{a.e. in } Q. \quad (2.42)$$

Letting now $\beta \rightarrow +\infty$, we deduce also that

$$q_-(x) \leq u(x) \leq q_+(x) \quad \text{a.e. in } Q. \quad (2.43)$$

Step 4. Strong convergence of the truncations.

We will prove that

$$\lim_{n \rightarrow \infty} \int_Q [a(x, t, \nabla T_k(u_n)) - a(x, t, \nabla T_k(u))] [\nabla T_k(u_n) - \nabla T_k(u)] dx dt = 0. \quad (2.44)$$

Fix $k > 0$ and let $\varphi(s) = \exp(\delta s^2)$, $\delta > 0$. Let $l > k$ and define the function $R_l(s) = \int_0^s \rho_l(t) dt$. Let us consider $\omega_\mu^m = T_k(H_m(u)_\mu)$, where v_μ is the mollification with respect to time v . Letting $v_\mu^{m,n} = \rho_l(H_m(u_n))\varphi(T_k(H_m(u_n)) - \omega_\mu^m)$ as test function in the problem (R_n) , we get

$$\begin{aligned} & \int_0^T \left\langle \frac{\partial H_m(u_n)}{\partial t}, \rho_l(H_m(u_n))\varphi(T_k(H_m(u_n)) - \omega_\mu^m) \right\rangle dt \\ & + \int_Q a(x, t, \nabla u_n) \nabla u_n h^2(u_n) \rho'_l(H_m(u_n))\varphi(T_k(H_m(u_n)) - \omega_\mu^m) dx dt \\ & + \int_Q a(x, t, \nabla u_n) (\nabla T_k(H_m(u_n)) - \nabla \omega_\mu^m) \\ & \times h_m(u_n) \rho_l(H_m(u_n))\varphi'(T_k(H_m(u_n)) - \omega_\mu^m) dx dt \\ & + \int_Q a(x, t, \nabla u_n) \nabla u_n h'_m(u_n) \rho_l(H_m(u_n))\varphi(T_k(H_m(u_n)) - \omega_\mu^m) dx dt \\ & + \int_Q |h(x, u_n)|^{n-1} h(x, u_n) |G(x, t, u_n, \nabla u_n)| \\ & \times h_m(u_n) \rho_l(H_m(u_n))\varphi(T_k(H_m(u_n)) - \omega_\mu^m) dx dt \\ & = \int_Q f v_\mu^{m,n} h_m(u_n) dx dt. \end{aligned} \quad (2.45)$$

We deal now with the estimate of each term of the last equalities.

Since $H_m(u_n) \in L^p(0, T; W_0^{1,p}(\Omega))$ and $\partial H_m(u_n)/\partial t \in L^{p'}(0, T; W^{-1,p'}(\Omega)) + L^1(Q)$, there exists a smooth function $H_m(u_n)_\sigma$ such that as $\sigma \rightarrow 0$,

$$\begin{aligned} H_m(u_n)_\sigma &\longrightarrow H_m(u_n) \quad \text{strongly in } L^p(0, T; W_0^{1,p}(\Omega)), \\ \frac{\partial H_m(u_n)_\sigma}{\partial t} &\longrightarrow \frac{\partial H_m(u_n)}{\partial t} \quad \text{strongly in } L^{p'}(0, T; W^{-1,p'}(\Omega)) + L^1(Q). \end{aligned} \quad (2.46)$$

This implies that

$$\begin{aligned} I &= \int_0^T \left\langle \frac{\partial H_m(u_n)}{\partial t}, \rho_l(H_m(u_n)) \varphi(T_k(H_m(u_n)) - \omega_\mu^m) \right\rangle dt \\ &= \lim_{\sigma \rightarrow 0^+} \int_Q (H_m(u_n)_\sigma)' \rho_l(H_m(u_n)_\sigma) \varphi(T_k(H_m(u_n)_\sigma) - \omega_\mu^m) dx dt \\ &= \lim_{\sigma \rightarrow 0^+} \int_Q [R_l(H_m(u_n)_\sigma)]' \varphi(T_k(H_m(u_n)_\sigma) - \omega_\mu^m) dx dt \\ &= \lim_{\sigma \rightarrow 0^+} \int_Q [R_l(H_m(u_n)_\sigma) - T_k(H_m(u_n)_\sigma)]' \varphi(T_k(H_m(u_n)_\sigma) - \omega_\mu^m) dx dt \\ &\quad + \int_Q [T_k(H_m(u_n)_\sigma)]' \varphi(T_k(H_m(u_n)_\sigma) - \omega_\mu^m) dx dt \\ &= \lim_{\sigma \rightarrow 0^+} \left\{ \int_\Omega [R_l(H_m(u_n)_\sigma) - T_k(H_m(u_n)_\sigma) \varphi(T_k(H_m(u_n)_\sigma) - \omega_\mu^m)]_0^T dx dt \right. \\ &\quad - \int_Q [R_l(H_m(u_n)_\sigma) - T_k(H_m(u_n)_\sigma)]' \varphi'(T_k(H_m(u_n)_\sigma) - \omega_\mu^m) (T_k(H_m(u_n)_\sigma) \\ &\quad \left. - \omega_\mu^m)' dx dt + \int_Q [T_k(H_m(u_n)_\sigma)]' \varphi(T_k(H_m(u_n)_\sigma) - \omega_\mu^m) dx dt \right\} \\ &= \lim_{\sigma \rightarrow 0^+} \{I_1(\sigma) + I_2(\sigma) + I_3(\sigma)\}. \end{aligned} \quad (2.47)$$

Observe that for $|s| \leq k$ we have $R_l(s) = T_k(s) = s$ and for $|s| > k$ we have $|R_l(s)| \geq |T_k(s)|$ and, since both $R_l(s)$ and $T_k(s)$ have the same sign of s , we deduce that $\text{sign}(s)(R_l(s) - T_k(s)) \geq 0$. Consequently,

$$I_1(\sigma) = \int_{\{|H_m(u_n)_\sigma| > k\}} [R_l(H_m(u_n)_\sigma) - T_k(H_m(u_n)_\sigma) \varphi(T_k(H_m(u_n)_\sigma) - \omega_\mu^m)]_0^T dx dt \geq 0. \quad (2.48)$$

12 Parabolic inequalities in L^1

We have, since $(R_l(s) - T_k(s))(T_k(s))' = 0$, for all s ,

$$\begin{aligned}
 I_2(\sigma) &= \int_{\{|H_m(u_n)_\sigma| > k\}} [R_l(H_m(u_n)_\sigma) - T_k(H_m(u_n)_\sigma) \varphi'(T_k(H_m(u_n)_\sigma) \\
 &\quad - \omega_\mu^m)] (\omega_\mu^m)' dx dt \\
 &= \mu \int_{\{|H_m(u_n)_\sigma| > k\}} [R_l(H_m(u_n)_\sigma) - T_k(H_m(u_n)_\sigma) \varphi'(T_k(H_m(u_n)_\sigma) \\
 &\quad - \omega_\mu^m)] (T_k(H_m(u_n)_\sigma) - \omega_\mu^m) dx dt,
 \end{aligned} \tag{2.49}$$

by using the fact that $\varphi' \geq 0$ and that

$$(R_l(H_m(u_n)_\sigma) - T_k(H_m(u_n)_\sigma))(T_k(H_m(u_n)_\sigma) - \omega_\mu^m) \chi_{\{|H_m(u_n)_\sigma| > k\}} \geq 0, \tag{2.50}$$

the last integral is of the form $\epsilon(m, n)$. We deduce that

$$\limsup_{\sigma \rightarrow 0^+} I_2(\sigma) \geq \epsilon(m, n). \tag{2.51}$$

For $I_3(\sigma)$, one has

$$\begin{aligned}
 I_3(\sigma) &= \int_Q [T_k(H_m(u_n)_\sigma)]' \varphi(T_k(H_m(u_n)_\sigma) - \omega_\mu^m) dx dt \\
 &= \int_Q [T_k(H_m(u_n)_\sigma) - \omega_\mu^m]' \varphi(T_k(H_m(u_n)_\sigma) - \omega_\mu^m) dx dt \\
 &\quad + \int_Q (\omega_\mu^m)' \varphi(T_k(H_m(u_n)_\sigma) - \omega_\mu^m) dx dt.
 \end{aligned} \tag{2.52}$$

Let $\Phi(s) = \int_0^s \varphi(t) dt$. Remark that $(T_k(H_m(u_n)_\sigma) - \omega_\mu^m) \varphi(T_k(H_m(u_n)_\sigma) - \omega_\mu^m) \geq 0$.

Integrating by parts, using the fact that $\Phi \geq 0$, and following the same way as above, we have

$$\limsup_{\sigma \rightarrow 0^+} I_3(\sigma) \geq \epsilon(m, n). \tag{2.53}$$

Combining these estimates, we conclude that

$$\int_0^T \left\langle \frac{\partial H_m(u_n)}{\partial t}, \rho_l(H_m(u_n)) \varphi(T_k(H_m(u_n)) - \omega_\mu^m) \right\rangle dt \geq \epsilon(m, n). \tag{2.54}$$

We set

$$I_4(m) = \int_Q |h(x, u_n)|^{n-1} h(x, u_n) \rho_l(H_m(u_n)) \varphi(T_k(H_m(u_n)_\sigma) - \omega_\mu^m) |G(x, t, u_n, \nabla u_n)|, \quad (2.55)$$

so we have

$$\limsup_{m \rightarrow \infty} I_4(m) \geq I_4^1 + I_4^2, \quad (2.56)$$

where

$$\begin{aligned} I_4^1 &= \int_{\{|u_n| < k, 0 \leq u_n \leq T_k(u)_\mu\}} \\ &\quad \times |h(x, u_n)|^{n-1} h(x, u_n) \varphi(T_k(u_n) - T_k(u)_\mu) \rho_l(u_n) |G(x, t, u_n, \nabla u_n)| dx dt, \\ I_4^2 &= \int_{\{|u_n| < k, T_k(u)_\mu \leq u_n \leq 0\}} \\ &\quad \times |h(x, u_n)|^{n-1} h(x, u_n) \varphi(T_k(u_n) - T_k(u)_\mu) \rho_l(u_n) |G(x, t, u_n, \nabla u_n)| dx dt. \end{aligned} \quad (2.57)$$

Since $q_- \leq T_k(u)_\mu \leq q_+$ (recall that $q_- \leq T_k(u) \leq q_+$) and $0 \leq \rho_l(u_n) \leq 1$, one easily has

$$\begin{aligned} |I_4^1| &\leq \int_{\{|u_n| < k\}} c(x, t) |\varphi(T_k(u_n) - T_k(u)_\mu)| \\ &\quad + \frac{b(k)}{\alpha} \int_{\{|u_n| < k\}} |\nabla u_n|^p |\varphi(T_k(u_n) - T_k(u)_\mu)| \\ &\leq \frac{b(k)}{\alpha} \int_Q [a(x, t, \nabla T_k(u_n)) - a(x, t, \nabla T_k(u)_\mu)] [\nabla T_k(u_n) - \nabla T_k(u)_\mu] \\ &\quad \times |\varphi(T_k(u_n) - T_k(u)_\mu)| dx dt + \epsilon(n, \mu), \end{aligned} \quad (2.58)$$

and also we have the same estimation of I_4^2 .

Then

$$\begin{aligned} |I_4^1| + |I_4^2| &\leq 2 \frac{b(k)}{\alpha} \int_Q [a(x, t, \nabla T_k(u_n)) - a(x, t, \nabla T_k(u)_\mu)] [\nabla T_k(u_n) - \nabla T_k(u)_\mu] \\ &\quad \times |\varphi(T_k(u_n) - T_k(u)_\mu)| dx dt + \epsilon(n, \mu). \end{aligned} \quad (2.59)$$

By denoting by J_1 the third term of (2.45), one can write

$$\begin{aligned}
J_1 &= \int_Q a(x, t, \nabla u_n) (\nabla T_k(H_m(u_n)) - \nabla T_k(H_m(u)))_\mu \\
&\quad \times h_m(u_n) \rho_l(H_m(u_n)) \varphi'(T_k(H_m(u_n)) - T_k(H_m(u)))_\mu dx dt \\
&= \int_Q a(x, t, \nabla T_k(u_n)) (\nabla T_k(H_m(u_n)) - \nabla T_k(H_m(u)))_\mu h_m(u_n) \\
&\quad \times \rho_l(H_m(u_n)) \varphi'(T_k(H_m(u_n)) - T_k(H_m(u)))_\mu dx dt \\
&\quad + \int_{\{|u_n|>k\}} a(x, t, \nabla u_n) (\nabla T_k(H_m(u_n)) - \nabla T_k(H_m(u)))_\mu h_m(u_n) \\
&\quad \times \rho_l(H_m(u_n)) \varphi'(T_k(H_m(u_n)) - T_k(H_m(u)))_\mu dx dt, \\
J_1 &= \int_Q a(x, t, \nabla T_k(u_n)) (\nabla T_k(u_n) - \nabla T_k(u))_\mu \rho_l(u_n) \varphi'(T_k(u_n) - T_k(u))_\mu dx dt \\
&\quad - \int_{\{|u_n|>k\}} \rho_l(u_n) a(x, t, \nabla u_n) (\nabla T_k(u))_\mu \varphi'(T_k(u_n) - T_k(u))_\mu dx dt + \epsilon(m).
\end{aligned} \tag{2.60}$$

Since $a(x, t, \nabla u_n) \rho_l(u_n)$ is bounded in $L^{p'}(Q)$, we deduce that

$$a(x, t, \nabla u_n) \rho_l(u_n) \rightharpoonup a(x, t, \nabla u) \rho_l(u) \quad \text{weakly in } L^{p'}(Q), \tag{2.61}$$

and so

$$\begin{aligned}
J_1 &= \int_Q (a(x, t, \nabla T_k(u_n)) - a(x, t, \nabla T_k(u))_\mu) (\nabla T_k(u_n) - \nabla T_k(u))_\mu \\
&\quad \times \varphi'(T_k(u_n) - T_k(u))_\mu dx dt + \epsilon(m, n, \mu).
\end{aligned} \tag{2.62}$$

Concerning the second term of (2.45), one easily has

$$\begin{aligned}
&\int_Q a(x, t, \nabla u_n) \nabla u_n h_m^2(u_n) \rho_l'(H_m(u_n)) \varphi(T_k(H_m(u_n)) - \omega_\mu^m) dx dt \\
&\leq \varphi(2k) \int_{\{|l \leq |u_n| < l+1\}} a(x, t, \nabla u_n) \nabla u_n dx dt,
\end{aligned} \tag{2.63}$$

and since

$$\int_{\{|l \leq |u_n| < l+1\}} a(x, t, \nabla u_n) \nabla u_n dx dt \leq \int_{|u_n|>l} |f| dx dt, \tag{2.64}$$

we deduce that

$$\begin{aligned} & \left| \int_Q a(x, t, \nabla u_n) \nabla u_n h_m^2(u_n) \rho_l'(H_m(u_n)) \varphi(T_k(H_m(u_n))) - \omega_\mu^m dx dt \right| \\ & \leq \varphi(2k) \int_{|u_n| > l} |f| dx dt = \epsilon(n, l). \end{aligned} \quad (2.65)$$

The same result can be obtained for the fourth term of (2.45).

Combining (2.45)–(2.65), using the fact that $\phi' - 2(b(k)/\alpha)|\phi| \geq 1/2$ for $\delta \geq (b(k)/\alpha)^2$, we deduce that

$$\lim_{n \rightarrow \infty} \int_Q [a(x, t, \nabla T_k(u_n)) - a(x, t, \nabla T_k(u)_\mu)] [\nabla T_k(u_n) - \nabla T_k(u)_\mu] dx dt = 0. \quad (2.66)$$

On the other hand, we have

$$\begin{aligned} & \int_Q [a(x, t, \nabla T_k(u_n)) - a(x, t, \nabla T_k(u))] [\nabla T_k(u_n) - \nabla T_k(u)] dx dt \\ & - \int_Q [a(x, t, \nabla T_k(u_n)) - a(x, t, \nabla T_k(u)_\mu)] [\nabla T_k(u_n) - \nabla T_k(u)_\mu] dx dt \\ & = \int_Q a(x, t, \nabla T_k(u_n)) (\nabla T_k(u)_\mu - \nabla T_k(u)) dx dt \\ & - \int_Q a(x, t, \nabla T_k(u)) (\nabla T_k(u_n) - \nabla T_k(u)) dx dt \\ & + \int_Q a(x, t, \nabla T_k(u)_\mu) (\nabla T_k(u_n) - \nabla T_k(u)_\mu) dx dt = \epsilon(n, \mu). \end{aligned} \quad (2.67)$$

Consequently by [1, Lemma 5], we obtain

$$T_k(u_n) \longrightarrow T_k(u) \quad \text{strongly in } L^p(0, T; W_0^{1,p}(\Omega)) \text{ for every } k > 0. \quad (2.68)$$

Step 5 (passage to the limit). Letting $v \in D \cap \mathcal{H} \cap L^\infty(Q)$, and using $T_k(H_m(u_n) - \theta v)$ as test function in the problem (R_n) , we obtain

$$\begin{aligned} & \int_0^T \left\langle \frac{\partial H_m(u_n)}{\partial t}, T_k(H_m(u_n) - \theta v) \right\rangle dt + \int_Q a(x, t, \nabla u_n) \nabla T_k(H_m(u_n) - \theta v) h_m(u_n) dx dt \\ & + \int_Q a(x, t, \nabla u_n) \nabla u_n T_k(H_m(u_n) - \theta v) h_m'(u_n) dx dt \\ & + \int_Q |h(x, u_n)|^{n-1} h(x, u_n) |G(x, t, u_n, \nabla u_n)| h_m(u_n) T_k(H_m(u_n) - \theta v) dx dt \\ & \leq \int_Q f T_k(H_m(u_n) - \theta v) h_m(u_n) dx dt. \end{aligned} \quad (2.69)$$

We have

$$\begin{aligned}
& \int_Q |h(x, u_n)|^{n-1} h(x, u_n) |G(x, t, u_n, \nabla u_n)| h_m(u_n) T_k(H_m(u_n) - \theta v) dx dt \\
& \geq \int_{\{0 \leq H_m(u_n) \leq \theta v\}} |h(x, u_n)|^{n-1} \\
& \quad \times h(x, u_n) |G(x, t, u_n, \nabla u_n)| h_m(u_n) T_k(H_m(u_n) - \theta v) dx dt \\
& \quad + \int_{\{\theta v \leq H_m(u_n) \leq 0\}} |h(x, u_n)|^{n-1} \\
& \quad \times h(x, u_n) |G(x, t, u_n, \nabla u_n)| h_m(u_n) T_k(H_m(u_n) - \theta v) dx dt.
\end{aligned} \tag{2.70}$$

Now we deal with the estimation of the last two terms in the right-hand side of the last inequality which we denote, respectively, by $J'_1(m, n)$ and $J'_2(m, n)$. Let us define

$$\delta_1(x, t) = \sup_{0 \leq s \leq \theta v} h(x, s), \tag{2.71}$$

then we get $0 \leq \delta_1(x, t) < 1$ a.e. in Q .

We have

$$\begin{aligned}
\limsup_{m \rightarrow \infty} |J'_1(m, n)| & \leq k \int_{\{0 \leq u_n \leq \theta v\}} (\delta(x, t))^n (c(x, t) + |\nabla u_n|^p) \\
& \leq \int_{\{|u_n| \leq \|v\|_\infty\}} (\delta(x, t))^n (c(x, t) + |\nabla u_n|^p),
\end{aligned} \tag{2.72}$$

and by using the strong convergence of $T_{\|v\|_\infty}(u_n)$ in $L^p(0, T; W_0^{1,p}(\Omega))$, we deduce that

$$\limsup_{n \rightarrow \infty} \limsup_{m \rightarrow \infty} |J'_1(m, n)| = 0, \tag{2.73}$$

with the same technique (taking $\delta_2(x, t) = \inf_{\theta v \leq s \leq 0} h(x, s)$), we can see that

$$\limsup_{m \rightarrow \infty} |J'_2(n, m)| \rightarrow 0 \quad \text{as } n \rightarrow +\infty. \tag{2.74}$$

On the other hand,

$$\begin{aligned}
& \int_Q a(x, t, \nabla u_n) \nabla T_k(H_m(u_n) - \theta v) h_m(u_n) dx dt \\
& = \int_Q a(x, t, \nabla u_n) \nabla (H_m(u_n) - \theta v) \chi_{\{|H_m(u_n) - \theta v| \leq k\}} h_m(u_n) dx dt \\
& = \int_Q (a(x, t, \nabla u_n) - a(x, t, \theta \nabla v)) \nabla (H_m(u_n) - \theta v) \chi_{\{|H_m(u_n) - \theta v| \leq k\}} h_m(u_n) dx dt \\
& \quad + \int_Q a(x, t, \theta \nabla v) \nabla (H_m(u_n) - \theta v) \chi_{\{|H_m(u_n) - \theta v| \leq k\}} h_m(u_n) dx dt.
\end{aligned} \tag{2.75}$$

Since $a(x, t, \theta v)$ belongs to $(L^{p'}(Q))^N$, using Fatou's lemma in the first term of the last side gives

$$\liminf_{n, m \rightarrow +\infty} \int_0^T \langle Au_n, T_k(H_m(u_n) - \theta v) \rangle dt \geq \int_0^T \langle Au, T_k(u) - \theta v \rangle dt. \quad (2.76)$$

Go back to (2.69) and pass to the limit as $m, n \rightarrow \infty$ to obtain

$$\int_0^T \left\langle \theta \frac{\partial v}{\partial t}, T_k(u - \theta \nabla v) \right\rangle dt + \int_Q a(x, t, \nabla u) \nabla T_k(u - \theta v) dx dt \leq \int_Q f T_k(u - \theta v) dx dt. \quad (2.77)$$

Letting now θ tend to 1, we get

$$\int_0^T \left\langle \frac{\partial v}{\partial t}, T_k(u - v) \right\rangle dt + \int_Q a(x, t, \nabla u) \nabla T_k(u - v) dx dt \leq \int_Q f T_k(u - v) dx dt, \quad (2.78)$$

which completes the proof. $\square$

Remark 2.4. The same technique allows us to prove an existence result for solutions of the following parabolic inequalities:

$$\begin{aligned} q_-(x) &\leq u(x, t) \leq q_+(x) \quad \text{a.e. in } Q, \\ T_k(u) &\in L^p(0, T; W_0^{1,p}(\Omega)), \\ \int_0^T \left\langle \frac{\partial v}{\partial t}, T_k(u - v) \right\rangle dt + \int_Q a(x, t, \nabla u) \nabla T_k(u - v) dx dt + \int_Q H(x, t, u, \nabla u) T_k(u - v) dx dt \\ &\leq \int_Q f T_k(u - v) dx dt, \quad \forall v \in \mathcal{H} \cap D \cap L^\infty(Q), \end{aligned} \quad (2.79)$$

where H is a given Carathéodory function satisfying, for all $(s, \zeta) \in \mathbb{R} \times \mathbb{R}^N$ and a.e. $(x, t) \in Q$, the following conditions:

$$\begin{aligned} |H(x, t, s, \zeta)| &\leq \lambda(|s|)(\delta(x, t) + |\zeta|^p), \\ H(x, t, s, \zeta)s &\geq 0, \end{aligned} \quad (2.80)$$

with $\lambda : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ is a continuous increasing function and $\delta(x, t)$ is a given positive function in $L^1(Q)$.

Acknowledgments

This work has been supported by LERMA (EMI), LMA (FSR), and Volkswagen Foundation Grant no. I/79315.

References

- [1] L. Boccardo, F. Murat, and J.-P. Puel, *Existence of bounded solutions for nonlinear elliptic unilateral problems*, Annali di Matematica Pura ed Applicata **152** (1988), no. 1, 183–196.
- [2] F. Donati, *A penalty method approach to strong solutions of some nonlinear parabolic unilateral problems*, Nonlinear Analysis **6** (1982), no. 6, 585–597.
- [3] R. Landes, *On the existence of weak solutions for quasilinear parabolic initial-boundary value problems*, Proceedings of the Royal Society of Edinburgh. Section A. Mathematics **89** (1981), no. 3–4, 217–237.
- [4] R. Landes and V. Mustonen, *A strongly nonlinear parabolic initial-boundary value problem*, Arkiv för Matematik **25** (1987), no. 1, 29–40.
- [5] J.-L. Lions, *Quelques méthodes de résolution des problèmes aux limites non linéaires*, Dunod, Paris; Gauthier-Villars, Paris, 1969.
- [6] D. Meskine and A. Elmahi, *On the limit of some nonlinear parabolic problems*, Archives of Inequalities and Applications **2** (2004), no. 4, 499–515.
- [7] M. C. Palmeri, *Homographic approximation for some nonlinear parabolic unilateral problems*, Journal of Convex Analysis **7** (2000), no. 2, 353–373.
- [8] A. Porretta, *Existence results for nonlinear parabolic equations via strong convergence of truncations*, Annali di Matematica Pura ed Applicata **177** (1999), 143–172.
- [9] M. Rudd, *Nonlinear constrained evolution in Banach spaces*, Ph.D. thesis, University of Utah, Utah, 2003.
- [10] ———, *Weak and strong solvability of parabolic variational inequalities in Banach spaces*, Journal of Evolution Equations **4** (2004), no. 4, 497–517.
- [11] J. Simon, *Compact sets in the space $L^p(0, T; B)$* , Annali di Matematica Pura ed Applicata **146** (1987), no. 1, 65–96.

K. Azelmat: LMA, GAN, Département de Mathématiques et d’Informatique, Faculté des Sciences, Université Mohammed V Agdal, Avenue Ibn Battouta, Rabat BP 1014, Morocco
E-mail address: azelmat@fsr.ac.ma

M. Kbiri Alaoui: LERMA, Ecole Mohammadia d’Ingénieurs, Université Mohammed V Agdal, Avenue Ibn Sina, Agdal, Rabat BP 765, Morocco; LMA, GAN, Département de Mathématiques et d’Informatique, Faculté des Sciences, Université Mohammed V Agdal, Avenue Ibn Battouta, Rabat BP 1014, Morocco
E-mail address: mka_la@yahoo.fr

D. Meskine: LERMA, Ecole Mohammadia d’Ingénieurs, Université Mohammed V Agdal, Avenue Ibn Sina, Agdal, Rabat BP 765, Morocco; LMA, GAN, Département de Mathématiques et d’Informatique, Faculté des Sciences, Université Mohammed V Agdal, Avenue Ibn Battouta, Rabat BP 1014, Morocco
E-mail address: driss.meskine@laposte.net

A. Souissi: LMA, GAN, Département de Mathématiques et d’Informatique, Faculté des Sciences, Université Mohammed V Agdal, Avenue Ibn Battouta, Rabat BP 1014, Morocco
E-mail address: souissi@fsr.ac.ma