

GENERALIZED LIFTING MODULES

YONGDUO WANG AND NANQING DING

Received 6 March 2006; Accepted 12 March 2006

We introduce the concepts of lifting modules and (quasi-)discrete modules relative to a given left module. We also introduce the notion of SSRS-modules. It is shown that (1) if M is an amply supplemented module and $0 \rightarrow N' \rightarrow N \rightarrow N'' \rightarrow 0$ an exact sequence, then M is N -lifting if and only if it is N' -lifting and N'' -lifting; (2) if M is a Noetherian module, then M is lifting if and only if M is R -lifting if and only if M is an amply supplemented SSRS-module; and (3) let M be an amply supplemented SSRS-module such that $\text{Rad}(M)$ is finitely generated, then $M = K \oplus K'$, where K is a radical module and K' is a lifting module.

Copyright © 2006 Hindawi Publishing Corporation. All rights reserved.

1. Introduction and preliminaries

Extending modules and their generalizations have been studied by many authors (see [2, 3, 8, 7]). The motivation of the present discussion is from [2, 8], where the concepts of extending modules and (quasi-)continuous modules with respect to a given module and CESS-modules were studied, respectively. In this paper, we introduce the concepts of lifting modules and (quasi-)discrete modules relative to a given module and SSRS-modules. It is shown that (1) if $0 \rightarrow N' \rightarrow N \rightarrow N'' \rightarrow 0$ is an exact sequence and M an amply supplemented module, then M is N -lifting if and only if it is both N' -lifting and N'' -lifting; (2) if M is a Noetherian module, then M is lifting if and only if M is R -lifting if and only if M is an amply supplemented SSRS-module; and (3) let M be an amply supplemented SSRS-module such that $\text{Rad}(M)$ is finitely generated, then $M = K \oplus K'$, where K is a radical module and K' is a lifting module.

Throughout this paper, R is an associative ring with identity and all modules are unital left R -modules. We use $N \leq M$ to indicate that N is a submodule of M . As usual, $\text{Rad}(M)$ and $\text{Soc}(M)$ stand for the Jacobson radical and the socle of a module M , respectively.

Let M be a module and $S \leq M$. S is called *small* in M (notation $S \ll M$) if $M \neq S + T$ for any proper submodule T of M . Let N and L be submodules of M , N is called a *supplement* of L in M if $N + L = M$, and N is minimal with respect to this property. Equivalently,

2 Generalized lifting modules

$M = N + L$ and $N \cap L \ll N$. N is called a *supplement submodule* if N is a supplement of some submodule of M . M is called an *amply supplemented* module if for any two submodules A and B of M with $A + B = M$, B contains a supplement of A . M is called a *weakly supplemented module* (see [5]) if for each submodule A of M there exists a submodule B of M such that $M = A + B$ and $A \cap B \ll M$. Let $B \leq A \leq M$. If $A/B \ll M/B$, then B is called a *coessential submodule* of A and A is called a *coessential extension* of B in M . A submodule A of M is called *coclosed* if A has no proper coessential submodules in M . Following [5], B is called an *s-closure* of A in M if B is a coessential submodule of A and B is coclosed in M .

Let M be a module. M is called a *lifting module* (or satisfies (D_1)) (see [9]) if for every submodule A of M , there exists a direct summand K of M such that $K \leq A$ and $A/K \ll M/K$, equivalently, M is amply supplemented and every supplement submodule of M is a direct summand. M is called *discrete* if M is lifting and has the following condition.

(D_2) If $A \leq M$ such that M/A is isomorphic to a direct summand of M , then A is a summand of M .

M is called *quasidiscrete* if M is lifting and has the following condition:

(D_3) For each pair of direct summands A and B of M with $A + B = M$, $A \cap B$ is a direct summand of M . For more details on these concepts, see [9].

LEMMA 1.1 (see [12, 19.3]). *Let M be a module and $K \leq L \leq M$.*

- (1) $L \ll M$ if and only if $K \ll M$ and $L/K \ll M/K$.
- (2) If M' is a module and $\phi : M \rightarrow M'$ a homomorphism, then $\phi(L) \ll M'$ whenever $L \ll M$.

LEMMA 1.2 (see Lemma 1.1 in [5]). *Let M be a weakly supplemented module and $N \leq M$. Then the following statements are equivalent.*

- (1) N is a supplement submodule of M .
- (2) N is coclosed in M .
- (3) For all $X \leq N$, $X \ll M$ implies $X \ll N$.

LEMMA 1.3 (see Proposition 1.5 in [5]). *Let M be an amply supplemented module. Then every submodule of M has an s-closure.*

LEMMA 1.4 (see [12, 41.7]). *Let M be an amply supplemented module. Then every coclosed submodule of M is amply supplemented.*

2. Relative lifting modules

To define the concepts of relative lifting and (quasi-)discrete modules, we dualize the concepts of relative extending and (quasi-)continuous modules introduced in [8] in this section. We start with the following.

Let N and M be modules. We define the family

$$\$(N, M) = \left\{ A \leq M \mid \exists X \leq N, \exists f \in \text{Hom}(X, M), \ni \frac{A}{f(X)} \ll \frac{M}{f(X)} \right\}. \quad (2.1)$$

PROPOSITION 2.1. $\$(N, M)$ is closed under small submodules, isomorphic images, and coessential extensions.

Proof. We only show that $\$(N, M)$ is closed under coessential extensions. Let $A \in \$(N, M)$, $A \leq A' \leq M$, and $A'/A \ll M/A$. There exist $X \leq N$ and $f \in \text{Hom}(X, M)$ such that $f(X) \leq A$ and $A/f(X) \ll M/f(X)$ since $A \in \$(N, M)$. Note that $A'/A \ll M/A$, so $A'/f(X) \ll M/f(X)$ by Lemma 1.1(1). Thus $A' \in \$(N, M)$. \square

LEMMA 2.2. Let $A \in \$(N, M)$ and A be coclosed in M . Then $B \in \$(N, M)$ for any submodule B of A .

Proof. There exist $X \leq N$ and $f \in \text{Hom}(X, M)$ such that $f(X) \leq A$ and $A/f(X) \ll M/f(X)$ by hypothesis. Since A is coclosed in M , $f(X) = A$. Let B be any submodule of A and $Y = f^{-1}(B) \leq X \leq N$. Then $f|_Y : Y \rightarrow M$ is a homomorphism such that $f|_Y(Y) = B$ for $f(X) = A$. Clearly $B/f|_Y(Y) \ll M/f|_Y(Y)$. Therefore $B \in \$(N, M)$. \square

LEMMA 2.3. Let $C \leq A \leq B \leq M$ and A be a coessential submodule of B . If C is an s -closure of A , then it is also an s -closure of B .

Proof. It is clear by Lemma 1.1(1). \square

PROPOSITION 2.4. Let M be an amply supplemented module. Then every A in $\$(N, M)$ has an s -closure \bar{A} in $\$(N, M)$.

Proof. Since $A \in \$(N, M)$, there exist $X \leq N$ and $f \in \text{Hom}(X, M)$ such that $A/f(X) \ll M/f(X)$. Note that M is amply supplemented, and so $f(X)$ has an s -closure \bar{A} in M by Lemma 1.3. Thus \bar{A} is also an s -closure of A by Lemma 2.3. The verification for $\bar{A} \in \$(N, M)$ is analogous to that for $B \in \$(N, M)$ in Lemma 2.2. \square

Let N be a module. Consider the following conditions for a module M .

$(\$(N, M)-D_1)$ For every submodule $A \in \$(N, M)$, there exists a direct summand K of M such that $K \leq A$ and $A/K \ll M/K$.

$(\$(N, M)-D_2)$ If $A \in \$(N, M)$ such that M/A is isomorphic to a direct summand of M , then A is a direct summand of M .

$(\$(N, M)-D_3)$ If A and L are direct summands of M with $A \in \$(N, M)$ and $A + L = M$, then $A \cap L$ is a direct summand of M .

Definition 2.5. Let N be a module. A module M is said to be N -lifting, N -discrete, or N -quasidiscrete if M satisfies $\$(N, M)-D_1$, $\$(N, M)-D_1$ and $\$(N, M)-D_2$ or $\$(N, M)-D_1$ and $\$(N, M)-D_3$, respectively.

One easily obtains the hierarchy: M is N -discrete $\Rightarrow M$ is N -quasidiscrete $\Rightarrow M$ is N -lifting. Clearly, the notion of relative discreteness generalizes the concept of discreteness. For any module N , lifting modules are N -lifting. But the converse is not true as shown in the following examples.

Example 2.6. Since, for any module M , $\$(0, M) = \{A \mid A \ll M\}$ and 0 is a direct summand of M such that $A/0 \ll M/0$ for any $A \in \$(0, M)$, all modules are 0 -lifting. However, the \mathbb{Z} -module $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/8\mathbb{Z}$ is not lifting since the supplement submodule $\langle (1, 2) \rangle$

4 Generalized lifting modules

$\langle\langle(1,2)\rangle\rangle$ is a supplement of $\langle\langle(1,1)\rangle\rangle$ and is not a direct summand of it though it is amply supplemented.

Example 2.7. Let M be a module with zero socle and S a simple module. Then M is S -lifting since $\$(S, M)$ is a family only containing all small submodules of M . So all torsion-free \mathbb{Z} -modules are S -lifting for any simple \mathbb{Z} -module S (see [12, Exercise 21.17]). In particular, ${}_{\mathbb{Z}}\mathbb{Z}$ and ${}_{\mathbb{Z}}\mathbb{Q}$ are S -lifting for any simple \mathbb{Z} -module, but each one is not a lifting module.

LEMMA 2.8. *Let M be a module. Then $\$(M, M) = \{A \mid A \leq M\} = \bigcup_{N \in R\text{-Mod}} \(N, M) , where $R\text{-Mod}$ denotes the category of left R -module.*

Proof. It is straight forward. □

PROPOSITION 2.9. *Let M be a module. Then M is lifting or (quasi-)discrete if and only if M is M -lifting or M -(quasi-)discrete if and only if M is N -lifting or N -(quasi-)discrete for any module N .*

Proof. It is clear by Lemma 2.8. □

PROPOSITION 2.10. *Let M be an amply supplemented module. Then the condition $\$(N, M)-D_1$ is inherited by coclosed submodules of M .*

Proof. Let M satisfy $\$(N, M)-D_1$ and H be a coclosed submodule of M . H is amply supplemented by Lemma 1.4. For any $A \in \$(N, H)$, A has an s -closure $\bar{A} \in \$(N, H)$ in H by Proposition 2.4. Since $\bar{A} \in \$(N, H) \subseteq \(N, M) and M satisfies $\$(N, M)-D_1$, there is a direct summand K of M such that $K \leq \bar{A}$ and $\bar{A}/K \ll M/K$. By Lemma 1.2, $\bar{A}/K \ll H/K$. Now $\bar{A} = K$ since \bar{A} is coclosed in H . Thus H satisfies $\$(N, H)-D_1$. □

COROLLARY 2.11. *Let M be an amply supplemented module. Then the condition $\$(N, M)-D_1$ is inherited by direct summands of M .*

PROPOSITION 2.12. *Let M be an amply supplemented module. Then $\$(N, M)-D_i$ ($i = 2, 3$) is inherited by direct summands of M .*

Proof. (1) Let M satisfy $\$(N, M)-D_2$ and H be a direct summand of M . We will show that H satisfies $\$(N, H)-D_2$.

Let $A \in \$(N, H) \subseteq \(N, M) and H/A is isomorphic to a direct summand of H . Since H is a direct summand of M , there exists $H' \leq M$ such that $M = H \oplus H'$. Thus $M/A = (H \oplus H')/A \simeq (H/A) \oplus H'$, and so M/A is isomorphic to a direct summand of M . A is a direct summand of M since M satisfies $\$(N, M)-D_2$, and hence A is a direct summand of H .

(2) Let $A \in \$(N, H) \subseteq \(N, M) and A, L be direct summands of H with $A + L = H$. We will show that $A \cap L$ is a direct summand of H . Since H is a direct summand of M , there exists $H' \leq M$ such that $M = H \oplus H'$. Thus $M = (A + L) \oplus H' = A + (L \oplus H')$. Now $A \cap (L \oplus H')$ is a direct summand of M since M satisfies $\$(N, M)-D_3$. Note that $A \cap (L \oplus H') = A \cap L$, so $A \cap L$ is a direct summand of H . □

THEOREM 2.13. *Let M be an amply supplemented module and $A \in \$(N, M)$ a direct summand of M . If M is N -(quasi-)discrete, then A is (quasi-)discrete.*

Proof. The proof follows from Lemma 2.2, Corollary 2.11, and Proposition 2.12. \square

PROPOSITION 2.14. *Let $0 \rightarrow N' \rightarrow N \rightarrow N'' \rightarrow 0$ be an exact sequence. Then $\$(N', M) \cup \$(N'', M) \subseteq \$(N, M)$. Therefore, if M is N -lifting (resp., (quasi-)discrete), then M is N' -lifting and N'' -lifting (resp., (quasi-)discrete).*

Proof. Without loss of generality we can assume that $N' \leq N$ and $N'' = N/N'$. By definition, $N' \leq N$ implies $\$(N', M) \subseteq \(N, M) . Next, let $A_2 \in \$(N'', M)$. Then there exist $X \leq N'' = N/N'$ and $f \in \text{Hom}(X, M)$ such that $A_2/f(X) \ll M/f(X)$. Write $X = Y/N'$, $Y \leq N$ and let $\delta : Y \rightarrow Y/N'$ be the canonical homomorphism. It is clear that $g = f\delta \in \text{Hom}(Y, M)$ and $g(Y) = f(X)$, hence $A_2/g(Y) \ll M/g(Y)$. Thus $A_2 \in \$(N, M)$. Therefore $\$(N', M) \cup \$(N'', M) \subseteq \$(N, M)$. The rest is obvious. \square

Dual to [8, Proposition 2.7], we have the following.

THEOREM 2.15. *Let $0 \rightarrow N' \rightarrow N \rightarrow N'' \rightarrow 0$ be an exact sequence and M an amply supplemented module. Then M is N -lifting if and only if it is both N' -lifting and N'' -lifting.*

Proof. Let M be N -lifting. Then it is both N' -lifting and N'' -lifting by Proposition 2.14. Conversely suppose that M is both N' -lifting and N'' -lifting. For any submodule $A \in \$(N, M)$, A has an s -closure $\bar{A} \in \$(N, M)$ by Proposition 2.4. Since $\bar{A} \in \$(N, M)$, there exist $X \leq N$ and $f \in \text{Hom}(X, M)$ such that $\bar{A}/f(X) \ll M/f(X)$. Since \bar{A} is coclosed in M , $f(X) = \bar{A}$. Write $Y = X \cap N' \leq N'$ and $f|_Y : Y \rightarrow M$ is a homomorphism, then $f(Y) \leq f(X) = \bar{A}$. Let $\overline{f(Y)}$ be an s -closure of $f(Y)$ in \bar{A} (for \bar{A} is amply supplemented). Thus we conclude that $f(Y)/\overline{f(Y)} \ll M/\overline{f(Y)}$ and $\overline{f(Y)} \in \$(N', M)$. Since M is N' -lifting, there exists a direct summand K of M such that $\overline{f(Y)}/K \ll M/K$. It is easy to see $\overline{f(Y)}$ is coclosed in M , hence $\overline{f(Y)} = K$ is a direct summand of M . Write $M = \overline{f(Y)} \oplus K'$, $K' \leq M$ and $\bar{A} = \bar{A} \cap M = \overline{f(Y)} \oplus (\bar{A} \cap K')$. Define $h : W = (X + N')/N' \rightarrow M$ by $h(x + N') = \pi f(x)$, where $\pi : \bar{A} \rightarrow \bar{A} \cap K'$ denotes the canonical projection. It is clear that $h(W) = \bar{A} \cap K'$, thus $(\bar{A} \cap K')/h(W) \ll M/h(W)$, and hence $(\bar{A} \cap K') \in \$(N'', M)$. Since M is N'' -lifting, there exists a direct summand K'' of M such that $(\bar{A} \cap K')/K'' \ll M/K''$. Since $\bar{A} \cap K'$ is coclosed in M , $\bar{A} \cap K' = K''$. Now $\bar{A} \cap K'$ is a direct summand of K' . Thus \bar{A} is a direct summand of M . It follows that M is N -lifting. \square

COROLLARY 2.16. *Let M be an amply supplemented module. If M is N_i -lifting for $i = 1, 2, \dots, n$ and $N = \bigoplus_i^n N_i$, then M is N -lifting.*

COROLLARY 2.17. *Let M be an amply supplemented module. Then M is lifting if and only if M is N -lifting and M/N -lifting for every submodule N of M if and only if M is N -lifting and M/N -lifting for some submodule N of M .*

Recall that a module M is said to be *distributive* if $N \cap (K + L) = (N \cap K) + (N \cap L)$ for all submodules N, K, L of M . A module M has SSP (see [4]) if the sum of any pair of direct summands of M is a direct summand of M .

COROLLARY 2.18. *Let $0 \rightarrow N' \rightarrow N \rightarrow N'' \rightarrow 0$ be an exact sequence and let M be a distributive and amply supplemented module with SSP. If M is both N' -quasidiscrete and N'' -quasidiscrete, then M is N -quasidiscrete.*

6 Generalized lifting modules

Proof. We only need to show that M satisfies $\$(N, M)-D_3$ when M satisfies $\$(N', M)-D_3$ and $\$(N'', M)-D_3$ by Theorem 2.15. Let $A \in \$(N, M)$ and A, H be direct summands of M with $A + H = M$. We know that $A = A_1 \oplus A_2$, where $A_1 \in \$(N', M)$, $A_2 \in \$(N'', M)$ from the proof of Theorem 2.15. Since M is a distributive module with SSP, $A_1 \cap H$ and $A_2 \cap H$ are direct summands of M . This implies that $A \cap H$ is a direct summand of M . Thus M satisfies $\$(N, M)-D_3$. \square

3. SSRS-modules

In [2], a module is called a *CESS-module* if every complement with essential socle is a direct summand. As a dual of CESS-modules, the concept of SSRS-modules is given in this section. It is proven that: (1) let M be an amply supplemented SSRS-module such that $\text{Rad}(M)$ is finitely generated, then $M = K \oplus K'$, where K is a radical module and K' is a lifting module; (2) let M be a finitely generated amply supplemented module, then M is an SSRS-module if and only if M/K is a lifting module for every coclosed submodule K of M .

Definition 3.1. A module is called an *SSRS-module* if every supplement with small radical is a direct summand.

Lifting modules are SSRS-modules, but the converse is not true. For example, ${}_Z\mathbb{Z}$ is an SSRS-module which is not a lifting module.

PROPOSITION 3.2. *Let M be an SSRS-module. Then any direct summand of M is an SSRS-module.*

Proof. Let K be a direct summand of M and N a supplement submodule of K such that $\text{Rad}(N) \ll N$. Let L be a supplement of N in K , that is, $N + L = K$ and $N \cap L \ll N$. Since K is a direct summand of M , there exists $K' \leq M$ such that $M = K \oplus K'$. Note that $M = N + (L \oplus K')$ and $N \cap (L \oplus K') = N \cap L \ll N$. Therefore N is a supplement of $L \oplus K'$ in M . Thus N is a direct summand of M since M is an SSRS-module. So N is a direct summand of K . The proof is complete. \square

PROPOSITION 3.3. *Let M be a weakly supplemented SSRS-module and K a coclosed submodule of M . Then K is an SSRS-module.*

Proof. It follows from the assumption and [4, Lemma 2.6(3)]. \square

PROPOSITION 3.4. *Let M be an amply supplemented module. Then M is an SSRS-module if and only if for every submodule N with small radical, there exists a direct summand K of M such that $K \leq N$ and $N/K \ll M/K$.*

Proof. “ \Leftarrow .” Let N be a supplement submodule with small radical. By assumption, there exists a direct summand K of M such that $K \leq N$ and $N/K \ll M/K$. Since N is coclosed in M , $N = K$. Thus N is a direct summand of M .

“ \Rightarrow .” Let $N \leq M$ with $\text{Rad}(N) \ll N$. There exists an s -closure \overline{N} of N since M is amply supplemented. Since $\text{Rad}(N) \ll M$ (for $\text{Rad}(N) \ll N$) and $\text{Rad}(\overline{N}) \leq \text{Rad}(N)$,

$\text{Rad}(\overline{N}) \ll \overline{N}$ and \overline{N} is a supplement submodule by Lemma 1.2. Therefore \overline{N} is a direct summand of M by assumption. This completes the proof. \square

COROLLARY 3.5. *Let M be an amply supplemented SSRS-module. Then every simple submodule of M is either a direct summand or a small submodule of M .*

PROPOSITION 3.6. *Let M be an amply supplemented module. Then M is an SSRS-module if and only if for every submodule N of M , every s -closure of N with small radical is a lifting module and a direct summand of M .*

Proof. It is straight forward. \square

PROPOSITION 3.7. *Let M be an amply supplemented SSRS-module. Then $M = K \oplus K'$, where K is semisimple and K' has small socle.*

Proof. For $\text{Soc}(M)$, there exists a direct summand K of M such that $\text{Soc}(M)/K \ll M/K$ by Proposition 3.4. It is easy to see that K is semisimple. Since K is a direct summand of M , there exists $K' \leq M$ such that $M = K \oplus K'$. Note that $\text{Soc}(M) = \text{Soc}(K) \oplus \text{Soc}(K')$. So $\text{Soc}(M)/K = (K \oplus \text{Soc}(K'))/K \ll M/K = (K \oplus K')/K$. Thus $\text{Soc}(K') \ll K'$. \square

Recall that a module M is called a *radical module* if $\text{Rad}(M) = M$. Dual to [2, Theorem 2.6], we have the following.

THEOREM 3.8. *Let M be an amply supplemented SSRS-module such that $\text{Rad}(M)$ is finitely generated. Then $M = K \oplus K'$, where K is a radical module and K' is a lifting module.*

Proof. $\text{Rad}(\text{Rad}(M)) \ll \text{Rad}(M)$ since $\text{Rad}(M)$ is finitely generated. There exists a direct summand K of M such that $\text{Rad}(M)/K \ll M/K$ by Proposition 3.4. Since K is a direct summand of M , there exists $K' \leq M$ such that $M = K \oplus K'$. Note that $\text{Rad}(M) = \text{Rad}(K) \oplus \text{Rad}(K')$. Therefore $K = K \cap \text{Rad}(M) = \text{Rad}(K)$ and $\text{Rad}(M)/K = (\text{Rad}(K) \oplus \text{Rad}(K'))/K \ll M/K = (K \oplus K')/K$. Thus $\text{Rad}(K) = K$ and $\text{Rad}(K') \ll K'$.

Next, we show that K' is a lifting module. K' is amply supplemented since it is a direct summand of M . So we only prove that every supplement submodule of K' is a direct summand of K' . Let N be a supplement submodule of K' . By Lemma 1.2 and $\text{Rad}(K') \ll K'$, we know that $\text{Rad}(N) \ll N$. N is a direct summand of K' since K' is an SSRS-module by Proposition 3.2. The proof is complete. \square

COROLLARY 3.9. *Let M be an amply supplemented module with small radical. Then M is an SSRS-module if and only if M is a lifting module.*

THEOREM 3.10. *Let M be a finitely generated amply supplemented module. Then the following statements are equivalent.*

- (1) M is an SSRS-module.
- (2) M is a lifting module.
- (3) M/K is a lifting module for every coclosed submodule K of M .

Proof. (1) \Leftrightarrow (2) follows from Corollary 3.9.

(3) \Rightarrow (1) is clear.

(1) \Rightarrow (3) we only prove that any supplement submodule of M/K is a direct summand. Let A/K be a supplement submodule of M/K . A is coclosed in M since A/K is coclosed in

8 Generalized lifting modules

M/K and K is coclosed in M . $\text{Rad}(A) \ll A$ since M is finitely generated and A is coclosed in M . A is a direct summand of M by assumption. Thus A/K is a direct summand of M/K . \square

LEMMA 3.11. *Let M be a module. Then the following statements are equivalent.*

- (1) *For every cyclic submodule N of M , there exists a direct summand K of M such that $K \leq N$ and $N/K \ll M/K$.*
- (2) *For every finitely generated submodule N of M , there exists a direct summand K of M such that $K \leq N$ and $N/K \ll M/K$.*

Proof. See [12, 41.13]. \square

COROLLARY 3.12. *Let M be a Noetherian module. Then the following statements are equivalent.*

- (1) *M is R -lifting.*
- (2) *M is F -lifting, for any free module F .*
- (3) *M is lifting.*
- (4) *M is an amply supplemented SSRS-module.*

Proof. It is easy to see that $\$(R, M)$ and $\$(F, M)$ are closed under cyclic submodules. The rest follows immediately from Theorem 3.10 and Lemma 3.11. \square

COROLLARY 3.13. *Let R be a left perfect (semiperfect) ring. Then every SSRS-module (finitely generated SSRS-module) is a lifting module.*

Proof. It follows from the fact that every module over a left perfect ring has small radical, [11, Theorems 1.6 and 1.7] and Corollary 3.9. \square

A module M is *uniserial* (see [6]) if its submodules are linearly ordered by inclusion and it is *serial* if it is a direct sum of uniserial submodules. A ring R is *right (left) serial* if the right (left) R -module R_R (${}_R R$) is serial and it is serial if it is both right and left serial.

COROLLARY 3.14. *The following statements are equivalent for a ring R with radical J .*

- (1) *R is an artinian serial ring and $J^2 = 0$.*
- (2) *R is a left semiperfect ring and every finitely generated module is an SSRS-module.*
- (3) *R is a left perfect ring and every module is an SSRS-module.*

Proof. It holds by [6, Theorem 3.15], [10, Theorem 1 and Proposition 2.13], and Corollary 3.13. \square

References

- [1] F. W. Anderson and K. R. Fuller, *Rings and Categories of Modules*, Springer, New York, 1974.
- [2] C. Çelik, A. Harmancı, and P. F. Smith, *A generalization of CS-modules*, *Communications in Algebra* **23** (1995), no. 14, 5445–5460.
- [3] N. V. Dung, D. V. Huynh, P. F. Smith, and R. Wisbauer, *Extending Modules*, Pitman Research Notes in Mathematics Series, vol. 313, Longman Scientific & Technical, Harlow, 1994.
- [4] L. Ganesan and N. Vanaja, *Modules for which every submodule has a unique coclosure*, *Communications in Algebra* **30** (2002), no. 5, 2355–2377.
- [5] D. Keskin, *On lifting modules*, *Communications in Algebra* **28** (2000), no. 7, 3427–3440.

- [6] D. Keskin, P. F. Smith, and W. M. Xue, *Rings whose modules are \oplus -supplemented*, Journal of Algebra **218** (1999), no. 2, 470–487.
- [7] Z. K. Liu, *On X -extending and X -continuous modules*, Communications in Algebra **29** (2001), no. 6, 2407–2418.
- [8] S. R. López-Permouth, K. Oshiro, and S. T. Rizvi, *On the relative (quasi-)continuity of modules*, Communications in Algebra **26** (1998), no. 11, 3497–3510.
- [9] S. H. Mohamed and B. J. Müller, *Continuous and Discrete Modules*, London Mathematical Society Lecture Note Series, vol. 147, Cambridge University Press, Cambridge, 1990.
- [10] N. Vanaja and V. M. Purav, *Characterisations of generalised uniserial rings in terms of factor rings*, Communications in Algebra **20** (1992), no. 8, 2253–2270.
- [11] K. Varadarajan, *Modules with supplements*, Pacific Journal of Mathematics **82** (1979), no. 2, 559–564.
- [12] R. Wisbauer, *Foundations of Module and Ring Theory*, Algebra, Logic and Applications, vol. 3, Gordon and Breach, Pennsylvania, 1991.

Yongduo Wang: Department of Applied Mathematics, Lanzhou University of Technology,
Lanzhou 730050, China
E-mail address: ydwang333@sohu.com

Nanqing Ding: Department of Mathematics, Nanjing University, Nanjing 210093, China
E-mail address: nqding@nju.edu.cn



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

