ON THE MEAN VALUE PROPERTY OF SUPERHARMONIC AND SUBHARMONIC FUNCTIONS

ROBERT DALMASSO

Received 19 October 2005; Revised 25 January 2006; Accepted 16 February 2006

We prove a converse of the mean value property for superharmonic and subharmonic functions. The case of harmonic functions was treated by Epstein and Schiffer.

Copyright © 2006 Hindawi Publishing Corporation. All rights reserved.

Recall that a function u is harmonic (superharmonic, subharmonic) in an open set $U \subset \mathbb{R}^n$ ($n \geq 1$) if $u \in C^2(U)$ and $\Delta u = 0$ ($\Delta u \leq 0, \Delta u \geq 0$) on U . Denote by $H(U)$ the space of harmonic functions in U and $SH(U)$ ($sH(U)$) the subset of $C^2(U)$ consisting of superharmonic (subharmonic) functions in U . If $A \subset \mathbb{R}^n$ is Lebesgue measurable, $L^1(A)$ denotes the space of Lebesgue integrable functions on A and $|A|$ denotes the Lebesgue measure of A when A is bounded.

We also recall the mean value property of harmonic, superharmonic, and subharmonic functions in U ([2]): if $x \in U$ and $B(x, r) = \{y \in \mathbb{R}^n; \|y - x\| < r\}$, $r > 0$, is such that $\bar{B}(x, r) \subset U$, then for all $u \in H(U)$ ($SH(U), sH(U)$),

$$u(x) = (\geq, \leq) \frac{1}{|B(x, r)|} \int_{B(x, r)} u(y) dy. \quad (1)$$

Using the Lebesgue-dominated convergence theorem we see that the conclusion above holds whenever $B(x, r) \subset U$ if $u \in H(U) \cap L^1(B(x, r))$ ($SH(U) \cap L^1(B(x, r)), sH(U) \cap L^1(B(x, r))$). Epstein and Schiffer [1] proved the following converse.

THEOREM 1. *Let $\Omega \subset \mathbb{R}^n$ ($n \geq 1$) be a bounded open set. Suppose that there exists $x_0 \in \Omega$ such that*

$$u(x_0) = \frac{1}{|\Omega|} \int_{\Omega} u(x) dx \quad (2)$$

for every $u \in H(\Omega) \cap L^1(\Omega)$. Then Ω is a ball with center x_0 .

A more general result was obtained by Kuran [3]. In this note we give a proof of the following converse.

2 Mean value property of super (sub) harmonic functions

THEOREM 2. *Let $\Omega \subset \mathbb{R}^n$ ($n \geq 1$) be a bounded open set. Suppose that there exists $x_0 \in \Omega$ such that*

$$u(x_0) \geq (\leq) \frac{1}{|\Omega|} \int_{\Omega} u(x) dx \quad (3)$$

for every $u \in SH(\Omega) \cap L^1(\Omega) \setminus H(\Omega)$ ($sH(\Omega) \cap L^1(\Omega) \setminus H(\Omega)$). Then Ω is a ball with center x_0 .

Proof. Clearly it is enough to consider the case of superharmonic functions. Since Ω is bounded, there exists a largest open ball B centered at x_0 of radius r_1 which lies in Ω . The compactness of $\partial\Omega$ implies that there is some $x_1 \in \partial\Omega$ such that $\|x_1 - x_0\| = r_1$. We will show that $\Omega = B$. Define

$$h(x) = r_1^{n-2} \left(\|x - x_0\|^2 - r_1^2 \right) \|x - x_1\|^{-n} \quad (4)$$

for $x \in \mathbb{R}^n \setminus \{x_1\}$. Then $h \in H(\mathbb{R}^n \setminus \{x_1\})$ and $h(x_0) = -1$. Now let $R > r_1$ be such that $\Omega \subset B(x_0, R)$. For $k \in \mathbb{N}^*$ we set

$$u_k(x) = 1 + h(x) + \frac{1}{2nk} \left(R^2 - \|x - x_0\|^2 \right), \quad x \in \Omega. \quad (5)$$

Obviously $u_k \in C^2(\Omega)$ and $\Delta u_k = -1/k$ in Ω , hence $u_k \in SH(\Omega) \setminus H(\Omega)$. Moreover $u_k \in L^1(\Omega)$ and $u_k(x) \geq 1$ for $x \in \Omega \setminus B$. Since $1 + h \in H(\Omega) \cap L^1(\Omega)$, we have

$$0 = 1 + h(x_0) = \int_B (1 + h(x)) dx. \quad (6)$$

Now using (6) we can write

$$\begin{aligned} \frac{R^2}{2nk} = u_k(x_0) &\geq \frac{1}{|\Omega|} \int_{\Omega} u_k(x) dx = \frac{1}{|\Omega|} \int_{\Omega \setminus B} u_k(x) dx + \frac{1}{|\Omega|} \int_B u_k(x) dx \\ &= \frac{1}{|\Omega|} \int_{\Omega \setminus B} u_k(x) dx + \frac{1}{2nk|\Omega|} \int_B \left(R^2 - \|x - x_0\|^2 \right) dx \\ &\geq \frac{|\Omega \setminus B|}{|\Omega|} + \frac{\omega_n r_1^n}{2nk|\Omega|} \left(\frac{R^2}{n} - \frac{r_1^2}{n+2} \right) \\ &\geq \frac{|\Omega \setminus \bar{B}|}{|\Omega|} + \frac{\omega_n r_1^n}{2nk|\Omega|} \left(\frac{R^2}{n} - \frac{r_1^2}{n+2} \right) \end{aligned} \quad (7)$$

for all $k \in \mathbb{N}^*$, where ω_n denotes the measure of the unit sphere in $\mathbb{R}^n$. This implies that $|\Omega \setminus \bar{B}| = 0$. Then the open set $\Omega \setminus \bar{B}$ must be empty, hence $\Omega \subset \bar{B}$. Since Ω is open and $B \subset \Omega \subset \bar{B}$, we deduce that $\Omega = B$. $\square$

References

- [1] B. Epstein and M. Schiffer, *On the mean-value property of harmonic functions*, Journal d'Analyse Mathématique **14** (1965), 109–111.
- [2] D. Gilbarg and N. S. Trudinger, *Elliptic Partial Differential Equations of Second Order*, Grundlehren der Mathematischen Wissenschaften, vol. 224, Springer, Berlin, 1977.

- [3] Ü. Kuran, *On the mean-value property of harmonic functions*, The Bulletin of the London Mathematical Society **4** (1972), 311–312.

Robert Dalmaso: Laboratoire LMC-IMAG, Equipe EDP, Tour IRMA, BP 53,
38041 Grenoble Cedex 9, France

E-mail address: robert.dalmaso@imag.fr