Research Article

Further Properties of β -Pascu Convex Functions of Order α

H. Özlem Güney and Grigore Stefan Sălăgean

Received 2 March 2007; Accepted 17 April 2007

Recommended by Narendra K. Govil

We obtain several further properties of β -Pascu convex functions of order α which were recently introduced and studied by Ali et al. in (2006).

Copyright © 2007 H. Ö. Güney and G. S. Sălăgean. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

Let $\mathbb{N} := \{1, 2, \dots\}$, for $m, p \in \mathbb{N}$, $m \geq p + 1$, let $\mathcal{A}(p, m)$ be the class of all p -valent analytic functions $f(z) = z^p + \sum_{n=m}^{\infty} a_n z^n$ defined on the open unit disk $\mathbb{U} = \{z \in \mathbb{C} : |z| < 1\}$ and let $\mathcal{A} := \mathcal{A}(1, 2)$.

Let $\mathcal{T}(p, m)$ be the subclass of $\mathcal{A}(p, m)$ consisting of functions of the form

$$f(z) = z^p - \sum_{n=m}^{\infty} a_n z^n, \quad a_n \geq 0 \text{ for } n \geq m, \quad (1.1)$$

and let $\mathcal{T} := \mathcal{T}(1, 2)$.

A function $f \in \mathcal{A}(p, m)$ is β -Pascu convex function of order α if

$$\frac{1}{p} \operatorname{Re} \left\{ \frac{(1 - \beta)zf'(z) + (\beta/p)z(zf'(z))'}{(1 - \beta)f(z) + (\beta/p)zf'(z)} \right\} > \alpha \quad (\beta \geq 0, 0 \leq \alpha < 1). \quad (1.2)$$

We denote by $\mathcal{TPC}(p, m, \alpha, \beta)$ the subclass of $\mathcal{T}(p, m)$ consisting of β -Pascu convex function of order α . Clearly, $\mathcal{TS}^*(\alpha) := \mathcal{TPC}(1, 2, \alpha, 0)$ is the class of starlike functions with negative coefficients of order α and $\mathcal{TC}(\alpha) := \mathcal{TPC}(1, 2, \alpha, 1)$ is the class of convex functions with negative coefficients of order α (studied by Silverman [1]).

For the class $\mathcal{TPC}(p, m, \alpha, \beta)$, the following characterization was given by Ali et al. [2].

LEMMA 1.1. Let the function f be defined by (1.1). Then f is in the class $\mathcal{TPC}(p, m, \alpha, \beta)$ if and only if

$$\sum_{n=m}^{\infty} (n - p\alpha)[(1 - \beta)p + \beta n] a_n \leq p^2(1 - \alpha). \quad (1.3)$$

The result is sharp.

LEMMA 1.2. Let $f(z)$ be given by (1.1). If $f \in \mathcal{TPC}(p, m, \alpha, \beta)$, then

$$a_n \leq \frac{p^2(1 - \alpha)}{(n - p\alpha)[(1 - \beta)p + \beta n]} \quad (1.4)$$

with equality only for functions of the form

$$f_n(z) = z^p - \frac{p^2(1 - \alpha)}{(n - p\alpha)[(1 - \beta)p + \beta n]} z^n. \quad (1.5)$$

Many interesting properties such as coefficient estimate and distortion theorems for the class $\mathcal{TPC}(p, m, \alpha, \beta)$ were given by Ali et al. [2]. In the present sequel to these earlier works, we will derive several interesting properties and characteristic of the δ -neighborhood associated with the class $\mathcal{TPC}(p, m, \alpha, \beta)$.

2. Integral properties of the class $\mathcal{TPC}(p, m, \alpha, \beta)$

We recall the following definition of integral operator before we give integral properties of the class $\mathcal{TPC}(p, m, \alpha, \beta)$.

Let $\mathcal{I}_c : \mathcal{T}(p, m) \rightarrow \mathcal{T}(p, m)$ be integral operator defined by $g = \mathcal{I}_c(f)$, where $c \in (-p, \infty)$, $f \in \mathcal{T}(p, m)$ and

$$g(z) = \frac{c+p}{z^c} \int_0^z t^{c-1} f(t) dt. \quad (2.1)$$

We note that if $f \in \mathcal{T}(p, m)$ is a function of the form (1.1), then

$$g(z) = \mathcal{I}_c(f)(z) = z^p - \sum_{n=m}^{\infty} \frac{c+p}{c+n} a_n z^n. \quad (2.2)$$

THEOREM 2.1. Let $p, m \in \mathbb{N}$, $m \geq p+1$, $\alpha \in [0, 1)$, $\beta \in [0, \infty)$, and $c \in (-p, \infty)$. If $f \in \mathcal{TPC}(p, m, \alpha, \beta)$ and $g = \mathcal{I}_c(f)$, then $g \in \mathcal{TPC}(p, m, \lambda, \beta)$, where

$$\lambda = \lambda(p, m, \alpha, c) = 1 - \frac{(1 - \alpha)(c + p)(m - p)}{(m - p\alpha)(c + m) - (1 - \alpha)(c + p)p} \quad (2.3)$$

and $\alpha < \lambda$. The result is sharp.

Proof. From Lemma 1.1 and (2.2), we have $g \in \mathcal{TPC}(p, m, \lambda, \beta)$ if and only if

$$\sum_{n=m}^{\infty} \frac{(n - p\lambda)[(1 - \beta)p + \beta n](c + p)}{p^2(1 - \lambda)(c + n)} a_n \leq 1. \quad (2.4)$$

We find the largest λ such that (2.4) holds. We note that the inequalities

$$\frac{(n - p\lambda)[(1 - \beta)p + \beta n](c + p)}{p^2(1 - \lambda)(c + n)} \leq \frac{(n - p\alpha)[(1 - \beta)p + \beta n]}{p^2(1 - \alpha)} \quad (2.5)$$

imply (2.4), because $f \in \mathcal{TPC}(p, m, \alpha, \beta)$ and satisfy (1.3). But inequalities (2.5) are equivalent to

$$\frac{(n - p\lambda)(c + p)}{(1 - \lambda)(c + n)} \leq \frac{(n - p\alpha)}{(1 - \alpha)}. \quad (2.6)$$

Since $(n - p\alpha) > p(1 - \alpha)$ and $c + n > c + p$, we obtain $\lambda \leq \lambda(p, n, \alpha, c)$, where

$$\lambda(p, n, \alpha, c) = \frac{(n - p\alpha)(c + n) - (1 - \alpha)(c + p)n}{(n - p\alpha)(c + n) - (1 - \alpha)(c + p)p}. \quad (2.7)$$

Now we show that $\lambda(p, n, \alpha, c)$ is an increasing function of n , $n \geq m$. Indeed,

$$\lambda(p, n, \alpha, c) = 1 - (1 - \alpha)(c + p)E(p, n, \alpha, c), \quad (2.8)$$

where

$$E(p, n, \alpha, c) = \frac{(n - p)}{(n - p\alpha)(c + n) - (1 - \alpha)(c + p)p}, \quad (2.9)$$

and $\lambda(p, n, \alpha, c)$ increases when n increases if and only if $E(p, n, \alpha, c)$ is a strictly decreasing function of n .

Let $h(x) = E(p, x, \alpha, c)$, $x \in [m, \infty) \subset [p + 1, \infty)$, we have

$$h'(x) = -\frac{(x - p)^2}{[(x - p\alpha)(c + x) - (1 - \alpha)(c + p)p]^2} < 0. \quad (2.10)$$

We obtained

$$\lambda = \lambda(p, m, \alpha, c) \leq \lambda(p, n, \alpha, c), \quad n \geq m. \quad (2.11)$$

The result is sharp because

$$\mathcal{I}_c(f_\alpha) = f_\lambda, \quad (2.12)$$

where

$$\begin{aligned} f_\alpha(z) &= z^p - \frac{p^2(1 - \alpha)}{(m - p\alpha)[(1 - \beta)p + \beta m]} z^m, \\ f_\lambda(z) &= z^p - \frac{p^2(1 - \lambda)}{(m - p\lambda)[(1 - \beta)p + \beta m]} z^m \end{aligned} \quad (2.13)$$

are extremal functions of $\mathcal{TPC}(p, m, \alpha, \beta)$ and $\mathcal{TPC}(p, m, \lambda, \beta)$, respectively, and $\lambda = \lambda(p, m, \alpha, c)$.

Indeed, we have

$$\mathcal{G}_c(f_\alpha(z)) = z^p - \frac{p^2(1-\alpha)(c+p)}{(m-p\alpha)[(1-\beta)p+\beta m](c+m)} z^m. \quad (2.14)$$

We deduce

$$\frac{p^2(1-\lambda)}{(m-p\lambda)} = \frac{p^2(1-\alpha)(c+p)}{(m-p\alpha)(c+m)}, \quad (2.15)$$

and this implies (2.14).

From $\lambda = 1 - (1-\alpha)(c+p)(m-p)/((m-p\alpha)(c+m) - (1-\alpha)(c+p)p)$ we obtain $\lambda < 1$ and also $\lambda > \alpha$. Indeed,

$$\begin{aligned} \lambda - \alpha &= (1-\alpha) \left\{ 1 - \frac{(c+p)(m-p)}{(m-p\alpha)(c+m) - (1-\alpha)(c+p)p} \right\} \\ &= (1-\alpha) \frac{(m-p\alpha)(m-p)}{(m-p\alpha)(c+m) - (1-\alpha)(c+p)p} > 0. \end{aligned} \quad (2.16)$$

□

3. Integral means inequalities for the class $\mathcal{TPC}(p, m, \alpha, \beta)$

An analytic function g is said to be subordinate to an analytic function f (written $g < f$) if $g(z) = f(w(z))$, $z \in \mathbb{U}$, for some analytic function w with $|w(z)| \leq |z|$. In 1925, Littlewood [3] proved the following subordination result which will be required in our present investigation.

LEMMA 3.1. *If f and g are analytic in $\mathbb{U}$ with $g < f$, then*

$$\int_0^{2\pi} |g(re^{i\theta})|^\delta d\theta \leq \int_0^{2\pi} |f(re^{i\theta})|^\delta d\theta, \quad (3.1)$$

where $\delta > 0$, $z = re^{i\theta}$, and $0 < r < 1$.

Applying Lemmas 1.1 and 3.1, we prove the following.

THEOREM 3.2. *Let $\delta > 0$. If $f \in \mathcal{TPC}(p, m, \alpha, \beta)$ and $f_m(z) = z^p - (p^2(1-\alpha)/(m-p\alpha)[(1-\beta)p+\beta m])z^m$, then for $z = re^{i\theta}$ and $0 < r < 1$,*

$$\int_0^{2\pi} |f(re^{i\theta})|^\delta d\theta \leq \int_0^{2\pi} |f_m(re^{i\theta})|^\delta d\theta. \quad (3.2)$$

Proof. Let

$$\begin{aligned} f(z) &= z^p - \sum_{n=m}^{\infty} a_n z^n, \quad a_n \geq 0, \quad n \geq m, \\ f_m(z) &= z^p - \frac{p^2(1-\alpha)}{(m-p\alpha)[(1-\beta)p+\beta m]} z^m, \end{aligned} \quad (3.3)$$

then we must show that

$$\int_0^{2\pi} \left| 1 - \sum_{n=m}^{\infty} a_n z^{n-p} \right|^\delta d\theta \leq \int_0^{2\pi} \left| 1 - \frac{p^2(1-\alpha)}{(m-p\alpha)[(1-\beta)p+\beta m]} z^{m-p} \right|^\delta d\theta. \quad (3.4)$$

By Lemma 3.1, it suffices to show that

$$1 - \sum_{n=m}^{\infty} a_n z^{n-p} < 1 - \frac{p^2(1-\alpha)}{(m-p\alpha)[(1-\beta)p+\beta m]} z^{m-p}. \quad (3.5)$$

Set

$$1 - \sum_{n=m}^{\infty} a_n z^{n-p} = 1 - \frac{p^2(1-\alpha)}{(m-p\alpha)[(1-\beta)p+\beta m]} w(z)^{m-p}. \quad (3.6)$$

From (3.6) and (1.3), we obtain

$$\begin{aligned} |w(z)|^{m-p} &= \left| \frac{(m-p\alpha)[(1-\beta)p+\beta m]}{p^2(1-\alpha)} \right| \left| \sum_{n=m}^{\infty} a_n z^{n-p} \right| \\ &\leq |z^{m-p}| \sum_{n=m}^{\infty} \frac{(n-p\alpha)[(1-\beta)p+\beta n]}{p^2(1-\alpha)} a_n \leq |z^{m-p}| \leq |z|. \end{aligned} \quad (3.7)$$

This completes the proof of the theorem. $\square$

The proof for the first derivative is similar.

THEOREM 3.3. *Let $\delta > 0$. If $f \in \mathcal{TPC}(p, m, \alpha, \beta)$ and $f_m(z) = z^p - (p^2(1-\alpha)/(m-p\alpha)[(1-\beta)p+\beta m])z^m$, then for $z = re^{i\theta}$ and $0 < r < 1$,*

$$\int_0^{2\pi} |f'(re^{i\theta})|^\delta d\theta \leq \int_0^{2\pi} |f'_m(re^{i\theta})|^\delta d\theta. \quad (3.8)$$

Proof. It suffices to show that

$$1 - \sum_{n=m}^{\infty} \frac{n}{p} a_n z^{n-p} < 1 - \frac{mp(1-\alpha)}{(m-p\alpha)[(1-\beta)p+\beta m]} z^{m-p}. \quad (3.9)$$

This follows because

$$\begin{aligned} |w(z)|^{m-p} &= \left| \sum_{n=m}^{\infty} \frac{(n-p\alpha)[(1-\beta)p+\beta n]}{p^2(1-\alpha)} a_n z^{n-p} \right| \\ &\leq |z|^{n-p} \sum_{n=m}^{\infty} \frac{(n-p\alpha)[(1-\beta)p+\beta n]}{p^2(1-\alpha)} a_n \leq |z|^{n-p} \leq |z|. \end{aligned} \quad (3.10)$$

$\square$

4. Neighborhoods of the class $\mathcal{TPC}(p, m, \alpha, \beta)$

For $f \in \mathcal{T}(p, m)$ and $\gamma \geq 0$, Frasin [4] defined

$$M_\gamma^q(f) = \left\{ g \in \mathcal{T}(p, m) : g(z) = z^p - \sum_{n=m}^{\infty} b_n z^n, \sum_{n=m}^{\infty} n^{q+1} |a_n - b_n| \leq \gamma \right\}, \quad (4.1)$$

which was called q - γ -neighborhood of f . So, for $e(z) = z$, we see that

$$M_\gamma^q(e) = \left\{ g \in \mathcal{T}(p, m) : g(z) = z^p - \sum_{n=m}^{\infty} b_n z^n, \sum_{n=m}^{\infty} n^{q+1} |b_n| \leq \gamma \right\}, \quad (4.2)$$

where q is a fixed positive integer. Note that $M_\gamma^0(f) \equiv N_\gamma(f)$ and $M_\gamma^1(f) \equiv M_\gamma(f)$. $N_\gamma(f)$ is called a γ -neighborhood of f by Ruscheweyh [5] and $M_\gamma(f)$ was defined by Silverman [6].

Now, we consider q - γ -neighborhood for function in the class $\mathcal{TPC}(p, m, \alpha, \beta)$.

THEOREM 4.1. *Let*

$$\gamma = \frac{m^{q+1} p^2 (1 - \alpha)}{(m - p\alpha)[(1 - \beta)p + \beta m]}, \quad (4.3)$$

then $\mathcal{TPC}(p, m, \alpha, \beta) \subset M_\gamma^q(e)$.

Proof. If $f \in \mathcal{TPC}(p, m, \alpha, \beta)$, then

$$\sum_{n=m}^{\infty} n^{q+1} a_n \leq \frac{m^{q+1} p^2 (1 - \alpha)}{(m - p\alpha)[(1 - \beta)p + \beta m]} = \gamma. \quad (4.4)$$

This gives that $\mathcal{TPC}(p, m, \alpha, \beta) \subset M_\gamma^q(e)$. □

Putting $p = 1$, $m = 2$ and $\beta = 0$ in Theorem 4.1, we have the following.

COROLLARY 4.2. $\mathcal{TS}^*(\alpha) \subset M_\gamma^q(e)$, where $\gamma = 2^{q+1}(1 - \alpha)/(2 - \alpha)$.

Putting $p = 1$, $m = 2$, and $\beta = 1$ in Theorem 4.1, we have the following.

COROLLARY 4.3. $\mathcal{TC}(\alpha) \subset M_\gamma^q(e)$, where $\gamma = 2^q(1 - \alpha)/(2 - \alpha)$.

References

- [1] H. Silverman, "Univalent functions with negative coefficients," *Proceedings of the American Mathematical Society*, vol. 51, no. 1, pp. 109–116, 1975.
- [2] R. M. Ali, M. H. Khan, V. Ravichandran, and K. G. Subramanian, "A class of multivalent functions with negative coefficients defined by convolution," *Bulletin of the Korean Mathematical Society*, vol. 43, no. 1, pp. 179–188, 2006.
- [3] J. E. Littlewood, "On inequalities in the theory of functions," *Proceedings of the London Mathematical Society*, vol. 23, no. 1, pp. 481–519, 1925.
- [4] B. A. Frasin, "Family of analytic functions of complex order," *Acta Mathematica. Academiae Paedagogicae Nyrtregyháziensis*, vol. 22, no. 2, pp. 179–191, 2006.

- [5] S. Ruscheweyh, "Neighborhoods of univalent functions," *Proceedings of the American Mathematical Society*, vol. 81, no. 4, pp. 521–527, 1981.
- [6] H. Silverman, "Neighborhoods of classes of analytic functions," *Far East Journal of Mathematical Sciences*, vol. 3, no. 2, pp. 165–169, 1995.

H. Özlem Güney: Department of Mathematics, Faculty of Science and Arts, University of Dicle,
21280 Diyarbakir, Turkey
Email address: ozlemg@dicle.edu.tr

Grigore Stefan Sălăgean: Faculty of Mathematics and Computer Science, Babes-Bolyai University,
1 M. Kogalniceanu Street, 400084 Cluj-Napoca, Romania
Email address: salagean@math.ubbcluj.ro