Research Article

Lebesgue Measurability of Separately Continuous Functions and Separability

V. V. Mykhaylyuk

Received 4 September 2006; Accepted 22 April 2007

Recommended by Peter Johnson

A connection between the separability and the countable chain condition of spaces with L -property (a topological space X has L -property if for every topological space Y , separately continuous function $f : X \times Y \rightarrow \mathbb{R}$ and open set $I \subseteq \mathbb{R}$, the set $f^{-1}(I)$ is an F_σ -set) is studied. We show that every completely regular Baire space with the L -property and the countable chain condition is separable and constructs a nonseparable completely regular space with the L -property and the countable chain condition. This gives a negative answer to a question of M. Burke.

Copyright © 2007 V. V. Mykhaylyuk. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

A function $f : X \rightarrow \mathbb{R}$ defined on a topological space X is called a *first Baire class function* if there exists a sequence $(f_n)_{n=1}^\infty$ of continuous functions $f_n : X \rightarrow \mathbb{R}$ which converges pointwise to f on X ; and a *first Lebesgue class function* if $f^{-1}(G)$ is an F_σ -set for every open set $G \subseteq \mathbb{R}$. Standard reasons (see [1, page 394]) show that every first Baire class function is a first Lebesgue class function.

Investigations of Baire and Lebesgue classifications of separately continuous functions were started by Lebesgue in [2] and were continued in papers of many mathematicians (see [3]).

We say that a topological space X has the *B-property* (the *L-property*) if for every topological space Y each separately continuous function $f : X \times Y \rightarrow \mathbb{R}$ is a first Baire class function (a first Lebesgue class function).

It is known [4, 5] that any topological space X has the *B-property* (the *L-property*) if and only if the evaluation function $c_X : X \times C_p(X) \rightarrow \mathbb{R}$, $c_X(x, y) = y(x)$ is a first Baire

class function (a first Lebesgue class function), where $C_p(X)$ means the space of continuous on X functions with the pointwise convergence topology.

Baire and Lebesgue classifications of separately continuous function were investigated in [6]. In particular, it was shown in [6] that any completely regular space X with the B -property and the countable chain condition is separable (topological space X has a countable chain condition (CCC) if every system of disjoint open-in- X sets is at most countable). In this connection the following question arose in [6, Problem 4.6].

Question 1. Is every completely regular space X with the L -property and the countable chain condition a separable space?

In this paper, we show that if a space X is a Baire space, then Question 1 has a positive answer and construct an example which gives a negative answer to the question in general case.

2. Density of Baire spaces with the L -property

The minimal cardinal $\aleph \geq \aleph_0$ for which any system of disjoint open in a topological space X sets has the cardinality at most $\aleph$ is called a *Souslin number of X* and is denoted by $c(X)$. Note that the countable chain condition of X means that $c(X) = \aleph_0$. It is easy to see that $c(X) \leq d(X)$, where $d(X)$ is the density of X .

The following result implies that for a Baire space X Question 1 has a positive answer.

THEOREM 2.1. *Let X be a completely regular Baire space with the L -property. Then $c(X) = d(X)$.*

Proof. Since the evaluation function c_X is a first Lebesgue class function, the set $E = \{(x, y) : y(x) = 0\}$ is a G_δ -set in $X \times Y$, where $Y = C_p(X)$. Choose a sequence $(W_n)_{n=1}^\infty$ of open-in- $X \times Y$ sets W_n such that $E = \bigcap_{n=1}^\infty W_n$. Denote by y_0 the null-function on Y . For every $n \in \mathbb{N}$ and an $x \in X$ find open neighborhoods $U(x, n)$ and $V(x, n)$ of x and y_0 in X and Y , respectively, such that $U(x, n) \times V(x, n) \subseteq W_n$.

Fix an $n \in \mathbb{N}$ and show that there exists a set $A_n \subseteq X$ with $|A_n| \leq c(X) = \aleph$ such that the open set $G_n = \bigcup_{x \in A_n} U(x, n)$ is dense in X . Consider a system $\mathcal{U}$ of all open-in- X nonempty sets U such that $U \subseteq U(x, n)$ for some $x \in X$ and choose a maximal system $\mathcal{U}' \subseteq \mathcal{U}$ which consists of disjoint sets. It is clear that $|\mathcal{U}'| \leq \aleph$. For every $U \in \mathcal{U}'$ find an $x = x(U) \in X$ such that $U \subseteq U(x, n)$ and put $A_n = \{x(U) : U \in \mathcal{U}'\}$. Then $|A_n| \leq |\mathcal{U}'| \leq \aleph$. Besides, it follows from the maximality of $\mathcal{U}'$ that G_n is dense in X .

Since X is a Baire space, the set $X_0 = \bigcap_{n=1}^\infty G_n$ is dense in X . For every $n \in \mathbb{N}$ and $x \in X$ choose a finite set $B(x, n) \subseteq X$ such that $y \in V(x, n)$ for each $y \in Y$ with $y|_{B(x, n)} = y_0|_{B(x, n)}$. Put $B = \bigcup_{n \in \mathbb{N}} \bigcup_{x \in A_n} B(x, n)$. Note that $|B| \leq \aleph_0 \cdot \aleph = \aleph$.

Show that B is dense in X . Since X is a completely regular space, it is enough to prove that y_0 is a unique continuous on X function which equals to 0 at every point from B . Let $y \in Y$ be a function such that $y(b) = 0$ for every $b \in B$. Fix a point $x \in X_0$ and an integer $n \in \mathbb{N}$. Find $a \in A_n$ such that $x \in U(a, n)$. Then $B(a, n) \subseteq B$ implies $y \in V(a, n)$. Therefore, $(x, y) \in W_n$. Thus $X_0 \times \{y\} \subseteq \bigcap_{n=1}^\infty W_n = E$, that is, $y(x) = 0$ for every $x \in X_0$. Hence $y = y_0$ because X_0 is dense in X .

Thus $d(X) \leq |B| \leq c(X)$. Therefore, $c(X) = d(X)$. □

COROLLARY 2.2. *Every completely regular Baire space with the L -property and the countable chain condition is a separable space.*

3. Nonseparable spaces with the L -property and CCC

The following notion was introduced in [4], where some properties of spaces with the B -property were studied.

A topological space X with a topology τ is called *quarter-stratifiable* if there exists a function $g : \mathbb{N} \times X \rightarrow \tau$ such that

- (i) $X = \bigcup_{x \in X} g(n, x)$ for every $n \in \mathbb{N}$;
- (ii) if $x \in g(n, x_n)$ for each $n \in \mathbb{N}$, then $x_n \rightarrow x$.

The following result follows from [7, Proposition 2.1].

PROPOSITION 3.1. *Every quarter-stratifiable space X has the L -property.*

A topological space X is called σ -discrete if there exists an increasing sequence $(X_n)_{n=1}^\infty$ of closed discrete subspaces X_n of X such that $X = \bigcup_{n=1}^\infty X_n$.

PROPOSITION 3.2. *Every σ -discrete space is a quarter-stratifiable space.*

Proof. Let $(X_n)_{n=1}^\infty$ be an increasing sequence of closed discrete subspaces X_n of X such that $X = \bigcup_{n=1}^\infty X_n$. For every $n \in \mathbb{N}$ and $x \in X_n$ denote by $U(x, n)$ an open-in- X neighborhood of x such that $U(x, n) \cap X_n = \{x\}$. We define a function $g : \mathbb{N} \times X \rightarrow \tau$, where τ is the topology of X , by $g(x, n) = U(x, n)$ if $x \in X_n$ and $g(x, n) = X \setminus X_n$ if $x \notin X_n$. It is easy to see that g satisfies (i) and (ii). $\square$

Show now that Question 1 has a negative answer.

THEOREM 3.3. *There exists a completely regular nonseparable space with the L -property and with the countable chain condition.*

Proof. Let Γ_0 be a set with $|\Gamma_0| \geq \aleph_1$, let $(a_n)_{n=1}^\infty$ be a sequence of distinct points $a_n \notin \Gamma_0$, $\Gamma_n = \Gamma_0 \cup \{a_k : 1 \leq k \leq n\}$, and let $\mathcal{A}_n$ be a system of all subsets $A \subseteq \Gamma_{n-1}$ such that $|A| = n$. Denote by X_n a set of all function $x \in \{0, 1\}^\Gamma$ such that $x = \chi_{A \cup \{a_n\}}$ for some $A \in \mathcal{A}_n$, where χ_B means the characteristic function of B , and put $X = \bigcup_{n=1}^\infty X_n$.

Show that X is a σ -discrete space. For every $n \in \mathbb{N}$ put $Y_n = \bigcup_{k=1}^n X_k$. Fix an integer $n \in \mathbb{N}$ and for each $1 \leq k \leq n$ put $G_k = \{x \in X : x(a_k) = 1, x(a_i) = 0, k < i \leq n\}$. It is easy to see that $G_k \cap Y_n = X_k$. Since all spaces X_k are discrete, Y_n is discrete in X too. Besides, Y_n is closed in X . Thus, X has the L -property by Propositions 3.1 and 3.2.

Note that X is dense in $Y = \{0, 1\}^\Gamma$. Indeed, let $A \subseteq \Gamma$ be a finite set and $y : A \rightarrow \{0, 1\}$. Choosing $n \geq |A|$ with $A \subseteq \Gamma_n$ find $x \in X_{n+1}$ such that $x|_A = y$. Then $c(X) = \aleph_0$ since $c(Y) = \aleph_0$ and X is dense in Y .

It remains to note that X is nonseparable because for every separable subspace Z of X there exists a countable set $B \subseteq \Gamma$ such that $z(y) = 0$ for every $y \in \Gamma \setminus B$. $\square$

This example shows that there exists a quarter-stratifiable space which has not the B -property. Thus, Proposition 3.1 cannot be generalized for spaces with the B -property.

A family $(A_i : i \in I)$ of sets A_i is called *pointwise finite* if $\bigcap_{i \in J} A_i = \emptyset$ for each infinite set $J \subseteq I$. A cardinal

$$p(X) = \sup \{ |\mathcal{A}| : \mathcal{A} \text{ is a pointwise finite family of nonempty open-in-} X \text{ sets} \} \quad (3.1)$$

is called a *point-finite cellularity of a topological space* X . Clearly $c(X) \leq p(X)$. Besides, it is known that $p(X) = c(X)$ for each Baire space X . Therefore, the following question arises naturally from Theorem 2.1 and the fact that $p(X) = |\Gamma| > \aleph_0$ for the space X from Theorem 3.3.

Question 2. Is every completely regular space X with the L -property and $p(X) = \aleph_0$ a separable space?

References

- [1] K. Kuratowski, *Topology. Vol. I*, Mir, Moscow, Russia, 1966.
- [2] H. Lebesgue, "Sur l'approximation des fonctions," *Bulletin des Sciences Mathématiques*, vol. 22, pp. 278–287, 1898.
- [3] V. K. Maslyuchenko, O. V. Maslyuchenko, V. V. Mykhaylyuk, and O. V. Sobchuk, "Paracompactness and separately continuous mappings," in *General Topology in Banach Spaces*, pp. 147–169, Nova Science Publishers, Huntington, NY, USA, 2001.
- [4] T. O. Banakh, "(Metrically) quarter-stratifiable spaces and their applications in the theory of separately continuous functions," *Matematichnī Studii*, vol. 18, no. 1, pp. 10–28, 2002.
- [5] M. R. Burke, "Borel measurability of separately continuous functions," *Topology and Its Applications*, vol. 129, no. 1, pp. 29–65, 2003.
- [6] M. R. Burke, "Borel measurability of separately continuous functions. II," *Topology and Its Applications*, vol. 134, no. 3, pp. 159–188, 2003.
- [7] O. O. Karlova, "The first functional Lebesgue class and Baire classification of separately continuous mappings," *Naukovyj Visnyk Chernivets'kogo Universytetu. Matematika*, vol. 191–192, pp. 52–60, 2004 (Ukrainian).

V. V. Mykhaylyuk: Department of Mathematical Analysis, Chernivtsi National University,
Kotsjubyns'koho 2, 58012 Chernivtsi, Ukraine
Email address: mathan@chnu.cv.ua