Research Article

On the Generalized Ulam-Gavruta-Rassias Stability of Mixed-Type Linear and Euler-Lagrange-Rassias Functional Equations

Paisan Nakmahachalasint

Received 18 April 2007; Revised 22 April 2007; Accepted 23 April 2007

Recommended by Martin J. Bohner

In this paper, the mixed-type linear and Euler-Lagrange-Rassias functional equations introduced by J. M. Rassias is generalized to the following n -dimensional functional equation: $f(\sum_{i=1}^n x_i) + (n-2)\sum_{i=1}^n f(x_i) = \sum_{1 \leq i < j \leq n} f(x_i - x_j)$ when $n > 2$. We prove the general solutions and investigate its generalized Ulam-Gavruta-Rassias stability.

Copyright © 2007 Paisan Nakmahachalasint. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

In 1940, Ulam [1] proposed the famous Ulam stability problem of linear mappings. In 1941, Hyers [2] considered the case of approximately additive mappings $f : E \rightarrow E'$, where E and E' are Banach spaces and f satisfies *Hyers inequality* $\|f(x+y) - f(x) - f(y)\| \leq \varepsilon$ for all $x, y \in E$. It was shown that the limit $L(x) = \lim_{n \rightarrow \infty} 2^{-n} f(2^n x)$ exists for all $x \in E$ and that $L : E \rightarrow E'$ is the unique additive mapping satisfying $\|f(x) - L(x)\| \leq \varepsilon$. In 1982–1998, Rassias [3–9] generalized the result to include the following theorem.

THEOREM 1.1. *Let X be a real-normed linear space and let Y be a real-complete-normed linear space. Assume in addition that $f : X \rightarrow Y$ is an approximately additive mapping for which there exist constants $\theta \geq 0$ and $p, q \in \mathbb{R}$ such that $r = p + q \neq 1$, and f satisfies the Cauchy-Gavruta-Rassias inequality*

$$\|f(x+y) - f(x) - f(y)\| \leq \theta \|x\|^p \|y\|^q \quad (1.1)$$

for all $x, y \in X$. Then, there exists a unique additive mapping $L : X \rightarrow Y$ satisfying

$$f(x) - L(x) \leq \frac{\theta}{|2^r - 2|} \|x\|^r \quad \forall x \in X. \quad (1.2)$$

If in addition $f : X \rightarrow Y$ is a mapping such that the transformation $t \rightarrow f(tx)$ is continuous in $t \in \mathbb{R}$ for each fixed $x \in X$, then L is $\mathbb{R}$ -linear mapping.

In 2002, Rassias [10] established the Ulam stability of the following *mixed-type* functional equation:

$$f\left(\sum_{i=1}^3 x_i\right) + \sum_{i=1}^3 f(x_i) = \sum_{1 \leq i < j \leq 3} f(x_i + x_j) \quad (1.3)$$

on restricted domains. In this paper, we will generalize Rassias' work to the following n -dimensional mixed-type functional equation:

$$f\left(\sum_{i=1}^n x_i\right) + (n-2) \sum_{i=1}^n f(x_i) = \sum_{1 \leq i < j \leq n} f(x_i + x_j) \quad (1.4)$$

when $n > 2$, and will investigate its generalized Ulam-Gavruta-Rassias stability.

2. The general solution

THEOREM 2.1. *Let $n > 2$ be a positive integer, and let X and Y be vector spaces.*

A function $f : X \rightarrow Y$ satisfies the functional equation

$$f\left(\sum_{i=1}^n x_i\right) + (n-2) \sum_{i=1}^n f(x_i) = \sum_{1 \leq i < j \leq n} f(x_i + x_j) \quad (2.1)$$

if and only if the even part of f , defined by $f_e(x) = (1/2)(f(x) + f(-x))$ for all $x \in X$, satisfies the classical quadratic functional equation, which is also a special Euler-Lagrange-Rassias equation [7, 9],

$$f(x+y) + f(x-y) = 2f(x) + 2f(y), \quad (2.2)$$

and the odd part of f , defined by $f_o(x) = (1/2)(f(x) - f(-x))$ for all $x \in X$, satisfies the Cauchy functional equation

$$f(x+y) = f(x) + f(y). \quad (2.3)$$

Proof. For the *if* part of the proof, suppose that $f : X \rightarrow Y$ satisfies (2.1), we can uniquely express f as $f(x) = f_e(x) + f_o(x)$ for all $x \in X$, where the even part, f_e , and the odd part, f_o , are defined as in the theorem. We will show that f_e satisfies (2.2) and f_o satisfies (2.3).

Setting $(x_1, x_2, \dots, x_n) = (0, 0, \dots, 0)$ in (2.1), we see that $f(0) = 0$. Setting $(x_1, x_2, \dots, x_n) = (x, y, -y, 0, 0, \dots, 0)$ in (2.1), we get

$$\begin{aligned} f(x) + (n-2)(f(x) + f(y) + f(-y)) &= f(x-y) + f(x+y) \\ &+ (n-3)(f(x) + f(y) + f(-y)), \end{aligned} \quad (2.4)$$

which is simplified to

$$2f(x) + f(y) + f(-y) = f(x+y) + f(x-y) \quad (2.5)$$

for all $x, y \in X$. Replacing x and y with $-x$ and $-y$, respectively, then taking half the sum and half the difference with (2.5), we have

$$\begin{aligned} 2f_e(x) + f_e(y) + f_e(-y) &= f_e(x+y) + f_e(x-y), \\ 2f_o(x) + f_o(y) + f_o(-y) &= f_o(x+y) + f_o(x-y). \end{aligned} \quad (2.6)$$

By the evenness of f_e , we immediately see that f_e satisfies the classical quadratic functional equation given by (2.2). By the oddness of f_o , we see that $2f_o(x) = f_o(x+y) + f_o(x-y)$ which is recognized as the Jensen functional equation. Since $f_o(0) = 0$, if we put $y = x$ in the above equation, then $f(2x) = 2f(x)$. By another substitution, $(x, y) = ((x+y)/2, (x-y)/2)$, we derive the Cauchy functional equation $f_o(x+y) = f_o(x) + f_o(y)$.

Now for the *only if* part of the proof, suppose that the even part and the odd part of $f : X \rightarrow Y$ satisfy (2.2) and (2.3), respectively, that is, $f_e(x+y) + f_e(x-y) = 2f_e(x) + 2f_e(y)$ and $f_o(x+y) = f_o(x) + f_o(y)$. We will show that f satisfies (2.1). Noting that a linear combination of two solutions of (2.1) yields just another solution, we will in turn prove that each part of f satisfies (2.1).

First, consider the odd part and make use of the linearity of the Cauchy functional equation. The left-hand side of (2.1) is

$$f_o\left(\sum_{i=1}^n x_i\right) + (n-2) \sum_{i=1}^n f_o(x_i) = \sum_{i=1}^n f_o(x_i) + (n-2) \sum_{i=1}^n f_o(x_i) = (n-1) \sum_{i=1}^n f_o(x_i), \quad (2.7)$$

and the right-hand side of (2.1) is

$$\sum_{1 \leq i < j \leq n} f_o(x_i + x_j) = \sum_{1 \leq i < j \leq n} (f_o(x_i) + f_o(x_j)) = \frac{2}{n} \binom{n}{2} \sum_{i=1}^n f_o(x_i) = (n-1) \sum_{i=1}^n f_o(x_i). \quad (2.8)$$

Thus, we have established (2.1) on the odd part of f .

For the even part, we will show by mathematical induction that (2.1) holds for every positive integer n . For $n = 1$, we take $\sum_{1 \leq i < j \leq 1} f_e(x_i + x_j)$ as 0; then $f_e(x_1) + (1-2)f_e(x_1) = 0$, which is trivially true. For $n = 2$, we have $f_e(x_1 + x_2) + 0 = f_e(x_1 + x_2)$, which is again trivially true. For $n \geq 3$, we assume that (2.1) holds for every number of variables from 1 to $n-1$, that is,

$$f_e\left(\sum_{i=1}^k x_i\right) + (k-2) \sum_{i=1}^k f_e(x_i) = \sum_{1 \leq i < j \leq k} f_e(x_i + x_j) \quad (2.9)$$

for $k = 1, 2, \dots, n-1$. For each $i, j = 1, 2, \dots, n$ with $i \neq j$, we have

$$f_e(x_i - x_j) + f_e(x_i + x_j) = 2(f_e(x_i) + f_e(x_j)). \quad (2.10)$$

Then,

$$\sum_{1 \leq i < j \leq n} (f_e(x_i - x_j) + f_e(x_i + x_j)) = 2 \sum_{1 \leq i < j \leq n} (f_e(x_i) + f_e(x_j)) = \frac{4}{n} \binom{n}{2} \sum_{i=1}^n f_e(x_i). \quad (2.11)$$

Thus,

$$\sum_{1 \leq i < j \leq n} (f_e(x_i - x_j) + f_e(x_i + x_j)) = 2(n-1) \sum_{i=1}^n f_e(x_i). \quad (2.12)$$

For each $j, k = 1, 2, \dots, n$ with $j \neq k$, we have

$$f_e\left(\sum_{i=1}^n x_i - 2x_j\right) + f_e\left(\sum_{i=1}^n x_i - 2x_k\right) = 2f_e\left(\sum_{i=1}^n x_i - x_j - x_k\right) + 2f_e(x_j - x_k). \quad (2.13)$$

Write down the above equation for every possible pair (j, k) and note that there are $\binom{n}{2}$ such pairs; so each $f_e(\sum_{i=1}^n x_i - 2x_j)$ appears $n-1$ times in all $\binom{n}{2}$ equations. Adding up the equations, we get

$$(n-1) \sum_{j=1}^n f_e\left(\sum_{i=1}^n x_i - 2x_j\right) = 2 \sum_{1 \leq j < k \leq n} f_e\left(\sum_{i=1}^n x_i - x_j - x_k\right) + 2 \sum_{1 \leq j < k \leq n} f_e(x_j - x_k). \quad (2.14)$$

For each $j = 1, 2, \dots, n$, we have

$$f_e\left(\sum_{i=1}^n x_i\right) + f\left(\sum_{i=1}^n x_i - 2x_j\right) = 2f_e\left(\sum_{i=1}^n x_i - x_j\right) + 2f_e(x_j). \quad (2.15)$$

Sum the above equation for all j 's and substitute the result from (2.12) and (2.14), then rearrange the resulting equation

$$\begin{aligned} n f_e\left(\sum_{i=1}^n x_i\right) + \frac{2}{n-1} \sum_{1 \leq j < k \leq n} f_e\left(\sum_{i=1}^n x_i - x_j - x_k\right) \\ = 2 \sum_{j=1}^n f_e\left(\sum_{i=1}^n x_i - x_j\right) + \frac{2}{n-1} \sum_{1 \leq i < j \leq n} f_e(x_i + x_j) - 2 \sum_{i=1}^n f_e(x_i). \end{aligned} \quad (2.16)$$

Note that $\sum_{j=1}^n f_e(\sum_{i=1}^n x_i - x_j)$ is the sum of f of x_i 's taken $n-1$ variables at a time, and $\sum_{1 \leq j < k \leq n} f_e(\sum_{i=1}^n x_i - x_j - x_k)$ is the sum of f of x_i 's taken $n-2$ variables at a time. From the induction assumption, (2.1) holds for $n-1$ and $n-2$ variables, that is,

$$\begin{aligned} \sum_{j=1}^n f_e\left(\sum_{i=1}^n x_i - x_j\right) + (n-1)(n-3) \sum_{i=1}^n f_e(x_i) &= (n-2) \sum_{1 \leq i < j \leq n} f_e(x_i + x_j), \\ \sum_{1 \leq j < k \leq n} f_e\left(\sum_{i=1}^n x_i - x_j - x_k\right) + \frac{(n-1)(n-2)(n-4)}{2} \sum_{i=1}^n f_e(x_i) \\ &= \frac{(n-2)(n-3)}{2} \sum_{1 \leq i < j \leq n} f_e(x_i + x_j). \end{aligned} \quad (2.17)$$

Substitute (2.17) into (2.16) and simplify, we will finally establish (2.1) on the even part of f . Thus, f satisfies (2.1) and the proof is complete. $\square$

3. The Ulam-Gavruta-Rassias stability

Rassias [10] established the Ulam stability of (2.1) in the special case when $n = 3$ on restricted domains. The following theorem provides a general condition for which a *true* general solution discussed in Theorem 2.1 exists near an approximate solution. For convenience, we define

$$Df(x_1, x_2, \dots, x_n) = f\left(\sum_{i=1}^n x_i\right) + (n-2) \sum_{i=1}^n f(x_i) - \sum_{1 \leq i < j \leq n} f(x_i + x_j). \quad (3.1)$$

From now on, we will refer to the even part and the odd part of a function by subscripts e and o , respectively.

THEOREM 3.1. *Let $n > 2$ be a positive integer, let X be a real vector space, let Y be a Banach space, let $\phi : X^n \rightarrow [0, \infty)$ be an even function. Define $\varphi(x) = \phi(x, x, -x, 0, \dots, 0)$ for all $x \in X$. If*

$$\sum_{i=0}^{\infty} 2^{-i} \varphi(2^i x) \text{ converges,} \quad \lim_{m \rightarrow \infty} 2^{-m} \phi(2^m x_1, \dots, 2^m x_n) = 0 \quad (3.2)$$

or

$$\sum_{i=1}^{\infty} 4^i \varphi(2^{-i} x) \text{ converges,} \quad \lim_{m \rightarrow \infty} 4^m \phi(2^{-m} x_1, \dots, 2^{-m} x_n) = 0 \quad (3.3)$$

for all $x_1, x_2, \dots, x_n \in X$, and a function $f : X \rightarrow Y$ satisfies $f(0) = 0$ and

$$\|Df(x_1, x_2, \dots, x_n)\| \leq \phi(x_1, x_2, \dots, x_n) \quad (3.4)$$

for all $x_1, x_2, \dots, x_n \in X$, then there exists a unique function $T : X \rightarrow Y$ that satisfies functional equation (2.1) and, if condition (3.2) holds,

$$\|f_e(x) - T_e(x)\| \leq \frac{1}{4} \sum_{i=0}^{\infty} 4^{-i} \varphi(2^i x), \quad \|f_o(x) - T_o(x)\| \leq \frac{1}{2} \sum_{i=0}^{\infty} 2^{-i} \varphi(2^i x_0) \quad (3.5)$$

or, if condition (3.3) holds,

$$\|f_e(x) - T_e(x)\| \leq \frac{1}{4} \sum_{i=1}^{\infty} 4^i \varphi(2^{-i} x), \quad \|f_o(x) - T_o(x)\| \leq \frac{1}{2} \sum_{i=1}^{\infty} 2^i \varphi(2^{-i} x). \quad (3.6)$$

The function T is given by

$$T(x) = \begin{cases} \lim_{m \rightarrow \infty} 4^{-m} f_e(2^m x) + 2^{-m} f_o(2^m x) & \text{if condition (3.2) holds,} \\ \lim_{m \rightarrow \infty} 4^m f_e(2^{-m} x) + 2^m f_o(2^{-m} x) & \text{if condition (3.3) holds} \end{cases} \quad (3.7)$$

for all $x \in X$.

Proof. We will prove the theorem for a function ϕ satisfying condition (3.2) and accordingly inequality (3.5). A proof for conditions (3.3) and (3.6) can be reproduced in a similar manner. Setting $(x_1, x_2, \dots, x_n) = (x, x, -x, 0, 0, \dots, 0)$ in (3.4) and simplifying, we have $\|3f(x) + f(-x) - f(2x)\| \leq \varphi(x)$. Replacing x by $-x$, we have $\|3f(-x) + f(x) - f(-2x)\| \leq \varphi(-x) = \varphi(x)$. Then,

$$\begin{aligned} & \|4f_e(x) - f_e(2x)\| \\ &= \frac{1}{2} \|(3f(x) + f(-x) - f(2x)) + (3f(-x) + f(x) - f(-2x))\| \\ &\leq \frac{1}{2} \|3f(x) + f(-x) - f(2x)\| + \frac{1}{2} \|3f(-x) + f(x) - f(-2x)\| \\ &\leq \frac{1}{2} \varphi(x) + \frac{1}{2} \varphi(x) = \varphi(x), \end{aligned} \quad (3.8)$$

$$\begin{aligned} & \|2f_o(x) - f_o(2x)\| \\ &= \frac{1}{2} \|(3f(x) + f(-x) - f(2x)) - (3f(-x) + f(x) - f(-2x))\| \\ &\leq \frac{1}{2} \|3f(x) + f(-x) - f(2x)\| + \frac{1}{2} \|3f(-x) + f(x) - f(-2x)\| \\ &\leq \frac{1}{2} \varphi(x) + \frac{1}{2} \varphi(x) = \varphi(x). \end{aligned}$$

Rewrite the inequality on f_e as $\|f_e(x) - 4^{-1}f_e(2x)\| \leq 4^{-1}\varphi(x)$ for all $x \in X$. Suppose that $\|f_e(x) - 4^{-m}f_e(2^m x)\| \leq (1/4) \sum_{i=0}^{m-1} 4^{-i}\varphi(2^i x)$ for a positive integer m . Then,

$$\begin{aligned} & \|f_e(x) - 4^{-(m+1)}f_e(2^{m+1}x)\| \\ &\leq \|f_e(x) - 4^{-m}f_e(2^m x)\| + \|4^{-m}f_e(2^m x) - 4^{-(m+1)}f_e(2^{m+1}x)\| \\ &\leq \|f_e(x) - 4^{-m}f_e(2^m x)\| + 4^{-m} \|f_e(2^m x) - 4^{-1}f_e(2 \cdot 2^m x)\| \\ &\leq \frac{1}{4} \sum_{i=0}^{m-1} 4^{-i}\varphi(2^i x) + 4^{-m}\varphi(2^m x) = \frac{1}{4} \sum_{i=0}^m 4^{-i}\varphi(2^i x). \end{aligned} \quad (3.9)$$

Hence, $\|f_e(x) - 4^{-m}f_e(2^m x)\| \leq (1/4) \sum_{i=0}^{m-1} 4^{-i}\varphi(2^i x)$ for every positive integer m .

If we rewrite the inequality for f_o as $\|f_o(x) - 2^{-1}f_o(2x)\| \leq 2^{-1}\varphi(x)$ and repeat the same steps as in the case of f_e , we will have $\|f_o(x) - 2^{-m}f_o(2^m x)\| \leq (1/2) \sum_{i=0}^{m-1} 2^{-i}\varphi(2^i x)$ for every positive integer m .

The convergence of the sequence $\{4^{-m}f_e(2^m x)\}$ can be settled as follows. For every positive integer p ,

$$\begin{aligned}
 \|4^{-(m+p)}f_e(2^{m+p}x) - 4^{-m}f_e(2^m x)\| &= 4^{-m}\|4^{-p}f_e(2^p \cdot 2^m x) - f_e(2^m x)\| \\
 &\leq 4^{-m} \cdot \frac{1}{4} \sum_{i=0}^{p-1} 4^{-i}\phi(2^i \cdot 2^m x) \\
 &\leq \frac{1}{4} \sum_{i=0}^{\infty} 4^{-(i+m)}\phi(2^{i+m}x).
 \end{aligned} \tag{3.10}$$

By the definition of ϕ and condition (3.2), the right-hand side approaches 0 as m goes to infinity, hence, we have a Cauchy sequence in a Banach space. Let $T_e(x) = \lim_{m \rightarrow \infty} 4^{-m}f_e(2^m x)$ for all $x \in X$, and thus $\|f_e(x) - T_e(x)\| \leq (1/4) \sum_{i=0}^{\infty} 4^{-i}\phi(2^i x)$. We can similarly show that $\{2^{-m}f_o(2^m x)\}$ converges; so let $T_o(x) = \lim_{m \rightarrow \infty} 2^{-m}f_o(2^m x)$ for all $x \in X$, and thus $\|f_o(x) - T_o(x)\| \leq (1/2) \sum_{i=0}^{\infty} 2^{-i}\phi(2^i x)$. Define $T(x) = T_e(x) + T_o(x)$ for all $x \in X$.

In order to show that T satisfies (2.1), we will in turn show that T_e and T_o satisfy (2.1). For convenience, define Df_e and Df_o as the even part and the odd part of Df in (3.1), respectively. For T_e , consider

$$\begin{aligned}
 4^{-m}\|Df_e(2^m x_1, \dots, 2^m x_n)\| \\
 &= 4^{-m} \cdot \frac{1}{2} \|Df(2^m x_1, \dots, 2^m x_n) + Df(-2^m x_1, \dots, -2^m x_n)\| \\
 &\leq 4^{-m}\phi(2^m x_1, \dots, 2^m x_n).
 \end{aligned} \tag{3.11}$$

As m tend to infinity, the left-hand side approaches $\|DT_e(x_1, \dots, x_n)\|$ and, by condition (3.2), the right-hand side approaches 0. Thus,

$$DT_e(x_1, x_2, \dots, x_n) = T_e\left(\sum_{i=1}^n x_i\right) + (n-2) \sum_{i=1}^n T_e(x_i) - \sum_{1 \leq i < j \leq n} T_e(x_i + x_j) = 0, \tag{3.12}$$

which shows that T_e satisfies (2.1).

We can similarly show that T_o satisfies (2.1) by considering

$$\begin{aligned}
 2^{-m}\|Df_o(2^m x_1, \dots, 2^m x_n)\| \\
 &= 2^{-m} \cdot \frac{1}{2} \|Df(2^m x_1, \dots, 2^m x_n) - Df(-2^m x_1, \dots, -2^m x_n)\| \\
 &\leq 2^{-m}\phi(2^m x_1, \dots, 2^m x_n),
 \end{aligned} \tag{3.13}$$

and take the limit as $m \rightarrow \infty$. Hence, $T = T_e + T_o$ satisfies (2.1) as desired.

To prove the uniqueness of T , suppose that there exists another function $S : X \rightarrow Y$ such that S satisfies (2.1) and satisfies the inequality (3.5) with T replaced by S . Then,

$$\begin{aligned} \|S(x) - T(x)\| &\leq \|S(x) - f(x)\| + \|T(x) - f(x)\| \\ &\leq \|S_e(x) - f_e(x)\| + \|S_o(x) - f_o(x)\| \\ &\quad + \|T_e(x) - f_e(x)\| + \|T_o(x) - f_o(x)\|. \end{aligned} \quad (3.14)$$

It is straightforward to show that every solution of the *quadratic* functional equation $f(x+y) + f(x-y) = 2f(x) + 2f(y)$ has the *quadratic* property $f(nx) = n^2 f(x)$ and every solution of the *linear* functional equation $f(x+y) = f(x) + f(y)$ has the *linear* property $f(nx) = nf(x)$ for every positive integer n and for every x in the domain. We thus obtain

$$\begin{aligned} \|S(x) - T(x)\| &\leq 4^{-m} \|S_e(2^m x) - f_e(2^m x)\| + 2^{-m} \|S_o(2^m x) - f_o(2^m x)\| \\ &\quad + 4^{-m} \|T_e(2^m x) - f_e(2^m x)\| + 2^{-m} \|T_o(2^m x) - f_o(2^m x)\| \\ &\leq 2 \left(4^{-m} \cdot \frac{1}{4} \sum_{i=0}^{\infty} 4^{-i} \varphi(2^i \cdot 2^m x) + \frac{1}{2^m} \cdot \frac{1}{2} \sum_{i=0}^{\infty} 2^{-i} \varphi(2^i \cdot 2^m x) \right) \\ &= \frac{1}{2} \sum_{i=0}^{\infty} 4^{-(i+m)} \varphi(2^{i+m} x) + \sum_{i=0}^{\infty} 2^{-(i+m)} \varphi(2^{i+m} x) \end{aligned} \quad (3.15)$$

for all $x \in X$. As m goes to infinity, the right-hand side approaches 0, and $S(x) = T(x)$ for all $x \in X$. This completes the proof. $\square$

The following corollary proves the Hyers-Ulam stability of (2.1).

COROLLARY 3.2. *If a function $f : X \rightarrow Y$ satisfies $f(0) = 0$ and the functional equation*

$$\|Df(x_1, x_2, \dots, x_n)\| \leq \varepsilon \quad (3.16)$$

for some $\varepsilon > 0$ and for all $x_1, x_2, \dots, x_n \in X$, then there exists a unique function $T : X \rightarrow Y$ that satisfies functional equation (2.1) and, for all $x \in X$,

$$\|f_e(x) - T_e(x)\| \leq \frac{\varepsilon}{3}, \quad \|f_o(x) - T_o(x)\| \leq \varepsilon. \quad (3.17)$$

Proof. Let $\phi(x_1, x_2, \dots, x_n) = \varepsilon$, then condition (3.2) in Theorem 3.1 holds. Hence, it follows from the theorem that there exists a unique function $T : X \rightarrow Y$ such that

$$\|f_e(x) - T_e(x)\| \leq \frac{1}{4} \sum_{i=0}^{\infty} 4^{-i} \cdot \varepsilon = \frac{\varepsilon}{3}, \quad \|f_o(x) - T_o(x)\| \leq \frac{1}{2} \sum_{i=0}^{\infty} 2^{-i} \varepsilon = \varepsilon. \quad (3.18)$$

$\square$

The following corollary proves the Hyers-Ulam-Rassias stability of (2.1).

COROLLARY 3.3. *Let p be a positive real number with $0 < p < 1$ or $p > 2$. If a function $f : X \rightarrow Y$ satisfies the inequality*

$$\|Df(x_1, x_2, \dots, x_n)\| \leq \varepsilon \sum_{i=1}^n \|x_i\|^p \quad (3.19)$$

for some $\varepsilon > 0$ and for all $x_1, x_2, \dots, x_n \in X$, then there exists a unique function $T : X \rightarrow Y$ that satisfies functional equation (2.1) and, for all $x \in X$,

$$\|f_e(x) - T_e(x)\| \leq \frac{3\varepsilon}{4|1 - 2^{p-2}|} \|x\|^p, \quad \|f_o(x) - T_o(x)\| \leq \frac{3\varepsilon}{2|1 - 2^{p-1}|} \|x\|^p. \quad (3.20)$$

Proof. Substituting $x_1 = x_2 = \dots = x_n = 0$ into (3.19), we get

$$f(0) + (n-2) \cdot n f(0) = \binom{n}{2} f(0). \quad (3.21)$$

Since $n > 2$, it follows that $1 + n(n-2) > \binom{n}{2}$, hence, $f(0) = 0$.

Let $\phi(x_1, x_2, \dots, x_n) = \varepsilon \sum_{i=1}^n \|x_i\|^p$. If $0 < p < 1$, then condition (3.2) in Theorem 3.1 holds and it follows that

$$\begin{aligned} \|f_e(x) - T_e(x)\| &\leq \frac{1}{4} \sum_{i=0}^{\infty} 4^{-i} (3\varepsilon \cdot 2^{ip} \|x\|^p) = \frac{3\varepsilon}{4(1 - 2^{p-2})} \|x\|^p, \\ \|f_o(x) - T_o(x)\| &\leq \frac{1}{2} \sum_{i=0}^{\infty} 2^{-i} (3\varepsilon \cdot 2^{ip} \|x\|^p) = \frac{3\varepsilon}{2(1 - 2^{p-1})} \|x\|^p. \end{aligned} \quad (3.22)$$

If $p > 1$, we apply Theorem 3.1 with condition (3.3) to get a similar result. $\square$

The following corollary proves the Ulam-Gavruta-Rassias stability of (2.1).

COROLLARY 3.4. *Let $p_1, p_2, \dots, p_n$ be nonnegative real numbers and $r = \sum_{i=1}^n p_i$ with $0 < r < 1$ or $r > 2$. If a function $f : X \rightarrow Y$ satisfies the inequality*

$$\|Df(x_1, x_2, \dots, x_n)\| \leq \varepsilon \prod_{i=1}^n \|x_i\|^{p_i} \quad (3.23)$$

for some $\varepsilon > 0$ and for all $x_1, x_2, \dots, x_n \in X$, then there exists a unique function $T : X \rightarrow Y$ that satisfies functional equation (2.1) and, for $n = 3$,

$$\|f_e(x) - T_e(x)\| \leq \frac{\varepsilon}{4|1 - 2^{r-2}|} \|x\|^r, \quad \|f_o(x) - T_o(x)\| \leq \frac{\varepsilon}{2|1 - 2^{r-1}|} \|x\|^r \quad (3.24)$$

for all $x \in X$.

Proof. We can show that $f(0) = 0$ by the same substitution used in the proof of Corollary 3.3. Let $\phi(x_1, x_2, \dots, x_n) = \varepsilon \prod_{i=1}^n \|x_i\|^{p_i}$. According to Theorem 3.1, if $0 < r < 1$, then condition (3.2) holds, and if $r > 2$, then condition (3.3) holds. If $n > 3$, then the desired result

immediately follows. However, for $n = 3$, we have

$$\begin{aligned} \|f_e(x) - T_e(x)\| &\leq \frac{1}{4} \sum_{i=0}^{\infty} 4^{-i} (\varepsilon \cdot 2^{ir} \|x\|^r) = \frac{\varepsilon}{4(1 - 2^{r-2})} \|x\|^r, \\ \|f_o(x) - T_o(x)\| &\leq \frac{1}{2} \sum_{i=0}^{\infty} 2^{-i} (\varepsilon \cdot 2^{ir} \|x\|^r) = \frac{\varepsilon}{2(1 - 2^{r-1})} \|x\|^r \end{aligned} \quad (3.25)$$

when $0 < r < 1$, and a similar result when $r > 1$. □

Acknowledgment

The author would like to thank the referee for providing valuable comments and helping in improving the content of this paper.

References

- [1] S. M. Ulam, *Problems in Modern Mathematics*, chapter 6, John Wiley & Sons, New York, NY, USA, 1964.
- [2] D. H. Hyers, "On the stability of the linear functional equation," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 27, no. 4, pp. 222–224, 1941.
- [3] J. M. Rassias, "On approximation of approximately linear mappings by linear mappings," *Journal of Functional Analysis*, vol. 46, no. 1, pp. 126–130, 1982.
- [4] J. M. Rassias, "On approximation of approximately linear mappings by linear mappings," *Bulletin des Sciences Mathématiques*, vol. 108, no. 4, pp. 445–446, 1984.
- [5] J. M. Rassias, "Solution of a problem of Ulam," *Journal of Approximation Theory*, vol. 57, no. 3, pp. 268–273, 1989.
- [6] J. M. Rassias, "Solution of a stability problem of Ulam," *Discussiones Mathematicae*, vol. 12, pp. 95–103, 1992.
- [7] J. M. Rassias, "On the stability of the Euler-Lagrange functional equation," *Chinese Journal of Mathematics*, vol. 20, no. 2, pp. 185–190, 1992.
- [8] J. M. Rassias, "Complete solution of the multi-dimensional problem of Ulam," *Discussiones Mathematicae*, vol. 14, pp. 101–107, 1994.
- [9] J. M. Rassias, "Solution of the Ulam stability problem for Euler-Lagrange quadratic mappings," *Journal of Mathematical Analysis and Applications*, vol. 220, no. 2, pp. 613–639, 1998.
- [10] J. M. Rassias, "On the Ulam stability of mixed type mappings on restricted domains," *Journal of Mathematical Analysis and Applications*, vol. 276, no. 2, pp. 747–762, 2002.

Paisan Nakmahachalasint: Department of Mathematics, Faculty of Science,
Chulalongkorn University, Bangkok 10330, Thailand
Email address: paisan.n@chula.ac.th