Research Article

Subordination Properties for Certain Analytic Functions

A. A. Attiya,^{1,2} Nak Eun Cho,³ and M. A. Kutbi⁴

¹Department of Mathematics, Faculty of Science, University of Mansoura, Mansoura 35516, Egypt

²Department of Mathematics, Teachers' College in Abha, King Khalid University, Abha, P.O. Box 249, Saudi Arabia

³Department of Applied Mathematics, Pukyong National University, Pusan 608-737, South Korea

⁴Department of Mathematics, Faculty of Science, King AbdulAziz University, P.O. Box 80203, Jeddah 21589, Saudi Arabia

Correspondence should be addressed to A. A. Attiya, aattiy@mans.edu.eg

Received 13 September 2007; Accepted 7 November 2007

Recommended by Brigitte Forster-Heinlein

The purpose of the present paper is to derive a subordination result for functions in the class $H_n^*(\alpha, \lambda, b)$ of normalized analytic functions in the open unit disk $\mathbb{U}$. A number of interesting applications of the subordination result are also considered.

Copyright © 2008 A. A. Attiya et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

Let A denote the class of functions of the form

$$f(z) = z + \sum_{n=2}^{\infty} a_n z^n \quad (1.1)$$

which are analytic in the unit disc $\mathbb{U} = \{z : |z| < 1\}$. We also denote by K the class of functions $f \in A$ that are convex in $\mathbb{U}$.

Given two functions $f, g \in A$, where f is given by (1.1) and g is defined by

$$g(z) = z + \sum_{n=2}^{\infty} b_n z^n, \quad (1.2)$$

the Hadamard product (or convolution) $f * g$ is defined by

$$(f * g)(z) = z + \sum_{n=2}^{\infty} a_n b_n z^n \quad (z \in \mathbb{U}). \quad (1.3)$$

By using the Hadamard product, Ruscheweyh [1] defined

$$D^\alpha f(z) = \frac{z}{(1-z)^{\alpha+1}} * f(z) \quad (\alpha \geq -1). \quad (1.4)$$

From the definition of (1.4), we observe that

$$D^n f(z) = \frac{z(z^{n-1}f(z))^{(n)}}{n!}, \quad (1.5)$$

when $n = \alpha \in \mathbb{N}_0 = \mathbb{N} \cup \{0\} = \{0, 1, 2, \dots\}$. The symbol $D^n f(z)$ ($n \in \mathbb{N}_0$) was called the n th-order Ruscheweyh derivative of f by Al-Amiri [2]. We also note that $D^0 f(z) = f(z)$ and $D^1 f(z) = zf'(z)$.

Definition 1.1. Suppose that $f \in A$. Then the function f is said to be a member of the class $H_n(\alpha, \lambda, b)$ if it satisfies

$$\left| \frac{\lambda(n+1)(D^{n+1}f(z)/z) + [1 - \lambda(n+1)](D^n f(z)/z) - 1}{\lambda(n+1)(D^{n+1}f(z)/z) + [1 - \lambda(n+1)](D^n f(z)/z) + 2b(1 - \alpha) - 1} \right| < 1 \quad (1.6)$$

$$(z \in \mathbb{U}; 0 \leq \alpha < 1; \lambda \geq 0; b \in \mathbb{C} \setminus \{0\}; n \in \mathbb{N}_0).$$

By specializing α, λ, b , and n , one can obtain various subclasses studied by many authors (see, e.g., [3–11]).

Definition 1.2. Let g be analytic and univalent in $\mathbb{U}$. If f is analytic in $\mathbb{U}$, $f(0) = g(0)$, and $f(\mathbb{U}) \subset g(\mathbb{U})$, then one says that f is subordinate to g in $\mathbb{U}$, and we write $f < g$ or $f(z) < g(z)$. One also says that g is superordinate to f in $\mathbb{U}$.

Definition 1.3. An infinite sequence $\{b_k\}_{k=1}^\infty$ of complex numbers will be called a subordinating factor sequence if for every univalent function f in K , one has

$$\sum_{k=1}^{\infty} b_k a_k z^k < f(z) \quad (z \in \mathbb{U}; a_1 = 1). \quad (1.7)$$

Lemma 1.4 (see [12]). *The sequence $\{b_k\}_{k=1}^\infty$ is a subordinating factor sequence if and only if*

$$\operatorname{Re} \left\{ 1 + 2 \sum_{k=1}^{\infty} b_k z^k \right\} > 0 \quad (z \in \mathbb{U}). \quad (1.8)$$

Now, we prove the following lemma which gives a sufficient condition for functions belonging to the class $H_n(\alpha, \lambda, b)$.

Lemma 1.5. *If the function f which is defined by (1.1) satisfies the following condition:*

$$\sum_{k=2}^{\infty} [1 + \lambda(k-1)] C(n, k) |a_k| \leq (1 - \alpha)|b| \quad (0 \leq \alpha < 1; \lambda \geq 0; b \in \mathbb{C} \setminus \{0\}; n \in \mathbb{N}_0), \quad (1.9)$$

where

$$C(n, k) = \prod_{j=2}^k \frac{(j+n-1)}{(k-1)!} \quad (k = 2, 3, \dots), \quad (1.10)$$

then $f \in H_n(\alpha, \lambda, b)$.

Proof. Suppose that the inequality (1.9) holds. Using the identity

$$z(D^n f(z))' = (n+1)D^{n+1}f(z) - nD^n f(z), \quad (1.11)$$

we have for $z \in \mathbb{U}$,

$$\begin{aligned} & \left| (1 - \lambda) \frac{D^n f(z)}{z} + \lambda(D^n f(z))' - 1 \right| - \left| 2b(1 - \alpha) + (1 - \lambda) \frac{D^n f(z)}{z} + \lambda(D^n f(z))' - 1 \right| \\ &= \left| \sum_{k=2}^{\infty} [1 + \lambda(k-1)] C(n, k) a_k z^{k-1} \right| - \left| 2b(1 - \alpha) + \sum_{k=2}^{\infty} [1 + \lambda(k-1)] C(n, k) a_k z^{k-1} \right| \\ &\leq \sum_{k=2}^{\infty} [1 + \lambda(k-1)] C(n, k) |a_k| |z|^{k-1} \\ &\quad - \left\{ 2|b|(1 - \alpha) - \sum_{k=2}^{\infty} [1 + \lambda(k-1)] C(n, k) |a_k| |z|^{k-1} \right\} \\ &\leq 2 \left\{ \sum_{k=2}^{\infty} [1 + \lambda(k-1)] C(n, k) |a_k| - |b|(1 - \alpha) \right\} \leq 0, \end{aligned} \quad (1.12)$$

which shows that f belongs to $H_n(\alpha, \lambda, b)$. $\square$

Let $H_n^*(\alpha, \lambda, b)$ denote the class of functions f in A whose Taylor-Maclaurin coefficients a_k satisfy the condition (1.9).

We note that

$$H_n^*(\alpha, \lambda, b) \subseteq H_n(\alpha, \lambda, b). \quad (1.13)$$

Example 1.6. (i) For $0 \leq \alpha < 1$, $\lambda > 0$, $b \in \mathbb{C} \setminus \{0\}$, and $n \in \mathbb{N}_0$, the following function defined by:

$$f_0(z) = z + \frac{2b(1 - \alpha)}{(n+1)(\lambda+1)} z^2 {}_3F_2 \left(1, 2, 1 + \frac{1}{\lambda}; 2 + \frac{1}{\lambda}, n+2; z \right) \quad (z \in \mathbb{U}), \quad (1.14)$$

is in the class $H_n(\alpha, \lambda, b)$.

(ii) For $0 \leq \alpha < 1$, $\lambda > 0$, $b \in \mathbb{C} \setminus \{0\}$, and $n \in \mathbb{N}_0$, the following functions defined by:

$$\begin{aligned} f_1(z) &= z \pm \frac{(1 - \alpha)|b|}{(\lambda+1)(n+1)} z^2 \quad (z \in \mathbb{U}), \\ f_2(z) &= z \pm \frac{(1 - \alpha)|b|}{(2\lambda+1)(n+1)(n+2)} z^3 \quad (z \in \mathbb{U}), \\ f_3(z) &= z \pm \frac{1}{(\lambda+1)(n+1)} z^2 \pm \frac{2[(1 - \alpha)|b| - 1]}{(2\lambda+1)(n+1)(n+2)} z^3 \quad (z \in \mathbb{U}) \end{aligned} \quad (1.15)$$

are in the class $H_n^*(\alpha, \lambda, b)$.

In this paper, we obtain a sharp subordination result associated with the class $H_n^*(\alpha, \lambda, b)$ by using the same techniques as in [13] (see also [14–16]). Some applications of the main result which give important results of analytic functions are also investigated.

2. Main theorem

Theorem 2.1. *Let $f \in H_n^*(\alpha, \lambda, b)$. Then*

$$\frac{(\lambda+1)(n+1)}{2[(\lambda+1)(n+1) + |b|(1-\alpha)]} (f * g)(z) < g(z) \quad (z \in \mathbb{U}) \quad (2.1)$$

for every function g in K , and

$$\operatorname{Re} f(z) > -\frac{(\lambda+1)(n+1) + |b|(1-\alpha)}{(\lambda+1)(n+1)}. \quad (2.2)$$

The constant $(\lambda+1)(n+1)/2[(\lambda+1)(n+1) + |b|(1-\alpha)]$ cannot be replaced by a larger one.

Proof. Let $f \in H_n^*(\alpha, \lambda, b)$ and let

$$g(z) = z + \sum_{k=2}^{\infty} c_k z^k \quad (2.3)$$

be any function in the class K . Then we readily have

$$\frac{(\lambda+1)(n+1)}{2[(\lambda+1)(n+1) + |b|(1-\alpha)]} (f * g)(z) = \frac{(\lambda+1)(n+1)}{2[(\lambda+1)(n+1) + |b|(1-\alpha)]} \left(z + \sum_{k=2}^{\infty} a_k c_k z^k \right). \quad (2.4)$$

Thus, by Definition 1.2, the subordination result (2.1) will hold true if the sequence

$$\left\{ \frac{(\lambda+1)(n+1)a_k}{2[(\lambda+1)(n+1) + |b|(1-\alpha)]} \right\}_{k=1}^{\infty} \quad (2.5)$$

is a subordinating factor sequence, with $a_1 = 1$. In view of Lemma 1.4, this is equivalent to the following inequality:

$$\operatorname{Re} \left\{ 1 + \sum_{k=1}^{\infty} \frac{(\lambda+1)(n+1)}{[(\lambda+1)(n+1) + |b|(1-\alpha)]} a_k z^k \right\} > 0 \quad (z \in \mathbb{U}). \quad (2.6)$$

Now, since

$$[1 + \lambda(k-1)]C(n, k) \quad (\lambda \geq 0, n \in \mathbb{N}_0) \quad (2.7)$$

is an increasing function of k , we have

$$\begin{aligned}
& \operatorname{Re} \left\{ 1 + \sum_{k=1}^{\infty} \frac{(\lambda+1)(n+1)}{[(\lambda+1)(n+1) + |b|(1-\alpha)]} a_k z^k \right\} \\
&= \operatorname{Re} \left\{ 1 + \frac{(\lambda+1)(n+1)}{[(\lambda+1)(n+1) + |b|(1-\alpha)]} z \right. \\
&\quad \left. + \frac{1}{[(\lambda+1)(n+1) + |b|(1-\alpha)]} \sum_{k=2}^{\infty} (\lambda+1)(n+1) a_k z^k \right\} \\
&> 1 - \frac{(\lambda+1)(n+1)}{[(\lambda+1)(n+1) + |b|(1-\alpha)]} r \\
&\quad - \frac{1}{[(\lambda+1)(n+1) + |b|(1-\alpha)]} \sum_{k=2}^{\infty} [1 + \lambda(k-1)] C(n, k) |a_k| r^k \\
&> 1 - \frac{(\lambda+1)(n+1)}{[(\lambda+1)(n+1) + |b|(1-\alpha)]} r - \frac{|b|(1-\alpha)}{[(\lambda+1)(n+1) + |b|(1-\alpha)]} r > 0 \quad (|z| = r).
\end{aligned} \tag{2.8}$$

This proves the inequality (2.6), and hence also the subordination result (2.1) asserted by Theorem 2.1. The inequality (2.2) follows from (2.1) by taking

$$g(z) = \frac{z}{1-z} \in K. \tag{2.9}$$

Next, we consider the function

$$f_1(z) = z - \frac{|b|(1-\alpha)}{(\lambda+1)(n+1)} z^2 \quad (0 \leq \alpha < 1; \lambda \geq 0; b \in \mathbb{C} \setminus \{0\}; n \in \mathbb{N}_0) \tag{2.10}$$

which is a member of the class $H_n^*(\alpha, \lambda, b)$. Then by using (2.1), we have

$$\frac{(\lambda+1)(n+1)}{2[(\lambda+1)(n+1) + |b|(1-\alpha)]} f_1(z) < \frac{z}{1-z} \quad (z \in \mathbb{U}). \tag{2.11}$$

It can be easily verified for the function $f_1(z)$ defined by (2.10) that

$$\inf_{z \in \mathbb{U}} \left\{ \operatorname{Re} \left(\frac{(\lambda+1)(n+1)}{2[(\lambda+1)(n+1) + |b|(1-\alpha)]} f_1(z) \right) \right\} = -\frac{1}{2} \quad (z \in \mathbb{U}) \tag{2.12}$$

which completes the proof of Theorem 2.1. $\square$

3. Some applications

Taking $n = 0$ in Theorem 2.1, we obtain the following.

Corollary 3.1. *If the function f defined by (1.1) satisfies*

$$\sum_{k=2}^{\infty} [1 + \lambda(k-1)] |a_k| \leq m \quad (\lambda \geq 0, m > 0) \tag{3.1}$$

then for every function g in K , one has

$$\begin{aligned} \frac{(\lambda + 1)}{2(\lambda + m + 1)}(f * g)(z) &< g(z), \quad (z \in \mathbb{U}), \\ \operatorname{Re} f(z) &> -\left(1 + \frac{m}{\lambda + 1}\right). \end{aligned} \quad (3.2)$$

The constant $(\lambda + 1)/2(\lambda + m + 1)$ cannot be replaced by larger one.

Putting $\lambda = 0$ in Theorem 2.1, we have the following corollary.

Corollary 3.2. *If the function f defined by (1.1) satisfies*

$$\sum_{k=2}^{\infty} C(n, k) |a_k| \leq m, \quad m > 0, \quad n \in \mathbb{N}_0, \quad (3.3)$$

where $C(n, k)$ is defined by (1.10), then for every function g in K , one has

$$\begin{aligned} \frac{(n + 1)}{2(n + m + 1)}(f * g)(z) &< g(z) \quad (z \in \mathbb{U}), \\ \operatorname{Re} f(z) &> -\left(1 + \frac{m}{n + 1}\right). \end{aligned} \quad (3.4)$$

The constant $(n + 1)/2(n + m + 1)$ cannot be replaced by larger one.

Next, letting $\lambda = 1$ and $n = 0$, in Theorem 2.1, we obtain the following corollary.

Corollary 3.3. *If the function f satisfies*

$$\sum_{k=2}^{\infty} k |a_k| \leq m \quad (m > 0), \quad (3.5)$$

then for every function g in K , one has

$$\begin{aligned} \frac{1}{(m + 2)}(f * g)(z) &< g(z) \quad (z \in \mathbb{U}), \\ \operatorname{Re} f(z) &> -\left(1 + \frac{m}{2}\right). \end{aligned} \quad (3.6)$$

The constant $1/(m + 2)$ cannot be replaced by a larger one.

Remark 3.4. Putting $\lambda = 1$, $m = 1$, and $n = 0$, in Theorem 2.1, we obtain the result due to Singh [17].

Also, by taking $\lambda = 0$ and $n = 0$, in Theorem 2.1, we have the following.

Corollary 3.5. *If the function f satisfies*

$$\sum_{k=2}^{\infty} |a_k| \leq m \quad (m > 0), \quad (3.7)$$

then for every function g in K , one has

$$\frac{1}{2(m+1)}(f*g)(z) < g(z) \quad (z \in \mathbb{U}),$$

$$\operatorname{Re} f(z) > -(1+m).$$
(3.8)

The constant $1/2(m+1)$ cannot be replaced by a larger one.

It is clearly from the proof of Theorem 2.1 that the function $f(z) = z - mz^2$ ($m > 0$, $z \in \mathbb{U}$) is the extremal function of Corollary 3.5. Also, the following example gives a nonpolynomial extremal function for the same corollary.

Example 3.6. Let the function h be defined by

$$h(z) = \frac{(m+1)z}{(m+1) + mz} \quad (m > 0, z \in \mathbb{U}),$$
(3.9)

the above function is analytic in $\mathbb{U}$ and it is equivalent to

$$h(z) = z + \sum_{k=2}^{\infty} \left(\frac{-m}{m+1} \right)^{k-1} z^k.$$
(3.10)

Then we have

$$\sum_{k=2}^{\infty} \left| \left(\frac{-m}{m+1} \right)^{k-1} \right| = m.$$
(3.11)

Therefore, the Taylor-Maclaurin coefficients of the function h satisfy the condition in Corollary 3.5. Moreover, it can be easily verified that

$$\inf_{z \in \mathbb{U}} \operatorname{Re} h(z) = h(-1) = -(m+1).$$
(3.12)

Then, the constant $-(m+1)$ cannot be replaced by a larger one. Therefore, the function h is the extremal function of Corollary 3.5.

References

- [1] S. Ruscheweyh, "New criteria for univalent functions," *Proceedings of the American Mathematical Society*, vol. 49, no. 1, pp. 109–115, 1975.
- [2] H. S. Al-Amiri, "On Ruscheweyh derivatives," *Annales Polonici Mathematici*, vol. 38, no. 1, pp. 88–94, 1980.
- [3] M. P. Chen, "On functions satisfying $\operatorname{Re}\{f(z)/z\} > \alpha$," *Tamkang Journal of Mathematics*, vol. 5, pp. 231–234, 1974.
- [4] M. P. Chen, "On the regular functions satisfying $\operatorname{Re}\{f(z)/z\} > \alpha$," *Bulletin of the Institute of Mathematics. Academia Sinica*, vol. 3, no. 1, pp. 65–70, 1975.
- [5] K. K. Dixit and S. K. Pal, "On a class of univalent functions related to complex order," *Indian Journal of Pure and Applied Mathematics*, vol. 26, no. 9, pp. 889–896, 1995.
- [6] T. G. Ezrohi, "Certain estimates in special classes of univalent functions regular in the circle $|z| < 1$," *Dopovidi Akademiji Nauk Ukrajin's'koji RSR*, vol. 1965, pp. 984–988, 1965.

- [7] R. M. Goel, "The radius of convexity and starlikeness for certain classes of analytic functions with fixed second coefficients," *Annales Universitatis Mariae Curie-Skłodowska. Sectio A*, vol. 25, pp. 33–39, 1971.
- [8] Y. C. Kim and F. Rønning, "Integral transforms of certain subclasses of analytic functions," *Journal of Mathematical Analysis and Applications*, vol. 258, no. 2, pp. 466–489, 2001.
- [9] T. H. MacGregor, "Functions whose derivative has a positive real part," *Transactions of the American Mathematical Society*, vol. 104, no. 3, pp. 532–537, 1962.
- [10] A. Swaminathan, "Certain sufficiency conditions on Gaussian hypergeometric functions," *Journal of Inequalities in Pure and Applied Mathematics*, vol. 5, no. 4, article 83, pp. 1–10, 2004.
- [11] K. Yamaguchi, "On functions satisfying $\operatorname{Re}\{f(z)/z\} < 0$," *Proceedings of the American Mathematical Society*, vol. 17, no. 3, pp. 588–591, 1966.
- [12] H. S. Wilf, "Subordinating factor sequences for convex maps of the unit circle," *Proceedings of the American Mathematical Society*, vol. 12, no. 5, pp. 689–693, 1961.
- [13] H. M. Srivastava and A. A. Attiya, "Some subordination results associated with certain subclasses of analytic functions," *Journal of Inequalities in Pure and Applied Mathematics*, vol. 5, no. 4, pp. 1–6, 2004, article 82.
- [14] R. M. Ali, V. Ravichandran, and N. Seenivasagan, "Subordination by convex functions," *International Journal of Mathematics and Mathematical Sciences*, vol. 2006, Article ID 62548, 6 pages, 2006.
- [15] A. A. Attiya, "On some applications of a subordination theorem," *Journal of Mathematical Analysis and Applications*, vol. 311, no. 2, pp. 489–494, 2005.
- [16] B. A. Frasin, "Subordination results for a class of analytic functions defined by a linear operator," *Journal of Inequalities in Pure and Applied Mathematics*, vol. 7, no. 4, article 134, pp. 1–7, 2006.
- [17] S. Singh, "A subordination theorem for spirallike functions," *International Journal of Mathematics and Mathematical Sciences*, vol. 24, no. 7, pp. 433–435, 2000.