Research Article

A Note on Four-Variable Reciprocity Theorem

Chandrashekar Adiga and P. S. Guruprasad

Department of Studies in Mathematics, University of Mysore, Manasagangotri, Mysore 570 006, India

Correspondence should be addressed to Chandrashekar Adiga, adiga.c@yahoo.com

Received 12 September 2009; Accepted 7 December 2009

Recommended by Teodor Bulboacă

We give new proof of a four-variable reciprocity theorem using Heine's transformation, Watson's transformation, and Ramanujan's ${}_1\psi_1$ -summation formula. We also obtain a generalization of Jacobi's triple product identity.

Copyright © 2009 C. Adiga and P. S. Guruprasad. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

Throughout the paper, we let $|q| < 1$ and we employ the standard notation:

$$\begin{aligned}(a)_0 &:= (a; q)_0 = 1, \\ (a)_\infty &:= (a; q)_\infty = \prod_{n=0}^{\infty} (1 - aq^n), \\ (a)_n &:= (a; q)_n = \frac{(a; q)_\infty}{(aq^n; q)_\infty}, \quad -\infty < n < \infty.\end{aligned}\tag{1.1}$$

Ramanujan [1] stated several q -series identities in his "lost" notebook. One of the beautiful identities is the two-variable reciprocity theorem.

Theorem 1.1 (see [2]). *For $ab \neq 0$,*

$$\rho(a, b) - \rho(b, a) = \left(\frac{1}{b} - \frac{1}{a} \right) \frac{(aq/b)_\infty (bq/a)_\infty (q)_\infty}{(-aq)_\infty (-bq)_\infty},\tag{1.2}$$

where

$$\rho(a, b) := \left(1 + \frac{1}{b}\right) \sum_{n=0}^{\infty} \frac{(-1)^n q^{n(n+1)/2} a^n b^{-n}}{(-aq)_n}. \quad (1.3)$$

In the recent past many new proofs of (1.2) have been found. The first proof of (1.2) was given by Andrews [3] using four-free-variable identity and Jacobi's triple product identity. Further, Andrews [4] applied (1.2) in proving Euler partition identity analogues stated in [1]. Somashekara and Fathima [5] established an equivalent version of (1.2) using Ramanujan's ${}_1\psi_1$ summation formula [6] and Heine's transformation [7, 8]. Berndt et al. [9] also derived (1.2) using the same above mentioned two transformations. In fact, Berndt et al. [9] in the same paper have given two more proofs of (1.2) one employing the Rogers-Fine identity [10] and the other is purely combinatorial. Using the q -binomial theorem:

$$\sum_{n=0}^{\infty} \frac{(a)_n}{(q)_n} t^n = \frac{(at)_{\infty}}{(t)_{\infty}}, \quad |t| < 1, \quad |q| < 1, \quad (1.4)$$

Kim et al. [11] gave a much different proof of (1.2). Guruprasad and Pradeep [12] also devised a proof of (1.2) using the q -binomial theorem. Adiga and Anitha [13] established (1.2) along the lines of Ismail's proof [14] of Ramanujan's ${}_1\psi_1$ summation formula. Further, they showed that the reciprocity theorem (1.2) leads to a q -integral extension of the classical gamma function. Kang [2] constructed a proof of (1.2) along the lines of Venkatachaliengar's proof of the Ramanujan ${}_1\psi_1$ summation formula [6, 15].

In [2] Kang proved the following three- and four-variable generalizations of (1.2).

For $|c| < |a| < 1$ and $|c| < |b| < 1$,

$$\rho_3(a, b, c) - \rho_3(b, a, c) = \left(\frac{1}{b} - \frac{1}{a}\right) \frac{(c)_{\infty} (aq/b)_{\infty} (bq/a)_{\infty} (q)_{\infty}}{(-c/a)_{\infty} (-c/b)_{\infty} (-aq)_{\infty} (-bq)_{\infty}}, \quad (1.5)$$

where

$$\rho_3(a, b, c) := \left(1 + \frac{1}{b}\right) \sum_{n=0}^{\infty} \frac{(c)_n (-1)^n q^{n(n+1)/2} a^n b^{-n}}{(-aq)_n (-c/b)_{n+1}}, \quad a, \frac{c}{b} \neq -q^{-n}, \quad (1.6)$$

and for $|c|, |d| < |a|, |b| < 1$,

$$\rho_4(a, b, c, d) - \rho_4(b, a, c, d) = \left(\frac{1}{b} - \frac{1}{a}\right) \frac{(d)_{\infty} (c)_{\infty} (cd/ab)_{\infty} (aq/b)_{\infty} (bq/a)_{\infty} (q)_{\infty}}{(-d/a)_{\infty} (-d/b)_{\infty} (-c/a)_{\infty} (-c/b)_{\infty} (-aq)_{\infty} (-bq)_{\infty}}, \quad (1.7)$$

where

$$\rho_4(a, b, c, d) := \left(1 + \frac{1}{b}\right) \sum_{n=0}^{\infty} \frac{(d)_n (c)_n (cd/ab)_n (1 + cdq^{2n}/b) (-1)^n q^{n(n+1)/2} a^n b^{-n}}{(-aq)_n (-c/b)_{n+1} (-d/b)_{n+1}}, \quad a, \frac{c}{b}, \frac{d}{b} \neq -q^{-n}. \quad (1.8)$$

Kang [2] established (1.5) on employing Ramanujan's ${}_1\psi_1$ summation formula and Jackson's transformation of ${}_2\phi_1$ and ${}_2\phi_2$ -series. Recently (1.5) was derived by Adiga and Guruprasad [16] using q -binomial theorem and Gauss summation formula. Somashekara and Mamta [17, 18] obtained (1.5) using the two-variable reciprocity theorem (1.2), Jackson's transformation, and again two-variable reciprocity theorem by parameter augmentation. Zhang [19] also established (1.5).

Kang [2] established (1.7) on employing Andrews's generalization of ${}_1\psi_1$ summation formula, Sears's transformation of ${}_3\phi_2$ -series, and a limiting case of Watson's transformation for a terminating very well-poised ${}_8\phi_7$ -series [8]:

$$\begin{aligned} & \sum_{n=0}^{\infty} \frac{(\alpha)_n (\beta)_n (\gamma)_n (\delta)_n (\epsilon)_n (1 - \alpha q^{2n}) q^{n(n+3)/2}}{(\alpha q/\beta)_n (\alpha q/\gamma)_n (\alpha q/\delta)_n (\alpha q/\epsilon)_n (q)_n (1 - \alpha)} \left(\frac{-\alpha^2}{\beta \gamma \delta \epsilon} \right)^n \\ &= \frac{(\alpha q)_\infty (\alpha q/\delta \epsilon)_\infty}{(\alpha q/\delta)_\infty (\alpha q/\epsilon)_\infty} \sum_{n=0}^{\infty} \frac{(\delta)_n (\epsilon)_n (\alpha q/\beta \gamma)_n}{(\alpha q/\beta)_n (\alpha q/\gamma)_n (q)_n} \left(\frac{\alpha q}{\delta \epsilon} \right)^n. \end{aligned} \quad (1.9)$$

Recently Ma [20, 21] proved a six-variable generalization and a five-variable generalization of (1.2). The main purpose of this paper is to provide a new proof of (1.7) using (1.9), Heine's transformation:

$$\sum_{n=0}^{\infty} \frac{(\alpha)_n (\beta)_n}{(q)_n (\gamma)_n} z^n = \frac{(\gamma/\beta)_\infty (\beta z)_\infty}{(\gamma)_\infty (z)_\infty} \sum_{n=0}^{\infty} \frac{(\alpha \beta z/\gamma)_n (\beta)_n}{(\beta z)_n (q)_n} \left(\frac{\gamma}{\beta} \right)^n, \quad |q| < 1, |z| < 1, |\gamma| < |\beta| < 1 \quad (1.10)$$

and Ramanujan's ${}_1\psi_1$ summation formula:

$${}_1\psi_1(a; b; z) := \sum_{n=-\infty}^{\infty} \frac{(a)_n}{(b)_n} z^n = \frac{(b/a)_\infty (az)_\infty (q/az)_\infty (q)_\infty}{(q/a)_\infty (b/az)_\infty (b)_\infty (z)_\infty}, \quad |q| < 1, \left| \frac{b}{a} \right| < |z| < 1. \quad (1.11)$$

Jacobi's triple product identity states that

$$\sum_{n=-\infty}^{\infty} q^{n(n+1)/2} z^n = (q)_\infty (-zq)_\infty \left(-\frac{1}{z} \right)_\infty, \quad z \neq 0, |q| < 1. \quad (1.12)$$

Andrews [22] gave a proof of (1.12) using Euler identities. Combinatorial proofs of Jacobi's triple product identity were given by Wright [23], Cheema [24], and Sudler [25]. We can also find a proof of (1.12) in [26]. Using (1.12), Hirschhorn [27, 28] established Jacobi's two-square and four-square theorems.

Somashekara and Fathima [5] and Kim et al. [11] established

$$\sum_{n=0}^{\infty} \frac{(-1)^n a^n b^{-n} q^{n(n+1)/2}}{(-a)_{n+1}} - \sum_{n=0}^{\infty} \frac{(-1)^n a^{-(n+1)} b^{n+1} q^{n(n+1)/2}}{(-b)_{n+1}} = \frac{(aq/b)_{\infty} (b/a)_{\infty} (q)_{\infty}}{(-a)_{\infty} (-b)_{\infty}}. \quad (1.13)$$

Note that (1.13) which is equivalent to (1.2) may be considered as a two-variable generalization of (1.12). Corteel and Lovejoy [29, equation (1.5)] have given a bijective proof of (1.13) using representations of over partitions. All the reciprocity theorems (1.2), (1.5), and (1.7) are generalizations of Jacobi's triple product identity (1.12).

We also obtain a generalization of Jacobi's triple product identity (1.12) which is due to Kang [2].

2. Proof of (1.7)—The Four-Variable Reciprocity Theorem

On employing q -binomial theorem, we have

$$\begin{aligned} \sum_{n=0}^{\infty} \frac{(-cq)_n (-dq)_n}{(-aq)_n (-bq)_n} q^n &= \frac{(-dq)_{\infty}}{(-bq)_{\infty}} \sum_{n=0}^{\infty} \frac{(-cq)_n (-bq^{n+1})_{\infty}}{(-aq)_n (-dq^{n+1})_{\infty}} q^n \\ &= \frac{(-dq)_{\infty}}{(-bq)_{\infty}} \sum_{m=0}^{\infty} \frac{(b/d)_m}{(q)_m} (-dq)^m \sum_{n=0}^{\infty} \frac{(-cq)_n}{(-aq)_n} (q^{m+1})^n. \end{aligned} \quad (2.1)$$

On using Heine's transformation (1.10) with $\alpha = -cq$, $\beta = q$, $\gamma = -aq$, $z = q^{m+1}$, we have

$$\begin{aligned} \sum_{n=0}^{\infty} \frac{(-cq)_n}{(-aq)_n} (q^{m+1})^n &= \frac{(q^{m+2})_{\infty} (-a)_{\infty}}{(q^{m+1})_{\infty} (-aq)_{\infty}} \sum_{n=0}^{\infty} \frac{(cq^{m+2}/a)_n}{(q^{m+2})_n} (-a)^n \\ &= \frac{(q)_m (1+a) (-a)^{-m-1}}{(cq/a)_{m+1}} \sum_{n=0}^{\infty} \frac{(cq/a)_{n+m+1}}{(q)_{n+m+1}} (-a)^{n+m+1} \\ &= \frac{(q)_m (1+a) (-a)^{-m-1}}{(cq/a)_{m+1}} \left[\sum_{n=0}^{\infty} \frac{(cq/a)_n}{(q)_n} (-a)^n - \sum_{n=0}^m \frac{(cq/a)_n}{(q)_n} (-a)^n \right] \\ &= \frac{(q)_m (-a)^{-m-1} (-cq)_{\infty}}{(cq/a)_{m+1} (-aq)_{\infty}} - \frac{(q)_m (1+a) (-a)^{-m-1}}{(cq/a)_{m+1}} \sum_{n=0}^m \frac{(cq/a)_n}{(q)_n} (-a)^n. \end{aligned} \quad (2.2)$$

Substituting this in (2.1), we obtain

$$\sum_{n=0}^{\infty} \frac{(-cq)_n (-dq)_n}{(-aq)_n (-bq)_n} q^n = \frac{(-dq)_{\infty} (-cq)_{\infty}}{(-a) (-bq)_{\infty} (-aq)_{\infty}} \sum_{m=0}^{\infty} \frac{(b/d)_m}{(cq/a)_{m+1}} \left(\frac{dq}{a}\right)^m$$

$$+ \frac{(1+a^{-1})(-dq)_{\infty}}{(-bq)_{\infty}} \sum_{m=0}^{\infty} \sum_{n=0}^m \frac{(b/d)_m (dq/a)^m (cq/a)_n (-a)^n}{(cq/a)_{m+1} (q)_n}. \quad (2.3)$$

Now,

$$\frac{(1+a^{-1})(-dq)_{\infty}}{(-bq)_{\infty}} \sum_{m=0}^{\infty} \sum_{n=0}^m \frac{(b/d)_m (dq/a)^m (cq/a)_n (-a)^n}{(cq/a)_{m+1} (q)_n}$$

$$= \frac{(1+a^{-1})(-dq)_{\infty}}{(-bq)_{\infty}} \sum_{n=0}^{\infty} \frac{(b/d)_n (-dq)^n}{(q)_n (1-cq^{n+1}/a)} \sum_{m=0}^{\infty} \frac{(bq^n/d)_m (dq/a)^m}{(cq^{n+2}/a)_m}$$

$$= \frac{(1+a^{-1})(-dq)_{\infty}}{(-bq)_{\infty}} \sum_{n=0}^{\infty} \frac{(b/d)_n (-dq)^n}{(q)_n (1-cq^{n+1}/a)} \cdot \frac{(1-cq^{n+1}/a)}{(1-dq/a)} \sum_{m=0}^{\infty} \frac{(b/c)_m (cq^{n+1}/a)^m}{(dq^2/a)_m},$$

$$\left(\text{on using (1.10) with } \alpha = \frac{bq^n}{d}, \beta = q, \gamma = \frac{cq^{n+2}}{a}, z = \frac{dq}{a} \right) \quad (2.4)$$

$$= \frac{(1+a^{-1})(-dq)_{\infty}}{(-bq)_{\infty}} \sum_{m=0}^{\infty} \frac{(b/c)_m (cq/a)^m}{(dq/a)_{m+1}} \sum_{n=0}^{\infty} \frac{(b/d)_n (-dq^{m+1})^n}{(q)_n}$$

$$= \frac{(1+a^{-1})(-dq)_{\infty}}{(-bq)_{\infty}} \sum_{m=0}^{\infty} \frac{(b/c)_m (cq/a)^m}{(dq/a)_{m+1}} \cdot \frac{(-bq^{m+1})_{\infty}}{(-dq^{m+1})_{\infty}}$$

$$= \left(1+a^{-1}\right) \sum_{m=0}^{\infty} \frac{(b/c)_m (-dq)_m}{(-bq)_m (dq/a)_{m+1}} \left(\frac{cq}{a}\right)^m.$$

Substituting (2.4) in (2.3), we obtain

$$\sum_{n=0}^{\infty} \frac{(-cq)_n (-dq)_n}{(-aq)_n (-bq)_n} q^n = \frac{(-dq)_{\infty} (-cq)_{\infty}}{(-a) (-bq)_{\infty} (-aq)_{\infty}} \sum_{m=0}^{\infty} \frac{(b/d)_m}{(cq/a)_{m+1}} \left(\frac{dq}{a}\right)^m$$

$$+ \left(1+a^{-1}\right) \sum_{m=0}^{\infty} \frac{(b/c)_m (-dq)_m}{(-bq)_m (dq/a)_{m+1}} \left(\frac{cq}{a}\right)^m \quad (2.5)$$

$$= \frac{(-dq)_{\infty} (-cq)_{\infty}}{(-a) (-bq)_{\infty} (-aq)_{\infty}} \sum_{m=0}^{\infty} \frac{(b/c)_m}{(dq/a)_{m+1}} \left(\frac{cq}{a}\right)^m$$

$$+ \left(1+a^{-1}\right) \sum_{m=0}^{\infty} \frac{(b/c)_m (-dq)_m}{(-bq)_m (dq/a)_{m+1}} \left(\frac{cq}{a}\right)^m.$$

(Here, we used (1.10) with $\alpha = b/d, \beta = q, \gamma = cq^2/a, z = dq/a$.)

Changing c to $-c/q$, d to $-d/q$ in (2.5), we get

$$\begin{aligned} \sum_{n=0}^{\infty} \frac{(c)_n (d)_n}{(-aq)_n (-bq)_n} q^n &= \frac{(d)_{\infty} (c)_{\infty}}{(-a)(-bq)_{\infty} (-aq)_{\infty}} \sum_{m=0}^{\infty} \frac{(-bq/c)_m}{(-d/a)_{m+1}} \left(-\frac{c}{a}\right)^m \\ &\quad + \left(1 + a^{-1}\right) \sum_{m=0}^{\infty} \frac{(-bq/c)_m (d)_m}{(-bq)_m (-d/a)_{m+1}} \left(-\frac{c}{a}\right)^m. \end{aligned} \quad (2.6)$$

Interchanging a and b in (2.6), we have

$$\begin{aligned} \sum_{n=0}^{\infty} \frac{(c)_n (d)_n}{(-aq)_n (-bq)_n} q^n &= \frac{(d)_{\infty} (c)_{\infty}}{(-b)(-bq)_{\infty} (-aq)_{\infty}} \sum_{m=0}^{\infty} \frac{(-aq/c)_m}{(-d/b)_{m+1}} \left(-\frac{c}{b}\right)^m \\ &\quad + \left(1 + b^{-1}\right) \sum_{m=0}^{\infty} \frac{(-aq/c)_m (d)_m}{(-aq)_m (-d/b)_{m+1}} \left(-\frac{c}{b}\right)^m. \end{aligned} \quad (2.7)$$

Subtracting (2.6) from (2.7), we deduce that

$$\begin{aligned} &\frac{(d)_{\infty} (c)_{\infty}}{(-bq)_{\infty} (-aq)_{\infty}} \left[\frac{1}{b} \sum_{m=0}^{\infty} \frac{(-aq/c)_m}{(-d/b)_{m+1}} \left(-\frac{c}{b}\right)^m - \frac{1}{a} \sum_{m=0}^{\infty} \frac{(-bq/c)_m}{(-d/a)_{m+1}} \left(-\frac{c}{a}\right)^m \right] \\ &= \left(1 + b^{-1}\right) \sum_{m=0}^{\infty} \frac{(-aq/c)_m (d)_m}{(-aq)_m (-d/b)_{m+1}} \left(-\frac{c}{b}\right)^m - \left(1 + a^{-1}\right) \sum_{m=0}^{\infty} \frac{(-bq/c)_m (d)_m}{(-bq)_m (-d/a)_{m+1}} \left(-\frac{c}{a}\right)^m. \end{aligned} \quad (2.8)$$

Now change a to $-b/d$, b to $-c/a$, and z to $-d/a$ in (1.11) to obtain

$$\sum_{n=1}^{\infty} \frac{(-b/d)_n}{(-c/a)_n} \left(-\frac{d}{a}\right)^n + \sum_{n=0}^{\infty} \frac{(-aq/c)_n}{(-dq/b)_n} \left(-\frac{c}{b}\right)^n = \frac{(cd/ab)_{\infty} (b/a)_{\infty} (aq/b)_{\infty} (q)_{\infty}}{(-c/a)_{\infty} (-c/b)_{\infty} (-d/a)_{\infty} (-dq/b)_{\infty}}. \quad (2.9)$$

Changing n to $n+1$ in the first summation of the above identity and then multiplying both sides by $(1+d/b)^{-1}$, we find that

$$\begin{aligned} &\frac{1}{(1+d/b)} \sum_{n=0}^{\infty} \frac{(-b/d)_{n+1}}{(-c/a)_{n+1}} \left(-\frac{d}{a}\right)^{n+1} + \sum_{n=0}^{\infty} \frac{(-aq/c)_n}{(-d/b)_{n+1}} \left(-\frac{c}{b}\right)^n \\ &= \left(1 - \frac{b}{a}\right) \frac{(cd/ab)_{\infty} (bq/a)_{\infty} (aq/b)_{\infty} (q)_{\infty}}{(-c/a)_{\infty} (-c/b)_{\infty} (-d/a)_{\infty} (-d/b)_{\infty}}. \end{aligned} \quad (2.10)$$

Using (1.10) with $\alpha = -bq/c$, $\beta = q$, $\gamma = -dq/a$, and $z = -c/a$ in the first summation of the above identity and then multiplying both sides by $1/b$, we get

$$\begin{aligned} & \frac{1}{b} \sum_{n=0}^{\infty} \frac{(-aq/c)_n}{(-d/b)_{n+1}} \left(-\frac{c}{b}\right)^n - \frac{1}{a} \sum_{n=0}^{\infty} \frac{(-bq/c)_n}{(-d/a)_{n+1}} \left(-\frac{c}{a}\right)^n \\ &= \left(\frac{1}{b} - \frac{1}{a}\right) \frac{(cd/ab)_{\infty} (bq/a)_{\infty} (aq/b)_{\infty} (q)_{\infty}}{(-c/a)_{\infty} (-c/b)_{\infty} (-d/a)_{\infty} (-d/b)_{\infty}}. \end{aligned} \quad (2.11)$$

Substituting (2.11) in (2.8), we see that

$$\begin{aligned} & \left(\frac{1}{b} - \frac{1}{a}\right) \frac{(cd/ab)_{\infty} (c)_{\infty} (d)_{\infty} (bq/a)_{\infty} (aq/b)_{\infty} (q)_{\infty}}{(-aq)_{\infty} (-bq)_{\infty} (-c/a)_{\infty} (-c/b)_{\infty} (-d/a)_{\infty} (-d/b)_{\infty}} \\ &= \left(1 + b^{-1}\right) \sum_{m=0}^{\infty} \frac{(-aq/c)_m (d)_m}{(-aq)_m (-d/b)_{m+1}} \left(-\frac{c}{b}\right)^m - \left(1 + a^{-1}\right) \sum_{m=0}^{\infty} \frac{(-bq/c)_m (d)_m}{(-bq)_m (-d/a)_{m+1}} \left(-\frac{c}{a}\right)^m. \end{aligned} \quad (2.12)$$

Now setting $\alpha = -cd/b$, $\beta = cd/ab$, $\gamma = c$, $\delta = q$, and $\epsilon = d$ in (1.9) and then multiplying both sides by $1/(1 + d/b)(1 + c/b)$, we obtain

$$\begin{aligned} & \sum_{n=0}^{\infty} \frac{(cd/ab)_n (c)_n (d)_n (1 + cdq^{2n}/b) q^{n(n+1)/2} (-1)^n a^n b^{-n}}{(-aq)_n (-c/b)_{n+1} (-d/b)_{n+1}} \\ &= \sum_{n=0}^{\infty} \frac{(-aq/c)_n (d)_n}{(-aq)_n (-d/b)_{n+1}} \left(-\frac{c}{b}\right)^n. \end{aligned} \quad (2.13)$$

Interchanging a and b in (2.13), we have

$$\sum_{n=0}^{\infty} \frac{(cd/ab)_n (c)_n (d)_n (1 + cdq^{2n}/a) q^{n(n+1)/2} (-1)^n b^n a^{-n}}{(-bq)_n (-c/a)_{n+1} (-d/a)_{n+1}} = \sum_{n=0}^{\infty} \frac{(-bq/c)_n (d)_n}{(-bq)_n (-d/a)_{n+1}} \left(-\frac{c}{a}\right)^n. \quad (2.14)$$

Substituting (2.13) and (2.14) in (2.12), we deduce (1.7).

Theorem 2.1 (A four-variable generalization of Jacobi's triple product identity). For $|c|, |d| < |a|, |b| < 1$,

$$\begin{aligned} & \frac{(cd/ab)_{\infty} (c)_{\infty} (d)_{\infty} (b/a)_{\infty} (aq/b)_{\infty} (q)_{\infty}}{(-a)_{\infty} (-b)_{\infty} (-c/a)_{\infty} (-c/b)_{\infty} (-d/a)_{\infty} (-d/b)_{\infty}} \\ &= \sum_{m=0}^{\infty} \frac{(d)_m (-cq^{-m}/a)_m (-1)^m a^m b^{-m} q^{m(m+1)/2}}{(-a)_{m+1} (-d/b)_{m+1}} \\ &\quad - \sum_{m=0}^{\infty} \frac{(d)_m (-cq^{-m}/b)_m (-1)^m a^{-(m+1)} b^{m+1} q^{m(m+1)/2}}{(-b)_{m+1} (-d/a)_{m+1}}. \end{aligned} \quad (2.15)$$

Proof. Employing

$$\begin{aligned}\left(-\frac{aq}{c}\right)_m &= \left(\frac{a}{c}\right)^m q^{m(m+1)/2} \left(-\frac{cq^{-m}}{a}\right)_m, \\ \left(-\frac{bq}{c}\right)_m &= \left(\frac{b}{c}\right)^m q^{m(m+1)/2} \left(-\frac{cq^{-m}}{b}\right)_m\end{aligned}\tag{2.16}$$

in the right side of (2.12) and then multiplying both sides by $b/(1+a)(1+b)$, we obtain (2.15). $\square$

Acknowledgment

The authors thank the anonymous referee for several helpful comments.

References

- [1] S. Ramanujan, *The Lost Notebook and Other Unpublished Papers*, Springer, Berlin, Germany, 1988.
- [2] S.-Y. Kang, "Generalizations of Ramanujan's reciprocity theorem and their applications," *Journal of the London Mathematical Society*, vol. 75, no. 1, pp. 18–34, 2007.
- [3] G. E. Andrews, "Ramanujan's "lost" note book I: partial θ —functions," *Advances in Mathematics*, vol. 41, pp. 137–172, 1981.
- [4] G. E. Andrews, "Ramanujan's "lost" notebook. V. Euler's partition identity," *Advances in Mathematics*, vol. 61, no. 2, pp. 156–164, 1986.
- [5] D. D. Somashekara and S. N. Fathima, "An interesting generalization of Jacobi's triple product identity," *Far East Journal of Mathematical Sciences*, vol. 9, no. 3, pp. 255–259, 2003.
- [6] B. C. Berndt, *Ramanujan's Notebooks. Part III*, Springer, New York, NY, USA, 1991.
- [7] G. E. Andrews, R. Askey, and R. Roy, *Special Functions*, vol. 71 of *Encyclopedia of Mathematics and Its Applications*, Cambridge University Press, Cambridge, UK, 1999.
- [8] G. Gasper and M. Rahman, *Basic Hypergeometric Series*, vol. 96 of *Encyclopedia of Mathematics and Its Applications*, Cambridge University Press, Cambridge, UK, 2nd edition, 2004.
- [9] B. C. Berndt, S. H. Chan, B. P. Yeap, and A. J. Yee, "A reciprocity theorem for certain q -series found in Ramanujan's lost notebook," *Ramanujan Journal*, vol. 13, no. 1–3, pp. 27–37, 2007.
- [10] N. J. Fine, *Basic Hypergeometric Series and Applications*, vol. 27 of *Mathematical Surveys and Monographs*, American Mathematical Society, Providence, RI, USA, 1988.
- [11] T. Kim, D. D. Somashekara, and S. N. Fathima, "On a generalization of Jacobi's triple product identity and its applications," *Advanced Studies in Contemporary Mathematics*, vol. 9, no. 2, pp. 165–174, 2004.
- [12] P. S. Guruprasad and N. Pradeep, "A simple proof of Ramanujan's reciprocity theorem," *Proceedings of the Jangjeon Mathematical Society*, vol. 9, no. 2, pp. 121–124, 2006.
- [13] C. Adiga and N. Anitha, "On a reciprocity theorem of Ramanujan," *Tamsui Oxford Journal of Mathematical Sciences*, vol. 22, no. 1, pp. 9–15, 2006.
- [14] M. E. H. Ismail, "A simple proof of Ramanujan's ${}_1\psi_1$ sum," *Proceedings of the American Mathematical Society*, vol. 63, no. 1, pp. 185–186, 1977.
- [15] C. Adiga, B. C. Berndt, S. Bhargava, and G. N. Watson, "Chapter 16 of Ramanujan's second notebook: theta-functions and q -series," *Memoirs of the American Mathematical Society*, vol. 53, no. 315, 1985.
- [16] C. Adiga and P. S. Guruprasad, "On a three variable reciprocity theorem," *South East Asian Journal of Mathematics and Mathematical Sciences*, vol. 6, no. 2, pp. 57–61, 2008.
- [17] D. D. Somashekara and D. Mamta, "A note on three variable reciprocity theorem," to appear in *International Journal of Pure and Applied Mathematics*, 2009.
- [18] D. D. Somashekara and D. Mamta, "On the three variable reciprocity theorem and its applications," to appear in *The Australian Journal of Mathematical Analysis and Applications*.
- [19] Z. Zhang, "An identity related to Ramanujan's and its applications," to appear in *Indian Journal of Pure and Applied Mathematics*.

- [20] X. R. Ma, "Six-variable generalization of Ramanujan's reciprocity theorem and its variants," *Journal of Mathematical Analysis and Applications*, vol. 353, no. 1, pp. 320–328, 2009.
- [21] X. R. Ma, "A five-variable generalization of Ramanujan's reciprocity theorem and its applications," to appear.
- [22] G. E. Andrews, "A simple proof of Jacobi's triple product identity," *Proceedings of the American Mathematical Society*, vol. 16, pp. 333–334, 1965.
- [23] E. M. Wright, "An enumerative proof of an identity of Jacobi," *Journal of the London Mathematical Society*, vol. 40, pp. 55–57, 1965.
- [24] M. S. Cheema, "Vector partitions and combinatorial identities," *Mathematics of Computation*, vol. 18, pp. 414–420, 1964.
- [25] C. Sudler Jr., "Two enumerative proofs of an identity of Jacobi," *Proceedings of the Edinburgh Mathematical Society*, vol. 15, pp. 67–71, 1966.
- [26] G. H. Hardy and E. M. Wright, *An Introduction to the Theory of Numbers*, Oxford University Press, New York, NY, USA, 5th edition, 1979.
- [27] M. D. Hirschhorn, "A simple proof of Jacobi's two-square theorem," *The American Mathematical Monthly*, vol. 92, no. 8, pp. 579–580, 1985.
- [28] M. D. Hirschhorn, "A simple proof of Jacobi's four-square theorem," *Proceedings of the American Mathematical Society*, vol. 101, no. 3, pp. 436–438, 1987.
- [29] S. Corteel and J. Lovejoy, "Overpartitions," *Transactions of the American Mathematical Society*, vol. 356, no. 4, pp. 1623–1635, 2004.