Research Article

Unbounded Conditional Expectations for Partial O^* -Algebras

Mayumi Takakura

Department of Applied Mathematics, Fukuoka University, Fukuoka 814-0180, Japan

Correspondence should be addressed to Mayumi Takakura, mayumi@fukuoka-u.ac.jp

Received 18 September 2008; Accepted 26 January 2009

Recommended by Ingo Witt

The main purpose of this paper is to generalize studies of unbounded conditional expectations for O^* -algebras to those for partial O^* -algebras.

Copyright © 2009 Mayumi Takakura. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

In probability theory, conditional expectations play a fundamental role. Conditional expectations for von Neumann algebra have been studied in noncommutative probability theory. In particular, Takesaki [1] characterized the existence of conditional expectation using Tomita's modular theory. Thus a conditional expectation does not necessarily exist for a general von Neumann algebra. The study of conditional expectations for O^* -algebras was begun by Gudder and Hudson [2]. After that, in [3, 4] we have investigated an unbounded conditional expectation which is a positive linear map $\mathcal{E}$ of an O^* -algebra $\mathcal{M}$ onto a given O^* -subalgebra $\mathcal{N}$ of $\mathcal{M}$. In this paper we will consider conditional expectations for partial O^* -algebras. Suppose that $\mathcal{M}$ is a self-adjoint partial O^* -algebra containing identity I on dense subspace $\mathfrak{D}$ of Hilbert space $\mathcal{H}$ with a strongly cyclic vector ξ_0 , and $\mathcal{N}$ is a partial O^* -subalgebra of $\mathcal{M}$ such that $(\mathcal{N} \cap R^w(\mathcal{M}))\xi_0$ is dense in $\mathcal{H}_{\mathcal{N}} \equiv \overline{\mathcal{N}\xi_0}$, where $R^w(\mathcal{M})$ is the set of all right multiplier of $\mathcal{M}$. The definitions of (self-adjoint) partial O^* -algebra and a strongly cyclic vector are stated in Section 2. A map $\mathcal{E}$ of $\mathcal{M}$ onto $\mathcal{N}$ is said to be a *weak conditional-expectation* of $(\mathcal{M}, \xi_0)$ with respect to, $\mathcal{N}$ if it satisfies $(AX\xi_0 \mid Y\xi_0) = (\mathcal{E}(A)X\xi_0 \mid Y\xi_0)$, for all $A \in \mathcal{M}$, for all $X, Y \in \mathcal{N} \cap R^w(\mathcal{M})$; but, the range $\mathcal{E}(A)$ of the weak conditional-expectation $\mathcal{E}$ is not necessarily contained in $\mathcal{N}$, and so we have considered a map $\mathcal{E}$ of $\mathcal{M}$ onto $\mathcal{N}$ satisfying the following:

- (i) the domain $D(\mathcal{E})$ of $\mathcal{E}$ is a $\dagger$ -invariant subspace of $\mathcal{M}$ containing $\mathcal{N}$;
- (ii) $\mathcal{E}$ is a projection; that is, it is hermitian $(\mathcal{E}(A))^\dagger = \mathcal{E}(A^\dagger)$, for all $A \in D(\mathcal{E})$ and $\mathcal{E}(X) = X$, for all $X \in \mathcal{N}$;

- (iii) $\mathcal{E}(A \square X) = \mathcal{E}(A) \square X$, for all $A \in D(\mathcal{E})$, for all $X \in \mathcal{N} \cap R^w(\mathcal{M})$, $\mathcal{E}(X \square A) = X \square \mathcal{E}(A)$, for all $A \in D(\mathcal{E}) \cap R^w(\mathcal{N})$, for all $X \in \mathcal{N}$;
- (iv) $\omega_{\xi_0}(\mathcal{E}(A)) = \omega_{\xi_0}(A)$, for all $A \in D(\mathcal{E})$, where ω_{ξ_0} is a state on $\mathcal{M}$ defined by $\omega_{\xi_0}(A) = (A\xi_0 \mid \xi_0)$, $A \in \mathcal{M}$;

and call it an *unbounded conditional expectation* of $(\mathcal{M}, \xi_0)$ with respect to, $\mathcal{N}$. In particular, if $D(\mathcal{E}) = \mathcal{M}$, then $\mathcal{E}$ is said to be a *conditional expectation* of $(\mathcal{M}, \xi_0)$ with respect to, $\mathcal{N}$.

Finally, we will investigate the scale of the domain of unbounded conditional expectations of partial GW^* -algebra which is unbounded generalizations of von Neumann algebras.

2. Preliminaries

In this section we review the definitions and the basic theory of partial O^* -algebras, partial GW^* -algebras and partial EW^* -algebras. For more details, refer to [5].

A *partial $\ast$ -algebra* is a complex vector space $\mathfrak{A}$ with an involution $x \rightarrow x^*$ and a subset $\Gamma \subset \mathfrak{A} \times \mathfrak{A}$ such that

- (i) $(x, y) \in \Gamma$ implies $(y^*, x^*) \in \Gamma$;
- (ii) $(x, y_1), (x, y_2) \in \Gamma$ implies $(x, \lambda y_1 + \mu y_2) \in \Gamma$, for all $\lambda, \mu \in \mathbb{C}$;
- (iii) whenever $(x, y) \in \Gamma$, there exists a product $x \cdot y \in \mathfrak{A}$ with the usual properties of the multiplication: $x \cdot (y + \lambda z) = x \cdot y + \lambda(x \cdot z)$ and $(x \cdot y)^* = y^* \cdot x^*$ for $(x, y), (x, z) \in \Gamma$ and $\lambda \in \mathbb{C}$.

The element e of the $\mathfrak{A}$ is called a *unit* if $e^* = e$, $(e, x) \in \Gamma$ for all $x \in \mathfrak{A}$, and $e \cdot x = x \cdot e = x$, for all $x \in \mathfrak{A}$. Notice that the partial multiplication is not required to be associative. Whenever $(x, y) \in \Gamma$, x is called a *left multiplier* of y and y is called a *right multiplier* of x , and we write $x \in L(y)$ and $y \in R(x)$. For a subset $\mathcal{B} \subset \mathfrak{A}$, we write

$$L(\mathcal{B}) = \bigcap_{x \in \mathcal{B}} L(x), \quad R(\mathcal{B}) = \bigcap_{x \in \mathcal{B}} R(x). \quad (2.1)$$

Let $\mathcal{H}$ be a Hilbert space with inner product $(\cdot \mid \cdot)$ and $\mathfrak{D}$ a dense subspace of $\mathcal{H}$. We denote by $\mathcal{L}^\dagger(\mathfrak{D}, \mathcal{H})$ the set of all closable linear operators X such that $\mathfrak{D}(X) = \mathfrak{D}$, $\mathfrak{D}(X^*) \supseteq \mathfrak{D}$. The set $\mathcal{L}^\dagger(\mathfrak{D}, \mathcal{H})$ is a partial $\ast$ -algebra with respect to the following operations: the usual sum $X + Y$, the scalar multiplication λX , the involution $X \rightarrow X^\dagger (= X^* \upharpoonright \mathfrak{D})$, and the weak partial multiplication $X \square Y \equiv X^\dagger Y$, defined whenever Y is a weak right multiplier of X ($X \in L^w(Y)$ or $Y \in R^w(X)$), that is, if and only if $Y\mathfrak{D} \subset \mathfrak{D}(X^\dagger)$ and $X^*\mathfrak{D} \subset \mathfrak{D}(Y^*)$. A partial $\ast$ -subalgebra of $\mathcal{L}^\dagger(\mathfrak{D}, \mathcal{H})$ is called a *partial O^* -algebra* on $\mathfrak{D}$.

Let $\mathcal{M}$ be a partial O^* -algebra on $\mathfrak{D}$. The locally convex topology on $\mathfrak{D}$ defined by the family $\{\|\cdot\|_X; X \in \mathcal{M}\}$ of seminorms $\|\xi\|_X = \|\xi\| + \|X\xi\|$, $\xi \in \mathfrak{D}$ is called the *graph topology* on $\mathfrak{D}$ and denoted by $t_{\mathcal{M}}$. The completion of $\mathfrak{D}[t_{\mathcal{M}}]$ is denoted by $\tilde{\mathfrak{D}}[t_{\mathcal{M}}]$. If the locally convex space $\mathfrak{D}[t_{\mathcal{M}}]$ is complete, then $\mathcal{M}$ is called *closed*. We also define the following domains:

$$\begin{aligned} \widehat{\mathfrak{D}}(\mathcal{M}) &= \bigcap_{X \in \mathcal{M}} \overline{\mathfrak{D}(X)}, & \mathfrak{D}^*(\mathcal{M}) &= \bigcap_{X \in \mathcal{M}} \mathfrak{D}(X^*), \\ \mathfrak{D}^{**}(\mathcal{M}) &= \bigcap_{X \in \mathcal{M}} \mathfrak{D}((X^* \upharpoonright \mathfrak{D}^*(\mathcal{M}))^*), \end{aligned} \quad (2.2)$$

and then

$$\mathfrak{D} \subset \tilde{\mathfrak{D}}(\mathcal{M}) \subset \widehat{\mathfrak{D}}(\mathcal{M}) \subset \mathfrak{D}^{**}(\mathcal{M}) \subset \mathfrak{D}^*(\mathcal{M}). \quad (2.3)$$

The partial O^* -algebra $\mathcal{M}$ is called *fully closed* if $\mathfrak{D} = \widehat{\mathfrak{D}}(\mathcal{M})$, *self-adjoint* if $\mathfrak{D} = \mathfrak{D}^*(\mathcal{M})$, *essentially self-adjoint* if $\mathfrak{D}^*(\mathcal{M}) = \widehat{\mathfrak{D}}(\mathcal{M})$, and *algebraically self-adjoint* if $\mathfrak{D}^*(\mathcal{M}) = \mathfrak{D}^{**}(\mathcal{M})$.

We defined two weak commutants of $\mathcal{M}$. The *weak bounded commutant* $\mathcal{M}'_w$ of $\mathcal{M}$ is the set

$$\mathcal{M}'_w = \{C \in \mathcal{B}(\mathcal{H}); (CX\xi \mid \eta) = (C\xi \mid X^\dagger \eta) \text{ for every } X \in \mathcal{M}, \xi, \eta \in \mathfrak{D}\}; \quad (2.4)$$

but the partial multiplication is not required to be associative, so we define the *quasi-weak bounded commutant* $\mathcal{M}'_{qw}$ of $\mathcal{M}$ as the set

$$\mathcal{M}'_{qw} = \{C \in \mathcal{M}'_w; (CX_1^\dagger \xi \mid X_2 \eta) = (C\xi \mid (X_1 \square X_2) \eta) \ \forall X_1 \in L(X_2), \xi, \eta \in \mathfrak{D}\}. \quad (2.5)$$

In general, $\mathcal{M}'_{qw} \subsetneq \mathcal{M}'_w$.

A **-representation* of a partial $*$ -algebra $\mathfrak{A}$ is a $*$ -homomorphism of $\mathfrak{A}$ into $\mathcal{L}^+(\mathfrak{D}, \mathcal{H})$, satisfying $\pi(e) = I$ whenever $e \in \mathfrak{A}$, that is,

- (i) π is linear;
- (ii) $x \in L^w(y)$ in $\mathfrak{A}$ implies $\pi(x) \in L^w(\pi(y))$ and $\pi(x) \square \pi(y) = \pi(xy)$;
- (iii) $\pi(x^*) = \pi(x)^\dagger$ for every $x \in \mathfrak{A}$.

Let π be a $*$ -representation of a partial $*$ -algebra $\mathfrak{A}$ into $\mathcal{L}^+(\mathfrak{D}, \mathcal{H})$. Then we define

$\tilde{\mathfrak{D}}(\pi)$: the completion of $\mathfrak{D}$ with respect to the graph topology $t_{\pi(\mathfrak{A})}$,

$$\begin{aligned} \tilde{\pi}(x) &= \overline{\pi(x)}|_{\tilde{\mathfrak{D}}(\pi)}, \quad x \in \mathfrak{A}; \\ \widehat{\mathfrak{D}}(\pi) &= \bigcap_{x \in \mathfrak{A}} \overline{\mathfrak{D}(\pi(x))}, \\ \widehat{\pi}(x) &= \overline{\pi(x)}|_{\widehat{\mathfrak{D}}(\pi)}, \quad x \in \mathfrak{A}; \\ \mathfrak{D}^*(\pi) &= \bigcap_{x \in \mathfrak{A}} \mathfrak{D}(\pi(x)^*), \\ \pi^*(x) &= \pi(x^*)^*|_{\mathfrak{D}^*(\pi)}, \quad x \in \mathfrak{A}. \end{aligned} \quad (2.6)$$

We say that π is *closed* if $\mathfrak{D} = \tilde{\mathfrak{D}}(\pi)$; *fully closed* if $\mathfrak{D} = \widehat{\mathfrak{D}}(\pi)$; *essentially self-adjoint* if $\widehat{\mathfrak{D}}(\pi) = \mathfrak{D}^*(\pi)$; and *self-adjoint* if $\mathfrak{D} = \mathfrak{D}^*(\pi)$.

We introduce the weak and the quasi-weak commutants of a $*$ -representaion π of a partial $*$ -algebra $\mathfrak{A}$ as follows:

$$\begin{aligned}\pi(\mathfrak{A})'_w &= \{C \in \mathcal{B}(\mathcal{H}); (C\xi \mid \pi(x)\eta) = (C\pi(x^*)\xi \mid \eta), \forall x \in \mathfrak{A}, \xi, \eta \in \mathfrak{D}(\pi)\}, \\ \mathcal{C}_{qw}(\pi) &= \{C \in \pi(\mathfrak{A})'_w; (C\pi(x_1^*)\xi \mid \pi(x_2)\eta) = (C\xi \mid \pi(x_1x_2)\eta), \\ &\quad \forall x_1, x_2 \in \mathfrak{A} \text{ such that } x_1 \in L(x_2), \text{ and all } \xi, \eta \in \mathfrak{D}(\pi)\},\end{aligned}\tag{2.7}$$

respectively.

We define the notion of strongly cyclic vector for a partial O^* -algebra $\mathcal{M}$ on $\mathfrak{D}$ in $\mathcal{H}$. A vector ξ_0 in $\mathfrak{D}$ is said to be *strongly cyclic* if $R^w(\mathcal{M})\xi_0$ is dense in $\mathfrak{D}[t_{\mathcal{M}}]$, and ξ_0 is said to be *separating* if $\overline{\mathcal{M}'_w\xi_0} = \mathcal{H}$, where $R^w(\mathcal{M}) = \{Y \in \mathcal{M}; X \square Y \text{ is well-defined, for all } X \in \mathcal{M}\}$.

We introduce the notion of partial GW^* -algebras and partial EW^* -algebras which are unbounded generalizations of von Neumann algebras. A fully closed partial O^* -algebra $\mathcal{M}$ on $\mathfrak{D}$ is called a *partial GW^* -algebra* if there exists a von Neumann algebra $\mathcal{M}_0$ on $\mathcal{H}$ such that $\mathcal{M}'_0\mathfrak{D} \subset \mathfrak{D}$ and $\mathcal{M} = [\mathcal{M}_0[\mathfrak{D}]]^{s*}$. A partial O^* -algebra $\mathcal{M}$ on $\mathfrak{D}$ is said to be a partial EW^* -algebra if $\overline{\mathcal{M}_b} \equiv \{A \in \mathcal{B}(\mathcal{H}); A[\mathfrak{D} \in \mathcal{M}] \}$ is a von Neumann algebra, $\mathcal{M}_b\mathfrak{D} \subset \mathfrak{D}$ and $\overline{\mathcal{M}'_b}\mathfrak{D} \subset \mathfrak{D}$.

3. Weak Conditional Expectations

In this section, let $\mathcal{M}$ be a self-adjoint partial O^* -algebra containing the identity I on $\mathfrak{D}$ in $\mathcal{H}$ with a strongly cyclic vector ξ_0 and let $\mathcal{N}$ be a partial O^* -subalgebra of $\mathcal{M}$ such that

$$(N) \quad (\mathcal{N} \cap R^w(\mathcal{M}))\xi_0 \text{ is dense in } \mathcal{H}_{\mathcal{N}} \equiv \overline{\mathcal{N}\xi_0}.$$

The following is easily shown.

Lemma 3.1. *Put*

$$\begin{aligned}\mathfrak{D}(\pi_{\mathcal{N}}) &= (\mathcal{N} \cap R^w(\mathcal{M}))\xi_0, \\ \pi_{\mathcal{N}}(X)Y\xi_0 &= (X \square Y)\xi_0, \quad \forall X \in \mathcal{N}, \forall Y \in \mathcal{N} \cap R^w(\mathcal{M}).\end{aligned}\tag{3.1}$$

Then $\pi_{\mathcal{N}}$ is a $*$ -representations of $\mathcal{N}$ in the Hilbert space $\mathcal{H}_{\mathcal{N}} \equiv \overline{\mathfrak{D}(\pi_{\mathcal{N}})}$.

We denote by $P_{\mathcal{N}}$ the projection of $\mathcal{H}$ onto $\mathcal{H}_{\mathcal{N}} \equiv \overline{\mathfrak{D}(\pi_{\mathcal{N}})}$. This projection $P_{\mathcal{N}}$ plays an important role in this reserch. First we have the following.

Lemma 3.2. *It holds that $P_{\mathcal{N}}\mathfrak{D} \subset \mathfrak{D}^*(\pi_{\mathcal{N}})$ and $\pi_{\mathcal{N}}^*(X)P_{\mathcal{N}}\xi = P_{\mathcal{N}}X\xi$, for all $X \in \mathcal{N}$ and for all $\xi \in \mathfrak{D}$.*

Proof. Take arbitrary $X \in \mathcal{N}$ and $\xi \in \mathfrak{D}$. For any $Y \in \mathcal{N} \cap R^w(\mathcal{M})$, we have

$$(\pi_{\mathcal{N}}(X^\dagger)Y\xi_0 \mid P_{\mathcal{N}}\xi) = ((X^\dagger \square Y)\xi_0 \mid P_{\mathcal{N}}\xi) = (X^\dagger Y\xi_0 \mid \xi) = (Y\xi_0 \mid X\xi) = (Y\xi_0 \mid P_{\mathcal{N}}X\xi), \tag{3.2}$$

and so $P_{\mathcal{N}}\mathfrak{D} \subset \mathfrak{D}^*(\pi_{\mathcal{N}})$ and $\pi_{\mathcal{N}}^*(X)P_{\mathcal{N}}\xi = P_{\mathcal{N}}X\xi$. □

Definition 3.3. A map $\mathcal{E}$ of $\mathcal{M}$ into $\mathcal{L}^+(\mathfrak{D}(\pi_{\mathcal{N}}), \mathcal{H}_{\mathcal{N}})$ is said to be a weak conditional-expectation of $(\mathcal{M}, \xi_0)$ with respect to, $\mathcal{N}$ if it satisfies

$$(\mathcal{E}(AX\xi_0) \mid Y\xi_0) = (\mathcal{E}(A)X\xi_0 \mid Y\xi_0), \quad \forall A \in \mathcal{M}, \forall X, Y \in \mathcal{N} \cap R^w(\mathcal{M}). \quad (3.3)$$

For weak conditional-expectation we have the following.

Theorem 3.4. *There exists a unique weak conditional-expectation $\mathcal{E}(\cdot \mid \mathcal{N})$ of $(\mathcal{M}, \xi_0)$ with respect to, $\mathcal{N}$, and*

$$\mathcal{E}(A \mid \mathcal{N}) = P_{\mathcal{N}}A[\mathfrak{D}(\pi_{\mathcal{N}})], \quad \forall A \in \mathcal{M}. \quad (3.4)$$

The weak conditional-expectation $\mathcal{E}(\cdot \mid \mathcal{N})$ of $(\mathcal{M}, \xi_0)$ with respect to, $\mathcal{N}$ satisfies the following:

- (i) $\mathcal{E}(\cdot \mid \mathcal{N})$ is linear,
- (ii) $\mathcal{E}(\cdot \mid \mathcal{N})$ is hermitian, that is, $\mathcal{E}(A \mid \mathcal{N})^\dagger = \mathcal{E}(A^\dagger \mid \mathcal{N})$, for all $A \in \mathcal{M}$,
- (iii) $\mathcal{E}(X \mid \mathcal{N}) = X[\mathfrak{D}(\pi_{\mathcal{N}})]$, for all $X \in \mathcal{N}$,
- (iv) $\mathcal{E}(A^\dagger \square A \mid \mathcal{N}) \geq 0$, for all $A \in \mathcal{M}$ s.t. $A^\dagger \square A$ is well-defined,
- (v) $\mathcal{E}(A \mid \mathcal{N})^\dagger \square \mathcal{E}(A \mid \mathcal{N}) \leq \mathcal{E}(A^\dagger \square A \mid \mathcal{N})$, for all $A \in \mathcal{M}$ s.t. $A^\dagger \square A$ and $\mathcal{E}(A \mid \mathcal{N})^\dagger \square \mathcal{E}(A \mid \mathcal{N})$ are well-defined,
- (vi) $\mathcal{E}(A \mid \mathcal{N}) \square \pi_{\mathcal{N}}(X)$ is well-defined for any $A \in \mathcal{M}$ and $X \in \mathcal{N} \cap R^w(\mathcal{M})$, and $\mathcal{E}(A \mid \mathcal{N}) \square \pi_{\mathcal{N}}(X) = \mathcal{E}(A \square X \mid \mathcal{N})$,
- (vii) $\pi_{\mathcal{N}}(X) \square \mathcal{E}(A \mid \mathcal{N})$ is well-defined for any $A \in \mathcal{M} \cap R^w(\mathcal{N})$ and for all $X \in \mathcal{N}$, and $\pi_{\mathcal{N}}(X) \square \mathcal{E}(A \mid \mathcal{N}) = \mathcal{E}(X \square A \mid \mathcal{N})$,
- (viii) $\omega_{\xi_0}(\mathcal{E}(A \mid \mathcal{N})) = \omega_{\xi_0}(A)$, for all $A \in \mathcal{M}$.

Proof. We put

$$\mathcal{E}(A \mid \mathcal{N}) = P_{\mathcal{N}}A[\mathfrak{D}(\pi_{\mathcal{N}})], \quad \forall A \in \mathcal{M}. \quad (3.5)$$

By Lemma 3.2, $\mathcal{E}(A \mid \mathcal{N})$ is a linear map of $\mathfrak{D}(\pi_{\mathcal{N}})$ into $\mathfrak{D}^*(\pi_{\mathcal{N}})$ for any $A \in \mathcal{M}$, and furthermore we have $\mathcal{E}(A \mid \mathcal{N})^\dagger = \mathcal{E}(A^\dagger \mid \mathcal{N})$, for all $A \in \mathcal{M}$, so $\mathcal{E}(\cdot \mid \mathcal{N})$ is a map of $\mathcal{M}$ into $\mathcal{L}^+(\mathfrak{D}(\pi_{\mathcal{N}}), \mathcal{H}_{\mathcal{N}})$.

Since

$$(\mathcal{E}(A \mid \mathcal{N})X\xi_0 \mid Y\xi_0) = (P_{\mathcal{N}}AX\xi_0 \mid Y\xi_0) = (AX\xi_0 \mid Y\xi_0) \quad (3.6)$$

for each $A \in \mathcal{M}$, $X, Y \in \mathcal{N} \cap R^w(\mathcal{M})$, $\mathcal{E}(\cdot \mid \mathcal{N})$ is a weak conditional-expectation of $(\mathcal{M}, \xi_0)$ with respect to, $\mathcal{N}$. It is easily shown that if $\mathcal{E}$ is a weak conditional-expectation of $(\mathcal{M}, \xi_0)$ with respect to, $\mathcal{N}$, $\mathcal{E}(A) = \mathcal{E}(A \mid \mathcal{N})$ for each $A \in \mathcal{M}$. Thus the existence and uniqueness of weak conditional-expectations is shown. The statements (iii)–(viii) follow since $\mathcal{E}(A \mid \mathcal{N}) = P_{\mathcal{N}}A[\mathfrak{D}(\pi_{\mathcal{N}})]$, for all $A \in \mathcal{M}$. This completes the proof. $\square$

4. Unbounded Conditional Expectations for Partial O^* -Algebras

Let $\mathcal{M}$ be a self-adjoint partial O^* -algebra containing I on $\mathfrak{D}$ in $\mathcal{L}$ and let $\xi_0 \in \mathfrak{D}$ be a strongly cyclic and separating vector for $\mathcal{M}$ and suppose that $\mathcal{N} \ni I$ is a partial O^* -subalgebra of $\mathcal{M}$ satisfying (N): $(\mathcal{N} \cap R^w(\mathcal{M}))\xi_0$ is dense in $\mathcal{H}_{\xi_0}$. We introduce unbounded conditional expectations of $(\mathcal{M}, \xi_0)$ with respect to, $\mathcal{N}$.

Definition 4.1. A map $\mathcal{E}$ of $\mathcal{M}$ onto $\mathcal{N}$ is said to be an unbounded conditional expectation of $(\mathcal{M}, \xi_0)$ with respect to, $\mathcal{N}$ if

- (i) the domain $D(\mathcal{E})$ of $\mathcal{E}$ is a $\dagger$ -invariant subspace of $\mathcal{M}$ containing $\mathcal{N}$;
- (ii) $\mathcal{E}$ is a projection; that is, it is hermitian $(\mathcal{E}(A))^\dagger = \mathcal{E}(A^\dagger)$, for all $A \in D(\mathcal{E})$ and $\mathcal{E}(X) = X$, for all $X \in \mathcal{N}$;
- (iii) $\mathcal{E}(A \square X) = \mathcal{E}(A) \square X$, for all $A \in D(\mathcal{E})$, for all $X \in \mathcal{N} \cap R^w(\mathcal{M})$, $\mathcal{E}(X \square A) = X \square \mathcal{E}(A)$, for all $A \in D(\mathcal{E}) \cap R^w(\mathcal{N})$, for all $X \in \mathcal{N}$;
- (iv) $\omega_{\xi_0}(\mathcal{E}(A)) = \omega_{\xi_0}(A)$, for all $A \in D(\mathcal{E})$.

In particular, if $D(\mathcal{E}) = \mathcal{M}$, then $\mathcal{E}$ is said to be a *conditional expectation* of $(\mathcal{M}, \xi_0)$ with respect to, $\mathcal{N}$.

For unbounded conditional expectations we have the following.

Lemma 4.2. Let $\mathcal{E}$ be an unbounded conditional expectation of $(\mathcal{M}, \xi_0)$ with respect to, $\mathcal{N}$. Then,

$$\mathcal{E}(A)X\xi_0 = P_{\mathcal{N}}AX\xi_0 = \mathcal{E}(A | \mathcal{N})X\xi_0, \quad \forall A \in D(\mathcal{E}), \forall X \in \mathcal{N} \cap R^w(\mathcal{M}). \quad (4.1)$$

Proof. For all $A \in D(\mathcal{E})$ and $X, Y \in \mathcal{N} \cap R^w(\mathcal{M})$, we have

$$\begin{aligned} (\mathcal{E}(A)X\xi_0 | Y\xi_0) &= (\mathcal{E}(A \square X)\xi_0 | Y\xi_0) = (\mathcal{E}(Y^\dagger \square A \square X)\xi_0 | \xi_0) = ((Y^\dagger \square A \square X)\xi_0 | \xi_0) \\ &= (AX\xi_0 | Y\xi_0) = (AX\xi_0 | P_{\mathcal{N}}Y\xi_0) = (P_{\mathcal{N}}AX\xi_0 | Y\xi_0). \end{aligned} \quad (4.2)$$

Hence, $\mathcal{E}(A)X\xi_0 = P_{\mathcal{N}}AX\xi_0 = \mathcal{E}(A | \mathcal{N})X\xi_0$, for all $A \in D(\mathcal{E})$, for all $X \in \mathcal{N} \cap R^w(\mathcal{M})$. $\square$

Let $\mathfrak{E}$ be the set of all unbounded conditional expectations of $(\mathcal{M}, \xi_0)$ with respect to, $\mathcal{N}$. Then $\mathfrak{E}$ is an ordered set with the following order $\subset$:

$$\mathcal{E}_1 \subset \mathcal{E}_2 \quad \text{iff} \quad D(\mathcal{E}_1) \subset D(\mathcal{E}_2), \quad \mathcal{E}_1(A) = \mathcal{E}_2(A), \quad \forall A \in D(\mathcal{E}_1). \quad (4.3)$$

Theorem 4.3. There exists a maximal unbounded conditional expectation of $(\mathcal{M}, \xi_0)$ with respect to, $\mathcal{N}$, and it is denoted by $\mathcal{E}_{\mathcal{N}}$.

Proof. We put

$$D(\mathcal{E}_0) \equiv \{A \in \mathcal{M}; P_{\mathcal{N}}A[(\mathcal{N} \cap R^w(\mathcal{M}))\xi_0] \in \mathcal{N}[(\mathcal{N} \cap R^w(\mathcal{M}))\xi_0]\}. \quad (4.4)$$

Then, for any $A \in D(\mathcal{E}_0)$, there exists a unique map $\mathcal{E}_0$ such that

$$\mathcal{E}_0(A)X\xi_0 = P_{\mathcal{N}}AX\xi_0 = \mathcal{E}(A | \mathcal{N})X\xi_0, \quad \forall X \in \mathcal{N} \cap R^w(\mathcal{M}). \quad (4.5)$$

It is easily shown that $\mathcal{E}_0$ is an unbounded conditional expectation of $(\mathcal{M}, \xi_0)$ with respect to, $\mathcal{N}$. Furthermore, $\mathcal{E}_0$ is maximal in $\mathfrak{E}$. Indeed, let $\mathcal{E} \in \mathfrak{E}$. Take an arbitrary $A \in D(\mathcal{E})$. Then by Lemma 4.2 we have

$$\mathcal{E}(A)X\xi_0 = P_{\mathcal{N}}AX\xi_0 = \mathcal{E}(A | \mathcal{N})X\xi_0, \quad X \in \mathcal{N} \cap R^w(\mathcal{M}), \quad (4.6)$$

which implies $\mathcal{E}(A)X\xi_0 \in \mathcal{N}[(\mathcal{N} \cap R^w(\mathcal{M}))\xi_0]$. Hence $\mathcal{E} \subset \mathcal{E}_0$ and $\mathcal{E}_0$ is maximal in $\mathfrak{E}$. This completes the proof. $\square$

5. Existence of Conditional Expectations for Partial O^* -Algebras

Let $\mathcal{M}$ be a self-adjoint partial O^* -algebra containing I on $\mathfrak{D}$ in $\mathcal{H}$, $\xi_0 \in \mathfrak{D}$ be a strongly cyclic and separating vector for $\mathcal{M}$ and $\mathcal{N} \ni I$ a partial O^* -subalgebra of $\mathcal{M}$ such that

- (N) $(\mathcal{N} \cap R^w(\mathcal{M}))\xi_0$ is dense in $\mathcal{H}_{\mathcal{N}}$,
- (N₁) $\mathcal{N}'_w \widehat{\mathfrak{D}}(\mathcal{N}) \subset \widehat{\mathfrak{D}}(\mathcal{N})$,
- (N₂) $(\mathcal{N} \cap R^w(\mathcal{M}))\xi_0$ is essentially self-adjoint for $\mathcal{N}$,
- (N₃) $\Delta''_{\xi_0} \text{it}(\mathcal{N}'_w)' \Delta''_{\xi_0}^{-\text{it}} = (\mathcal{N}'_w)'$, for all $t \in \mathbb{R}$, where Δ''_{ξ_0} is the modular operator for the full Hilbert algebra $(\mathcal{M}'_w)'_{\xi_0}$.

Lemma 5.1. *It holds that $D(\mathcal{E}_{\mathcal{N}}) = \{A \in \mathcal{M}; P_{\mathcal{N}}A\xi_0 \in \mathcal{N}\xi_0\}$.*

Proof. We put

$$D(\mathcal{E}) = \{A \in \mathcal{M}; P_{\mathcal{N}}A\xi_0 \in \mathcal{N}\xi_0\}. \quad (5.1)$$

By Lemma 4.2, we have

$$P_{\mathcal{N}}A\xi_0 = \mathcal{E}_{\mathcal{N}}(A)\xi_0 \in \mathcal{N}\xi_0 \quad (5.2)$$

for each $A \in D(\mathcal{E}_{\mathcal{N}})$. Hence, $D(\mathcal{E}_{\mathcal{N}}) \subset D(\mathcal{E})$. We show the converse inclusion. Since ξ_0 is separating vector for $\mathcal{M}$, it follows that for any $A \in D(\mathcal{E})$, there exists a unique element $\mathcal{E}(A)$ of $\mathcal{N}$ such that $P_{\mathcal{N}}A\xi_0 = \mathcal{E}(A)\xi_0$. Indeed, since $\mathcal{E}_{\mathcal{N}}$ is maximal in $\mathfrak{E}$, it is sufficient to show that $\mathcal{E}$ is an unbounded conditional expectation of $(\mathcal{M}, \xi_0)$ with respect to, $\mathcal{N}$. By assumption (N₁) and [5, Proposition 2.3.5], we have

$$\overline{X} \text{ is affiliated with von Neumann algebra } (\mathcal{N}'_w)' \text{ for each } X \in \mathcal{N}, \quad (5.3)$$

$$\mathcal{N}'_w = \mathcal{N}'_{q^w}. \quad (5.4)$$

Since $\mathcal{M}$ is self-adjoint and $(\mathcal{N} \cap R^w(\mathcal{M}))\xi_0$ is dense in $\mathcal{H}_{\mathcal{N}}$, it follows that $(\mathcal{N} \cap R^w(\mathcal{M}))\xi_0$ is a reducing subspace for $\mathcal{N}$, that is,

$$\mathcal{N}(\mathcal{N} \cap R^w(\mathcal{M}))\xi_0 \subset \overline{(\mathcal{N} \cap R^w(\mathcal{M}))\xi_0} = \overline{\mathcal{N}\xi_0}, \quad (5.5)$$

which implies by assumption (N₂) and [5, Theorem 7.4.4] that

$$P_{\mathcal{N}} \in N'_{w'}, \quad P_{\mathcal{N}} \widehat{\mathfrak{D}}(\mathcal{N}) \subset \widehat{\mathfrak{D}}(\mathcal{N}). \quad (5.6)$$

Furthermore, by (5.3) and (5.6), we have

$$\overline{\mathcal{N}\xi_0} = \overline{(\mathcal{N}'_w)' \xi_0}, \quad \text{that is, } \mathcal{P}_{\mathcal{N}} = \mathcal{P}_{(\mathcal{N}'_w)'}. \quad (5.7)$$

Let S_{ξ_0} and S''_{ξ_0} be the closures of the maps:

$$\begin{aligned} S_{\xi_0} A \xi_0 &= A^\dagger \xi_0, \quad A \in \mathcal{M}, \\ S''_{\xi_0} B \xi_0 &= B^* \xi_0, \quad B \in (\mathcal{M}'_w)'. \end{aligned} \quad (5.8)$$

By (5.3) we have

$$S_{\xi_0} \subset S''_{\xi_0}. \quad (5.9)$$

Takesaki proved in [1] that assumption (N₃) implies

$$P_{(\mathcal{M}'_w)'} S''_{\xi_0} \subset S''_{\xi_0} P_{(\mathcal{M}'_w)'}. \quad (5.10)$$

and there exists a conditional expectation $\mathcal{E}''$ of the von Neumann algebra $((\mathcal{M}'_w)', \xi_0)$ with respect to, $(\mathcal{M}'_w)'$.

By (5.6), (5.9), and (5.10), we have

$$\begin{aligned} \mathcal{E}(A^\dagger) \xi_0 &= P_{\mathcal{N}} A^\dagger \xi_0 = P_{\mathcal{N}} S_{\xi_0} A \xi_0 = P_{\mathcal{N}} S''_{\xi_0} A \xi_0 \\ &= S''_{\xi_0} P_{\mathcal{N}} A \xi_0 = S''_{\xi_0} \mathcal{E}(A) \xi_0 = S_{\xi_0} \mathcal{E}(A) \xi_0 = \mathcal{E}(A)^\dagger \xi_0 \end{aligned} \quad (5.11)$$

for each $A \in D(\mathcal{E})$, which implies by the separateness of ξ_0 that $\mathcal{E}$ is hermitian.

It is clear that $\mathcal{E}(X) = X$, for all $X \in \mathcal{N}$. Take arbitrary $A \in D(\mathcal{E})$ and $X \in \mathcal{N} \cap L^w(\mathcal{M})$. Since

$$(P_{\mathcal{N}}(X \square A) \xi_0 \mid Y \xi_0) = (P_{\mathcal{N}} A \xi_0 \mid X^\dagger Y \xi_0) = (\mathcal{E}(A) \xi_0 \mid X^\dagger Y \xi_0) = ((X \square \mathcal{E}(A)) \xi_0 \mid Y \xi_0) \quad (5.12)$$

for each $Y \in \mathcal{N} \cap R^w(\mathcal{M})$, it follows that $X \square A \in D(\mathcal{E})$ and $\mathcal{E}(X \square A) = X \square \mathcal{E}(A)$. Furthermore, since $\mathcal{E}$ is hermitian, it follows that $A \square X \in D(\mathcal{E})$ and $\mathcal{E}(A \square X) = \mathcal{E}(A) \square X$ for each $A \in D(\mathcal{E})$ and $X \in \mathcal{N} \cap R^w(\mathcal{M})$. It is clear that $\omega_{\xi_0}(\mathcal{E}(A)) = \omega_{\xi_0}(A)$ for each $A \in D(\mathcal{E})$. Thus $\mathcal{E}$ is an unbounded conditional expectation of $(\mathcal{M}, \xi_0)$ with respect to, $\mathcal{N}$. This completes that proof. $\square$

By Lemma 5.1, we have the following.

Theorem 5.2. *Let $\mathcal{M}$ be a self-adjoint partial O^* -algebra containing I on $\mathfrak{D}$ in $\mathcal{L}$ and let $\xi_0 \in \mathfrak{D}$ be a strongly cyclic and separating vector for $\mathcal{M}$ and suppose that $\mathcal{N} \ni I$ is a partial O^* -subalgebra of $\mathcal{M}$ satisfying (N), (N_1) , (N_2) , and (N_3) . Then there exists a conditional expectation of $(\mathcal{M}, \xi_0)$ with respect to, $\mathcal{N}$ if and only if $P_{\mathcal{N}}\mathcal{M}\xi_0 = \mathcal{N}\xi_0$.*

It is important to investigate the scale of the domain of an unbounded conditional expectation. We consider the case of partial GW^* -algebras.

Theorem 5.3. *Let $\mathcal{M}$ be a partial GW^* -algebra on $\mathfrak{D}$ in $\mathcal{L}$ and let $\xi_0 \in \mathfrak{D}$ be a strongly cyclic and separating vector for $\mathcal{M}$ and suppose that $\mathcal{N}$ be a partial GW^* -subalgebra of $\mathcal{M}$ satisfying (N), (N_1) , (N_2) , and (N_3) .*

Then, $D(\mathcal{E}_{\mathcal{N}}) \supset$ linear span of $\{X \square A; X \in \mathcal{N}, A \in (\mathcal{M}'_{\mathcal{W}})' \text{ s.t. } X \square A \text{ and } X \square \mathcal{E}''(A) \text{ are well defined}\} \supset$ linear span of $(\mathcal{M}'_{\mathcal{W}})'$ and $\mathcal{N}$.

In particular, if $\mathcal{N}_{P_{\mathcal{N}}}$ is a partial GW^ -algebra on $P_{\mathcal{N}}\mathfrak{D}$, then $\mathcal{E}_{\mathcal{N}}$ is a conditional expectation of $(\mathcal{M}, \xi_0)$ with respect to, $\mathcal{N}$.*

Proof. Let $X \in \mathcal{N}$, and $A \in (\mathcal{M}'_{\mathcal{W}})'$ s.t. $X \square A$ and $X \square \mathcal{E}''(A)$ are all defined. Then, it follows since $\mathcal{N}$ is a partial GW^* -subalgebra of $\mathcal{M}$ that

$$P_{\mathcal{N}}(X \square A)\xi_0 = P_{\mathcal{N}}X^{\dagger*}A\xi_0 = X^{\dagger*}P_{\mathcal{N}}A\xi_0 = (X \square \mathcal{E}''(A))\xi_0 \in \mathcal{N}\xi_0, \quad (5.13)$$

which implies by Lemma 5.1 that $X \square A \in D(\mathcal{E}_{\mathcal{N}})$ and $P_{\mathcal{N}}(X \square A)\xi_0 = (X \square \mathcal{E}''(A))\xi_0$. Suppose that $\mathcal{N}_{P_{\mathcal{N}}}$ is a partial GW^* -algebra on $P_{\mathcal{N}}\mathfrak{D}$.

By the result of Takesaki [1] there exists a unique conditional expectation $\mathcal{E}''$ of the von Neumann algebra $(\mathcal{N}'_{\mathcal{W}})'$ such that $\mathcal{E}''(A_{\alpha})_{P_{\mathcal{N}}} = P_{\mathcal{N}}AP_{\mathcal{N}}$ for each $A \in (\mathcal{M}'_{\mathcal{W}})'$. Since $\mathcal{M}$ is a partial GW^* -algebra, for any $X \in \mathcal{M}$ there is a net $\{A_{\alpha}\} \in (\mathcal{M}'_{\mathcal{W}})'$ which converges strongly* to X . Then

$$\mathcal{E}''(A_{\alpha})_{P_{\mathcal{N}}} \in ((\mathcal{N}'_{\mathcal{W}})')_{P_{\mathcal{N}}} = ((\mathcal{N}_{P_{\mathcal{N}}})'_{\mathcal{W}})', \quad (5.14)$$

and $\mathcal{E}''(A_{\alpha})_{P_{\mathcal{N}}}$ converges strongly* to $P_{\mathcal{N}}X[P_{\mathcal{N}}\mathfrak{D}]$. Therefore, we have $P_{\mathcal{N}}X[P_{\mathcal{N}}\mathfrak{D}] \in \mathcal{N}$. Hence, $X \in D(\mathcal{E}_{\mathcal{N}})$ and $\mathcal{E}_{\mathcal{N}}$ is a conditional expectation of $(\mathcal{M}, \xi_0)$ with respect to, $\mathcal{N}$. This completes the proof. $\square$

Corollary 5.4. *Let $\mathcal{M}$ be a partial EW^* -algebra on $\mathfrak{D}$ in $\mathcal{L}$ and let $\xi_0 \in \mathfrak{D}$ be a strongly cyclic and separating vector for $\mathcal{M}$ and suppose that $\mathcal{N}$ be a partial EW^* -subalgebra of $\mathcal{M}$ satisfying (N_2) and (N_3) . Then,*

$$D(\mathcal{E}_{\mathcal{N}}) \supset \text{linear span of } \mathcal{M}_b\mathcal{N} \text{ and } \mathcal{N}\mathcal{M}_b. \quad (5.15)$$

Proof. Since $\mathcal{M}_b \subset R^w(\mathcal{M})$, it follows that $\mathcal{N} \cap R^w(\mathcal{M}) \supset \mathcal{N}_b$, and so clearly (N) holds. Furthermore, (N_1) holds since $\mathcal{N}'_{\mathcal{W}}\widehat{\mathfrak{D}}(\mathcal{N}) = \mathcal{N}_b\widehat{\mathfrak{D}}(\mathcal{N}) \subset \widehat{\mathfrak{D}}(\mathcal{N})$. This completes the proof. $\square$

We consider the case of the well-known Segal L^p -space defined by τ .

Example 5.5. Let $\mathcal{M}_0$ be a von Neumann algebra on a Hilbert space $\mathcal{H}$ with a faithful finite trace τ . We denote by $L^p(\tau)$ the Banach space completion of $\mathcal{M}_0$ with respect to, the norm

$$\|A\|_p \equiv \tau(|A|^p)^{1/p}, \quad A \in \mathcal{M}_0. \quad (5.16)$$

Then

$$\mathcal{M}_0 \equiv L^\infty(\tau) \subset L^p(\tau) \subset L^2(\tau) \subset L^q(\tau) \subset L^1(\tau), \quad 1 \leq q \leq 2 \leq p < \infty. \quad (5.17)$$

Let $2 \leq p < \infty$. Here we define a $*$ -representation π of $L^p(\tau)$ by

$$\pi(X)A = XA, \quad X \in L^p(\tau), \quad A \in L^\infty(\tau). \quad (5.18)$$

Then $\mathcal{M} \equiv \pi(L^p(\tau))$ is a partial EW*-algebra on $L^\infty(\tau)$ in $L^2(\tau)$ with $\mathcal{M}_b = \pi(L^\infty(\tau))$ which is integrable, that is, $\pi(X^\dagger) = \pi(X)^*$ for each $X \in L^p(\tau)$. Furthermore, $\pi(L^p(\tau))$ has a strongly cyclic and separating vector $\xi_0 \equiv \lambda_\tau(I)$, where I is an identity operator on $\mathcal{H}$. Let $\mathcal{N}_0$ be a von Neumann subalgebra of $\mathcal{M}_0$. We put

$$\mathcal{N} = \{\pi(X); X \in L^p(\tau), \pi(X)\lambda_\tau(I) \in L^p(\tau[\mathcal{N}_0])\}, \quad 2 \leq p \leq \infty. \quad (5.19)$$

Then $\mathcal{N}$ is an integrable partial EW*-subalgebra of $\mathcal{M}$ satisfying (N_2) and (N_3) and $P_{\mathcal{N}}\mathcal{M}\xi_0 = \mathcal{N}\xi_0$. By Theorem 5.2, there exists a conditional expectation of $(\mathcal{M}, \xi_0)$.

References

- [1] M. Takesaki, "Conditional expectations in von Neumann algebras," *Journal of Functional Analysis*, vol. 9, pp. 306–321, 1972.
- [2] S. P. Gudder and R. L. Hudson, "A noncommutative probability theory," *Transactions of the American Mathematical Society*, vol. 245, pp. 1–41, 1978.
- [3] A. Inoue, M. Takakura, and H. Ogi, "Unbounded conditional expectations for O^* -algebras," *Contemporary Mathematics*, vol. 427, pp. 225–234, 2007.
- [4] A. Inoue, H. Ogi, and M. Takakura, "Conditional expectations for unbounded operator algebras," *International Journal of Mathematics and Mathematical Sciences*, vol. 2007, Article ID 80152, 22 pages, 2007.
- [5] J.-P. Antoine, A. Inoue, and C. Trapani, *Partial $*$ -Algebras and Their Operator Realizations*, vol. 553 of *Mathematics and Its Applications*, Kluwer Academic Publishers, Dordrecht, The Netherlands, 2002.