Research Article

Convergence of Path and Approximation of Common Element of Null Spaces of Countably Infinite Family of m -Accretive Mappings in Uniformly Convex Banach Spaces

E. U. Ofoedu¹ and Y. Shehu²

¹ Department of Mathematics, Nnamdi Azikiwe University, Awka, Anambra State, Nigeria

² Department of Mathematics, University of Nigeria, Nsukka, Enugu, Nigeria

Correspondence should be addressed to E. U. Ofoedu, euofoedu@yahoo.com

Received 9 July 2009; Accepted 27 September 2009

Recommended by Gelu Popescu

We prove path convergence theorems and introduce a new iterative sequence for a countably infinite family of m -accretive mappings and prove strong convergence of the sequence to a common zero of these operators in uniformly convex real Banach space. Consequently, we obtain strong convergence theorems for a countably infinite family of pseudocontractive mappings. Our theorems extend and improve some important results which are announced recently by various authors.

Copyright © 2009 E. U. Ofoedu and Y. Shehu. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

Let E be a real Banach space with dual E^* . We denote by J the normalized duality mapping from E to 2^{E^*} defined by

$$Jx := \{f^* \in E^* : \langle x, f^* \rangle = \|x\|^2 = \|f^*\|^2\}, \quad (1.1)$$

where $\langle \cdot, \cdot \rangle$ denotes the generalized duality pairing between members of E and E^* . It is well known that if E^* is strictly convex, then J is single-valued (see, e.g., [1, 2]). In the sequel, we will denote the single-valued normalized duality mapping by j .

A mapping $A : D(A) \subseteq E \rightarrow E$ is called *accretive* if for all $x, y \in D(A)$ there exists $j(x - y) \in J(x - y)$ such that

$$\langle Ax - Ay, j(x - y) \rangle \geq 0. \quad (1.2)$$

By the result of Kato [3], (1.2) is equivalent to

$$\|x - y\| \leq \|x - y + s(Ax - Ay)\|, \quad \forall s > 0. \quad (1.3)$$

If E is a Hilbert space, accretive operators are also called *monotone*. An operator A is called *m-accretive* if it is accretive and $R(I + rA)$, range of $(I + rA)$, is E for all $r > 0$; and A is said to satisfy the range condition if $\text{cl}(D(A)) \subseteq R(I + rA)$, for all $r > 0$, where $\text{cl}(D(A))$ denotes the closure of the domain of A . It is easy to see that every *m-accretive* operator satisfies the range condition. An operator A is said to be *maximal accretive* if it is accretive and the inclusion $G(A) \subseteq G(B)$ implies $G(A) = G(B)$, where $G(A)$ denotes the graph of A and B is an accretive operator.

A mapping $T : D(T) \subseteq E \rightarrow R(T) \subseteq E$ is said to be *nonexpansive* if

$$\|Tx - Ty\| \leq \|x - y\|, \quad \forall x, y \in D(T). \quad (1.4)$$

It is not difficult to deduce from (1.3) that a mapping A is accretive if and only if its resolvent $J_r := (I + rA)^{-1}$, for all $r > 0$, is nonexpansive and single valued on the range of $(I + rA)$. Thus, in particular, $J_A = J_1 := (I + A)^{-1}$ is nonexpansive and single valued on the range of $(I + A)$. Furthermore, $F(J_A) := A^{-1}(0) := \{x \in D(A) : Ax = 0\}$. For more details see, for example, [4, 5].

Closely related to the class of accretive operators is the class of pseudocontractive maps. An operator T with domain $D(T)$ in E and range $R(T)$ in E is called *pseudocontractive* if $A := I - T$ is accretive. The importance of these operators in application is well known (see, e.g., [6–9] and the references contained therein).

It is well known that the class of pseudocontractive mappings properly contains the class of nonexpansive mappings (see, e.g., [4]). Construction of fixed points of nonexpansive mappings is an important subject in nonlinear operator theory and its applications, in particular, in image recovery and signal processing (see, e.g., [10]).

Iterative approximation of fixed points and zeros of nonlinear operators have been studied extensively by many authors to solve nonlinear operator equations as well as variational inequality problems (see, e.g., [11–15]). The iterative scheme

$$x_0 \in E, \quad x_{n+1} = J_{r_n} x_n, \quad n \geq 0, \quad (1.5)$$

(where J_{r_n} is the resolvent of an *m-accretive* operator A) for example, has been extensively studied over the past forty years or so for construction of zeros of accretive operators (see, e.g., [16–20]).

Kim and Xu [21] introduced a modification of Mann iterative scheme in a reflexive Banach space having weakly continuous duality mapping for finding a zero of an *m-accretive* operator A as follows:

$$x_0 = u \in E, \quad x_{n+1} = \alpha_n u + (1 - \alpha_n) J_{r_n} x_n, \quad n \geq 0. \quad (1.6)$$

They proved that the sequence $\{x_n\}_{n=0}^\infty$ generated by (1.6) converges to a zero of m -accretive operator A under the following conditions:

- (i) $\lim_{n \rightarrow \infty} \alpha_n = 0$,
- (ii) $\sum_{n=0}^\infty \alpha_n = \infty$ (equivalently, $\prod_{n=0}^\infty (1 - \alpha_n) = 0$),
- (iii) $\sum_{n=0}^\infty |\alpha_{n+1} - \alpha_n| < \infty$; $r_n \geq \epsilon$ for some $\epsilon > 0$ and for all $n \geq 0$,
- (iv) $\sum_{n=1}^\infty |1 - r_{n-1}/r_n| < \infty$,
- (v) $r_n \geq \epsilon$ for some $\epsilon > 0$ and for all $n \geq 0$ and $\sum_{n=1}^\infty |r_n - r_{n-1}| < \infty$.

In 2007, Qin and Su [22] also considered the following iterative scheme in either a uniformly smooth Banach space or a reflexive Banach space having a weakly sequentially continuous duality mapping:

$$\begin{aligned} x_0 &= u \in C, & y_n &= \beta_n x_n + (1 - \beta_n) J_{r_n} x_n, \\ x_{n+1} &= \alpha_n u + (1 - \alpha_n) y_n & n &\geq 0, \end{aligned} \quad (1.7)$$

where $\{\alpha_n\}_{n=0}^\infty$ and $\{\beta_n\}_{n=0}^\infty$ are sequences in $(0, 1)$. They proved that the sequence $\{x_n\}_{n=0}^\infty$ generated by (1.7) converges strongly to a zero of m -accretive operator A provided that $\{\alpha_n\}_{n=0}^\infty$ and $\{\beta_n\}_{n=0}^\infty$ satisfy conditions (i), (ii), and (iii), and $\{r_n\}_{n=0}^\infty$ satisfies condition (v).

Chen and Zhu [23] considered the following viscosity iterative scheme for resolvent J_{r_n} of m -accretive mapping A :

$$x_0 \in C, \quad x_{n+1} = \alpha_n f(x_n) + (1 - \alpha_n) J_{r_n} x_n, \quad n \geq 0, \quad (1.8)$$

where f is a contraction mapping defined on C . Under the assumption that $\{r_n\}_{n=0}^\infty$ satisfies condition (v), Chen and Zhu [23] proved in a reflexive Banach space having weakly sequentially continuous duality mapping that the sequence $\{x_n\}_{n=0}^\infty$ generated by (1.8) converges strongly to a zero of A , which solves a certain variational inequality.

Recently, Jung [24] introduced the following viscosity iterative method:

$$\begin{aligned} x_0 &\in C, & y_n &= \beta_n x_n + (1 - \beta_n) J_{r_n} x_n, \\ x_{n+1} &= \alpha_n f(x_n) + (1 - \alpha_n) y_n & n &\geq 0. \end{aligned} \quad (1.9)$$

Under certain appropriate conditions on the parameters $\{\alpha_n\}_{n=0}^\infty$, $\{\beta_n\}_{n=0}^\infty$, $\{r_n\}_{n=0}^\infty$ and the sequence $\{x_n\}_{n=0}^\infty$; Jung [24] established strong convergence of the sequence $\{x_n\}_{n=0}^\infty$ generated by (1.9) to a zero of A , which is a unique solution of a certain variational inequality problem, in either a reflexive Banach space having a weakly sequentially continuous duality mapping or a reflexive Banach space having a uniformly Gâteaux differentiable norm such that every weakly compact convex subset of E has the fixed point property for nonexpansive mappings.

In [5], Zegeye and Shahzad proved the following theorem.

Theorem ZS. *Let E be a strictly convex reflexive real Banach space which has uniformly Gâteaux differentiable norm and let K be a nonempty closed convex subset of E . Assume that every nonempty closed convex and bounded subset of E has the fixed point property for nonexpansive mappings.*

Let $A_i : K \rightarrow E, i = 1, 2, \dots, r$ be a finite family of m -accretive mappings with $\bigcap_{i=1}^r A_i^{-1}(0) \neq \emptyset$. For given $u, x_1 \in K$, let $\{x_n\}_{n \geq 1}$ be generated by the algorithm

$$x_{n+1} = \theta_n u + (1 - \theta_n) S_r x_n, \quad \forall n \geq 1, \quad (1.10)$$

where $S_r = a_0 I + a_1 J_{A_1} + \dots + a_r J_{A_r}$, with $J_{A_i} = (I + A_i)^{-1}, 0 < a_i < 1, i = 1, \dots, r, \sum_{i=1}^r a_i = 1$, and $\{\theta_n\}_{n \geq 1}$ is a sequence in $(0, 1)$ satisfying the following conditions:

- (i) $\lim_{n \rightarrow \infty} \theta_n = 0$;
- (ii) $\sum_{n=1}^{\infty} \theta_n = \infty$;
- (iii) $\sum_{n=1}^{\infty} |\theta_n - \theta_{n-1}| < \infty$ or $\lim_{n \rightarrow \infty} (|\theta_n - \theta_{n-1}| / \theta_n) = 0$.

Then, $\{x_n\}_{n \geq 1}$ converges strongly to a common solution of the equation $A_i x = 0$ for $i = 1, 2, \dots, r$.

Motivated by the results of the authors mentioned above, it is our purpose in this paper to prove new path convergence theorems and introduce a new iteration process for a countably infinite family of m -accretive mappings and prove strong convergence of the sequence to a common zero of these operators in uniformly convex real Banach spaces. As a result, we obtain strong convergence theorems for a countably infinite family of pseudocontractive mappings. Our theorems extend and improve some important results which are announced recently by various authors.

2. Preliminaries

Let E be a real normed linear space. Let $S := \{x \in E : \|x\| = 1\}$. E is said to have a *Gâteaux differentiable norm* (and E is called *smooth*) if the limit

$$\lim_{t \rightarrow 0} \frac{\|x + ty\| - \|x\|}{t} \quad (2.1)$$

exists for each $x, y \in S$; E is said to have a *uniformly Gâteaux differentiable norm* if for each $y \in S$ the limit is attained uniformly for $x \in S$. Furthermore, E is said to be *uniformly smooth* if the limit exists uniformly for $(x, y) \in S \times S$.

Let E be a real normed linear space. The modulus of convexity of E is the function $\delta_E : [0, 2] \rightarrow [0, 1]$ defined by

$$\delta_E(\epsilon) = \inf \left\{ 1 - \frac{\|x + y\|}{2} : \|x\| = \|y\| = 1, \epsilon = \|x - y\| \right\}. \quad (2.2)$$

The space E is said to be *uniformly convex* if and only if $\delta_E(\epsilon) > 0$ for all $\epsilon \in (0, 2]$. E is said to be *strictly convex* if for all $x, y \in E$ such that $\|x\| = \|y\| = 1$ and for all $\lambda \in (0, 1)$ we have $\|\lambda x + (1 - \lambda)y\| < 1$. It is well known that every uniformly convex Banach space is strictly convex.

A mapping T with domain $D(T)$ and range $R(T)$ in E is said to be *demiclosed* at p if whenever $\{x_n\}$ is a sequence in $D(T)$ such that $x_n \rightharpoonup x \in D(T)$ and $Tx_n \rightarrow p$, then $Tx = p$.

A mapping $T : D(T) \subseteq E \rightarrow E$ is said to be *demicompact* at h if for any bounded sequence $\{x_n\}$ in $D(T)$ such that $(x_n - Tx_n) \rightarrow h$ as $n \rightarrow \infty$, there exists a subsequence say $\{x_{n_j}\}$ of $\{x_n\}$ and $x^* \in D(T)$ such that $\{x_{n_j}\}$ converges strongly to x^* and $x^* - Tx^* = h$.

We need the following lemmas in the sequel.

Lemma 2.1. *Let E be a real normed space, then*

$$\|x + y\|^2 \leq \|x\|^2 + 2\langle y, j(x + y) \rangle, \quad (2.3)$$

for all $x, y \in E$ and for all $j(x + y) \in J(x + y)$.

Lemma 2.2 (Lemma 3 of Bruck [25]). *Let K be a nonempty closed convex subset of a strictly convex real Banach space E . Let $\{T_i\}_{i=1}^\infty$ be a sequence of nonexpansive mappings from K to E such that $\bigcap_{i=1}^\infty F(T_i) \neq \emptyset$. Let $\{\lambda_i\}_{i=1}^\infty$ be a sequence of positive numbers such that $\sum_{i=1}^\infty \lambda_i = 1$, then a mapping G on K defined by $Gx := \sum_{i=1}^\infty \lambda_i T_i x$ for all $x \in K$ is well defined, nonexpansive, and $F(G) = \bigcap_{i=1}^\infty F(T_i)$.*

Lemma 2.3 (Xu [26]). *Let $\{a_n\}$ be a sequence of nonnegative real numbers satisfying the following relation:*

$$a_{n+1} \leq (1 - \alpha_n)a_n + \alpha_n \sigma_n, \quad n \geq 1, \quad (2.4)$$

where $\{\alpha_n\}_{n=1}^\infty \subset [0, 1]$ and $\{\sigma_n\}_{n=1}^\infty$ is a sequence in $\mathbb{R}$ satisfying (i) $\sum \alpha_n = \infty$; (ii) $\limsup \sigma_n \leq 0$. Then, $a_n \rightarrow 0$ as $n \rightarrow \infty$.

Lemma 2.4 (Suzuki [27]). *Let $\{x_n\}$ and $\{y_n\}$ be bounded sequences in a Banach space X and let $\{\beta_n\}$ be a sequence in $[0, 1]$ with $0 < \liminf_{n \rightarrow \infty} \beta_n \leq \limsup_{n \rightarrow \infty} \beta_n < 1$. Suppose that $x_{n+1} = (1 - \beta_n)y_n + \beta_n x_n$ for all integers $n \geq 1$ and $\limsup_{n \rightarrow \infty} (\|y_{n+1} - y_n\| - \|x_{n+1} - x_n\|) \leq 0$. Then, $\lim_{n \rightarrow \infty} \|y_n - x_n\| = 0$.*

Lemma 2.5 (Cioranescu [28]). *Let A be a continuous accretive operator defined on a real Banach space E with $D(A) = E$. Then, A is m -accretive.*

Lemma 2.6 (C. E. Chidume and C. O. Chidume [29]). *Let K be a nonempty closed convex subset of a real Banach space E . For arbitrary $r > 0$, let $B_r(0) := \{x \in E : \|x\| \leq r\}$. Then, there exists a continuous strictly increasing function $g : [0, \infty) \rightarrow [0, \infty)$, $g(0) = 0$ such that for every $x, y \in B_r(0)$ and for $p \in (1, \infty)$, the following inequality holds:*

$$4 \cdot 2^p g\left(\frac{1}{2}\|x + y\|\right) \leq (p \cdot 2^p - 4)\|x\|^p + p \cdot 2^p \langle y, j_p(x) \rangle + 4\|y\|^p. \quad (2.5)$$

3. Path Convergence Theorems

We begin with the following lemma.

Lemma 3.1. *Let K be a nonempty closed convex subset of a reflexive strictly convex Banach space E . Let $T : K \rightarrow K$ be a nonexpansive mapping. Let $\{x_n\}_{n=1}^\infty$, a bounded sequence in K , be an approximate fixed point sequence of T , that is $\lim_{n \rightarrow \infty} \|x_n - Tx_n\| = 0$. Let $\varphi(x) = \mu_n \|x_n - x\|^2$, for all $x \in K$ and*

let $\Gamma = \{x \in K \cap B : \varphi(x) = \min_{z \in K} \varphi(z)\}$, where B is any bounded closed convex nonempty subset of E such that $x_n \in B$ for all $n \in \mathbb{N}$. Then T has a fixed point in Γ , provided that $F(T) \neq \emptyset$.

Proof. Since E is a reflexive Banach space, then Γ is a bounded closed convex nonempty subset of E . Since $\lim_{n \rightarrow \infty} \|x_n - Tx_n\| = 0$, we have that for all $x \in \Gamma$,

$$\begin{aligned} \varphi(Tx) &= \mu_n \|x_n - Tx\|^2 \leq \mu_n (\|x_n - Tx_n\| + \|Tx_n - Tx\|)^2 \\ &\leq \mu_n \|x_n - x\|^2 = \varphi(x). \end{aligned} \quad (3.1)$$

Hence, $Tx \in \Gamma$, for all $x \in \Gamma$, that is, Γ is invariant under T . Let $x^* \in F(T)$. Then since every closed convex nonempty subset of a reflexive strictly convex Banach space is a Chebyshev set (see, e.g., [30, Corollary 5.1.19]), there exists a unique $u^* \in \Gamma$ such that

$$\|x^* - u^*\| = \inf_{z \in \Gamma} \|x^* - z\|, \quad (3.2)$$

but $x^* = Tx^*$ and $Tu^* \in \Gamma$. Thus,

$$\|x^* - Tu^*\| = \|Tx^* - Tu^*\| \leq \|x^* - u^*\|. \quad (3.3)$$

So, $Tu^* = u^*$. Hence, $F(T) \cap \Gamma \neq \emptyset$. This completes the proof. $\square$

Proposition 3.2. Let K be a nonempty closed convex subset of a real Banach space E . Let $A_i : K \rightarrow E$, $i = 1, 2, \dots$, be a countably infinite family of m -accretive mappings and define $J_{A_i} := (I + A_i)^{-1}$, $i = 1, 2, \dots$. Let $\{\alpha_n\}_{n=1}^\infty$, $\{\sigma_{i,n}\}_{n=1}^\infty$, $i = 1, 2, \dots$ be sequences in $(0, 1)$ such that $\sum_{i=1}^\infty \sigma_{i,n} = (1 - \alpha_n)$. Fix $\delta \in [\gamma_1, \gamma_2]$, for some $\gamma_1, \gamma_2 \in (0, 1)$. For arbitrary fixed $u \in K$, define a map $T_n : K \rightarrow K$ by

$$T_n x = \alpha_n u + \sum_{i=1}^\infty \sigma_{i,n} ((1 - \delta)x + \delta J_{A_i} x), \quad \forall x \in K. \quad (3.4)$$

Then, T_n is a strict contraction on K .

Proof. Let $x, y \in K$, then

$$\begin{aligned} \|T_n x - T_n y\| &= \left\| \sum_{i=1}^\infty \sigma_{i,n} ((1 - \delta)(x - y) + \delta(J_{A_i} x - J_{A_i} y)) \right\| \\ &\leq \sum_{i=1}^\infty \sigma_{i,n} ((1 - \delta)\|x - y\| + \delta\|J_{A_i} x - J_{A_i} y\|) \\ &\leq (1 - \alpha_n)\|x - y\|. \end{aligned} \quad (3.5)$$

$\square$

Thus, for each $n \in \mathbb{N}$, there is a unique $z_n \in K$ satisfying

$$z_n = \alpha_n u + \sum_{i=1}^\infty \sigma_{i,n} ((1 - \delta)z_n + \delta J_{A_i} z_n). \quad (3.6)$$

Lemma 3.3. Let K be a nonempty closed convex subset of a real Banach space E . For each $i \geq 1$, let $A_i : K \rightarrow E$ be a countably infinite family of m -accretive mappings. For $n \in \mathbb{N}$, let $\{z_n\}$ be a sequence satisfying (3.6) and assume $\bigcap_{i=1}^{\infty} A_i^{-1}(0) \neq \emptyset$. Then, $\{z_n\}$ is bounded.

Proof. Let $x^* \in \bigcap_{i=1}^{\infty} A_i^{-1}(0) = \bigcap_{i=1}^{\infty} F(J_{A_i})$. Then, using (3.6), we obtain

$$\begin{aligned} \|z_n - x^*\|^2 &= \left\langle \alpha_n(u - x^*) + \sum_{i=1}^{\infty} \sigma_{i,n}((1 - \delta)z_n + \delta J_{A_i} z_n - x^*), j(z_n - x^*) \right\rangle \\ &\leq \alpha_n \langle u - x^*, j(z_n - x^*) \rangle + \sum_{i=1}^{\infty} \sigma_{i,n} \|z_n - x^*\|^2 \\ &= \alpha_n \langle u - x^*, j(z_n - x^*) \rangle + (1 - \alpha_n) \|z_n - x^*\|^2, \end{aligned} \quad (3.7)$$

which implies that $\|z_n - x^*\| \leq \|u - x^*\|$. Thus, $\{z_n\}$ is bounded. $\square$

Lemma 3.4. Let K be a nonempty closed convex subset of a uniformly convex real Banach space E . For each $i \geq 1$, let $A_i : K \rightarrow E$ be a countably infinite family of m -accretive mappings such that $\bigcap_{i=1}^{\infty} A_i^{-1}(0) \neq \emptyset$. Let $\{\alpha_n\}$ be a sequence in $(0, 1)$ such that $\lim_{n \rightarrow \infty} (\alpha_n / \sigma_{i,n}) = 0$, for all $i \geq 1$, $\sum_{i=1}^{\infty} \sigma_{i,n} = (1 - \alpha_n)$. Let $\{z_n\}$ be a sequence satisfying (3.6). Then, $\lim_{n \rightarrow \infty} \|z_n - J_{A_i} z_n\| = 0$, for all $i \geq 1$. Furthermore, if $\{\lambda_i\}_{i=1}^{\infty}$ is a sequence in $(0, 1)$ such that $\sum_{i=1}^{\infty} \lambda_i = 1$; $\lim_{n \rightarrow \infty} \sum_{i=1}^{\infty} |\sigma_{i,n} - \lambda_i| = 0$ and define $G := (1 - \delta)I + \delta T$, where $T := \sum_{i=1}^{\infty} \lambda_i J_{A_i}$, then $\lim_{n \rightarrow \infty} \|z_n - G z_n\| = 0$.

Proof. We start by showing that $\lim_{n \rightarrow \infty} \|z_n - J_{A_i} z_n\| = 0$, for all $i \geq 1$. For this, let $S_i := (1 - \delta)I + \delta J_{A_i}$, where I is the identity operator on K . Since $\{z_n\}_{n=1}^{\infty}$ is bounded, then for each $i \geq 1$ and $x^* \in \bigcap_{i=1}^{\infty} A_i^{-1}(0)$, we have the following using (2.5):

$$\begin{aligned} 4.2^p g\left(\frac{1}{2} \|S_i z_n - z_n\|\right) &= 4.2^p g\left(\frac{1}{2} \|S_i z_n - x^* + x^* - z_n\|\right) \\ &\leq (p.2^p - 4) \|x^* - z_n\|^p + p.2^p \langle S_i z_n - x^*, j_p(x^* - z_n) \rangle + 4 \|S_i z_n - x^*\|^p \\ &\leq (p.2^p - 4) \|x^* - z_n\|^p + p.2^p \langle S_i z_n - z_n + z_n - x^*, j_p(x^* - z_n) \rangle \\ &\quad + 4 \|S_i z_n - x^*\|^p \\ &= (p.2^p - 4) \|x^* - z_n\|^p + p.2^p \langle S_i z_n - z_n, j_p(x^* - z_n) \rangle \\ &\quad - p.2^p \langle x^* - z_n, j_p(x^* - z_n) \rangle + 4 \|S_i z_n - x^*\|^p \\ &\leq p.2^p \langle z_n - S_i z_n, j_p(z_n - x^*) \rangle. \end{aligned} \quad (3.8)$$

Hence,

$$\frac{4}{p} g\left(\frac{1}{2} \|S_i z_n - z_n\|\right) \leq \langle z_n - S_i z_n, j_p(z_n - x^*) \rangle, \quad (3.9)$$

and so,

$$\frac{4}{p} \sum_{i=1}^{\infty} \sigma_{i,n} g\left(\frac{1}{2} \|S_i z_n - z_n\|\right) \leq \frac{4}{p} \sum_{i=1}^{\infty} \sigma_{i,n} \langle z_n - S_i z_n, j_p(z_n - x^*) \rangle. \quad (3.10)$$

Using (3.6), we have

$$\begin{aligned} \langle z_n - x^*, j_p(z_n - x^*) \rangle &= \alpha_n \langle u - x^*, j_p(z_n - x^*) \rangle \\ &\quad + \sum_{i=1}^{\infty} \sigma_{i,n} \langle S_i z_n - z_n + z_n - x^*, j_p(z_n - x^*) \rangle \\ &= \alpha_n \langle u - x^*, j_p(z_n - x^*) \rangle + \sum_{i=1}^{\infty} \sigma_{i,n} \langle S_i z_n - z_n, j_p(z_n - x^*) \rangle \\ &\quad + (1 - \alpha_n) \langle z_n - x^*, j_p(z_n - x^*) \rangle, \end{aligned} \quad (3.11)$$

which implies

$$\sum_{i=1}^{\infty} \sigma_{i,n} \langle z_n - S_i z_n, j_p(z_n - x^*) \rangle = \alpha_n \langle u - z_n, j_p(z_n - x^*) \rangle. \quad (3.12)$$

Using this and (3.10), we get

$$\frac{4}{p} \sum_{i=1}^{\infty} \sigma_{i,n} g\left(\frac{1}{2} \|S_i z_n - z_n\|\right) \leq \alpha_n \langle u - z_n, j_p(z_n - x^*) \rangle. \quad (3.13)$$

Since $\{z_n\}$ is bounded, we have that

$$\sum_{i=1}^{\infty} \sigma_{i,n} g\left(\frac{1}{2} \|S_i z_n - z_n\|\right) \leq \alpha_n M, \quad (3.14)$$

for some constant $M > 0$. This yields

$$g\left(\frac{1}{2} \|S_i z_n - z_n\|\right) \leq \frac{\alpha_n}{\sigma_{i,n}} M. \quad (3.15)$$

Thus, since g is continuous, strictly increasing, $g(0) = 0$, and $\lim_{n \rightarrow \infty} (\alpha_n / \sigma_{i,n}) = 0$, for all $i \geq 1$, we have

$$2g\left(\frac{1}{2} \lim_{n \rightarrow \infty} \|S_i z_n - z_n\|\right) = 0. \quad (3.16)$$

So, $\lim_{n \rightarrow \infty} \|S_i z_n - z_n\| = 0$, for all $i \geq 1$, but

$$\begin{aligned} \|S_i z_n - z_n\| &= \|(1 - \delta)z_n + \delta J_{A_i} z_n - z_n\| \\ &= \|\delta(J_{A_i} z_n - z_n)\| \\ &= \delta \|J_{A_i} z_n - z_n\|. \end{aligned} \quad (3.17)$$

Thus,

$$\lim_{n \rightarrow \infty} \|J_{A_i} z_n - z_n\| = 0, \quad \forall i \geq 1. \quad (3.18)$$

Next, we show that $\lim_{n \rightarrow \infty} \|z_n - Gz_n\| = 0$. Observe that

$$z_n - Gz_n = \alpha_n u + \sum_{i=1}^{\infty} (\sigma_{i,n} - \lambda_i) [(1 - \delta)z_n + \delta J_{A_i} z_n]. \quad (3.19)$$

So,

$$\|z_n - Gz_n\| \leq \alpha_n \|u\| + M \sum_{i=1}^{\infty} |\sigma_{i,n} - \lambda_i| \quad (3.20)$$

for some $M > 0$. Hence,

$$\lim_{n \rightarrow \infty} \|z_n - Gz_n\| = 0. \quad (3.21)$$

This completes the proof. $\square$

Theorem 3.5. Let K be a nonempty closed convex subset of a uniformly convex real Banach space E with uniformly Gâteaux differentiable norm. Let $A_i : K \rightarrow E$, $i = 1, 2, \dots$, be a countably infinite family of m -accretive mappings such that $\bigcap_{i=1}^{\infty} A_i^{-1}(0) \neq \emptyset$. Let $\{z_n\}$ be a sequence satisfying (3.6). Let $\{\lambda_i\}_{i=1}^{\infty}$ be a sequence in $(0, 1)$ such that $\sum_{i=1}^{\infty} \lambda_i = 1$ and $\lim_{n \rightarrow \infty} \sum_{i=1}^{\infty} |\sigma_{i,n} - \lambda_i| = 0$. Let $G := (1 - \delta)I + \delta T$, where $T := \sum_{i=1}^{\infty} \lambda_i J_{A_i}$. Then, $\{z_n\}$ converges strongly to an element in $\bigcap_{i=1}^{\infty} A_i^{-1}(0)$.

Proof. Observe that by Lemma 2.2, $T := \sum_{i=1}^{\infty} \lambda_i J_{A_i}$ is well defined, nonexpansive, and $F(T) = \bigcap_{i=1}^{\infty} F(J_{A_i}) = \bigcap_{i=1}^{\infty} A_i^{-1}(0)$. Furthermore, it is easy to see that G is nonexpansive and that $F(G) = F(T) = \bigcap_{i=1}^{\infty} A_i^{-1}(0)$. Now, since $\{z_n\}$ is bounded and $\lim_{n \rightarrow \infty} \|Gz_n - z_n\| = 0$, we have by Lemma 3.1 that there exists a unique z^* in the set $\Omega^* := \{x \in K \cap B^* : \mu_n \|z_n - x\|^2 = \min_{y \in K} \|z_n - y\|^2\}$ such that $Gz^* = z^*$, where B^* is a bounded closed convex nonempty subset of E such that $u, z_n \in B^*$ for all $n \in \mathbb{N}$. Thus, $z^* \in F(G) = \bigcap_{i=1}^{\infty} A_i^{-1}(0)$. Let $t \in (0, 1)$, then by convexity of $K \cap B^*$, we have that $(1 - t)z^* + tu \in K \cap B^*$. Thus, $\mu_n \|z_n - z^*\|^2 \leq \mu_n \|z_n - ((1 - t)z^* + tu)\|^2 = \mu_n \|z_n - z^* - t(u - z^*)\|^2$. Moreover, we have, by Lemma 2.1 that

$$\|z_n - z^* - t(u - z^*)\|^2 \leq \|z_n - z^*\|^2 - 2t \langle u - z^*, j(z_n - z^* - t(u - z^*)) \rangle. \quad (3.22)$$

This implies that $\mu_n \langle u - z^*, j(z_n - z^* - t(u - z^*)) \rangle \leq 0$. Furthermore, since E has uniformly Gâteaux differentiable norm, we obtain that

$$\lim_{t \rightarrow 0} (\langle u - z^*, j(z_n - z^*) \rangle - \langle u - z^*, j(z_n - z^* - t(u - z^*)) \rangle) = 0. \quad (3.23)$$

Thus, given $\epsilon > 0$, there exists $\delta_\epsilon > 0$ such that for all $t \in (0, \delta_\epsilon)$ and for all $n \in \mathbb{N}$,

$$\langle u - z^*, j(z_n - z^*) \rangle < \epsilon + \langle u - z^*, j(z_n - z^* - t(u - z^*)) \rangle. \quad (3.24)$$

Taking Banach limit on both sides of this inequality, we obtain

$$\mu_n \langle u - z^*, j(z_n - z^*) \rangle \leq \epsilon; \quad (3.25)$$

and since $\epsilon > 0$ is arbitrary, we have that

$$\mu_n \langle u - z^*, j(z_n - z^*) \rangle \leq 0. \quad (3.26)$$

Now, using (3.6), we have that

$$\begin{aligned} \|z_n - z^*\|^2 &= \left\langle \alpha_n(u - z^*) + \sum_{i=1}^{\infty} \sigma_{i,n}(((1 - \delta)z_n + \delta J_{A_i} z_n) - z^*), j(z_n - z^*) \right\rangle \\ &\leq \alpha_n \langle u - z^*, j(z_n - z^*) \rangle + (1 - \alpha_n) \|z_n - z^*\|^2. \end{aligned} \quad (3.27)$$

So,

$$\|z_n - z^*\|^2 \leq \langle u - z^*, j(z_n - z^*) \rangle. \quad (3.28)$$

Again, taking Banach limit, we obtain

$$\mu_n \|z_n - z^*\|^2 \leq \mu_n \langle u - z^*, j(z_n - z^*) \rangle \leq 0, \quad (3.29)$$

so that $\mu_n \|z_n - z^*\|^2 = 0$. Hence, there exists a subsequence $\{z_{n_l}\}_{l=1}^{\infty}$ of $\{z_n\}_{n=1}^{\infty}$ such that $\lim_{l \rightarrow \infty} z_{n_l} = z^*$. We now show that $\{z_n\}_{n=1}^{\infty}$ actually converges to z^* . Suppose there is another subsequence $\{z_{n_k}\}_{k=1}^{\infty}$ of $\{z_n\}_{n=1}^{\infty}$ such that $\lim_{k \rightarrow \infty} z_{n_k} = u^*$. Then, since $\lim_{n \rightarrow \infty} \|J_{A_i} z_n - z_n\| = 0$ and J_{A_i} is continuous for all $i \in \mathbb{N}$, we have that $u^* \in \bigcap_{i=1}^{\infty} A_i^{-1}(0)$.

Claim 1 ($u^* = z^*$). Suppose for contradiction that $u^* \neq z^*$, then $\|u^* - z^*\| > 0$, but using (3.6), we have that

$$\begin{aligned}
 \|z_{n_l} - u^*\|^2 &= \left\langle \alpha_{n_l}(u - u^*) + \sum_{i=1}^{\infty} \sigma_{i,n_l}(((1-\delta)z_{n_l} + \delta J_{A_i} z_{n_l}) - u^*), j(z_{n_l} - u^*) \right\rangle \\
 &= \alpha_{n_l} \langle u - z^*, j(z_{n_l} - u^*) \rangle + \alpha_{n_l} \langle z^* - z_{n_l}, j(z_{n_l} - u^*) \rangle \\
 &\quad + \alpha_{n_l} \|z_{n_l} - u^*\|^2 + \sum_{i=1}^{\infty} \sigma_{i,n_l} \langle (1-\delta)z_{n_l} + \delta J_{A_i} z_{n_l} - u^*, j(z_{n_l} - u^*) \rangle \\
 &\leq \alpha_{n_l} \langle u - z^*, j(z_{n_l} - u^*) \rangle + \alpha_{n_l} \langle z^* - z_{n_l}, j(z_{n_l} - u^*) \rangle \\
 &\quad + \alpha_{n_l} \|z_{n_l} - u^*\|^2 + (1-\delta)(1-\alpha_{n_l}) \|z_{n_l} - u^*\|^2 + \delta(1-\alpha_{n_l}) \|z_{n_l} - u^*\|^2 \\
 &= \alpha_{n_l} \langle u - z^*, j(z_{n_l} - u^*) \rangle + \alpha_{n_l} \langle z^* - z_{n_l}, j(z_{n_l} - u^*) \rangle + \|z_{n_l} - u^*\|^2.
 \end{aligned} \tag{3.30}$$

Thus,

$$\langle u - z^*, j(u^* - z_{n_l}) \rangle \leq \|z_{n_l} - u^*\| \|z_{n_l} - z^*\|. \tag{3.31}$$

Using the fact that $\{z_n\}_{n=1}^{\infty}$ is bounded and that E has a uniformly Gâteaux differentiable norm, we obtain from (3.31) that

$$\langle u - z^*, j(u^* - z^*) \rangle \leq 0. \tag{3.32}$$

Similarly, we also obtain that $\langle u - u^*, j(z^* - u^*) \rangle \leq 0$ or rather

$$\langle u^* - u, j(u^* - z^*) \rangle \leq 0. \tag{3.33}$$

Adding (3.32) and (3.33), we have that $\|z^* - u^*\| \leq 0$, a contradiction. Thus, $z^* = u^*$. Hence, $\{z_n\}_{n=1}^{\infty}$ converges strongly to $z^* \in \bigcap_{i=1}^{\infty} A_i^{-1}(0)$. This completes the proof. $\square$

Theorem 3.6. Let K be a nonempty closed convex subset of a uniformly convex real Banach space E with uniformly Gâteaux differentiable norm. Let $A_i : K \rightarrow E$, $i = 1, 2, \dots$, be a countably infinite family of m -accretive mappings such that $\bigcap_{i=1}^{\infty} A_i^{-1}(0) \neq \emptyset$. Let $\{z_n\}$ be a sequence satisfying (3.6). If at least one of the mappings J_{A_i} is demicompact, then $\{z_n\}$ converges strongly to an element of $\bigcap_{i=1}^{\infty} A_i^{-1}(0)$.

Proof. For fixed $s \in \mathbb{N}$, let J_{A_s} be demicompact. Since $\lim_{n \rightarrow \infty} \|z_n - J_{A_s} z_n\| = 0$, there exists a subsequence say $\{z_{n_k}\}$ of $\{z_n\}$ that converges strongly to some point $z^* \in K$. By continuity of J_{A_i} and the fact that $\lim_{k \rightarrow \infty} \|z_{n_k} - J_{A_i} z_{n_k}\| = 0$, $i = 1, 2, \dots$, we have that $z^* \in \bigcap_{i=1}^{\infty} A_i^{-1}(0)$. Assuming that there is another subsequence $\{z_{n_j}\}_{j=1}^{\infty}$ of $\{z_n\}$ that converges strongly to a point say, q^* , then following the argument of the last part of proof of Theorem 3.5, we obtain that $\{z_n\}_{n=1}^{\infty}$ converges strongly to $z^* \in \bigcap_{i=1}^{\infty} A_i^{-1}(0)$. This completes the proof. $\square$

Theorem 3.7. Let K be a nonempty closed convex subset of a uniformly convex real Banach space E with uniformly Gâteaux differentiable norm and also admits weakly sequential continuous duality

mapping. Let $A_i : K \rightarrow E, i = 1, 2, \dots$, be a countably infinite family of m -accretive mappings such that $\bigcap_{i=1}^{\infty} A_i^{-1}(0) \neq \emptyset$. Let $\{z_n\}$ be a sequence satisfying (3.6). Then, $\{z_n\}$ converges strongly to an element of $\bigcap_{i=1}^{\infty} A_i^{-1}(0)$.

Proof. Since $\{z_n\}$ is bounded, there exists a subsequence say $\{z_{n_k}\}$ of $\{z_n\}$ that converges weakly to some point $z^* \in K$. Using the demiclosedness property of $(I - J_{A_i})$ at 0 for each $i \geq 1$ (since J_{A_i} is nonexpansive for each $i \in \mathbb{N}$, see, e.g., [31]) and the fact that $\lim_{k \rightarrow \infty} \|z_{n_k} - J_{A_i} z_{n_k}\| = 0$, we get that $z^* \in \bigcap_{i=1}^{\infty} A_i^{-1}(0)$. We also observe from (3.6) that

$$\begin{aligned} \|z_{n_k} - z^*\|^2 &= \left\langle \alpha_{n_k} u + \sum_{i=1}^{\infty} \sigma_{i,n_k} ((1 - \delta) z_{n_k} + \delta J_{A_i} z_{n_k}) - z^*, j(z_{n_k} - z^*) \right\rangle \\ &\leq \alpha_{n_k} \langle u - z^*, j(z_{n_k} - z^*) \rangle + \sum_{i=1}^{\infty} \sigma_{i,n_k} \|z_{n_k} - z^*\|^2 \\ &= \alpha_{n_k} \langle u - z^*, j(z_{n_k} - z^*) \rangle + (1 - \alpha_{n_k}) \|z_{n_k} - z^*\|^2. \end{aligned} \quad (3.34)$$

This implies that $\|z_{n_k} - z^*\|^2 \leq \langle u - z^*, j(z_{n_k} - z^*) \rangle$. Using the fact that j is weakly sequential continuous, then we have from the last inequality that $\{z_{n_k}\}$ converges strongly to z^* . Then following the argument of the last part of proof of Theorem 3.5, we obtain that $\{z_n\}_{n=1}^{\infty}$ converges strongly to $z^* \in \bigcap_{i=1}^{\infty} A_i^{-1}(0)$. This completes the proof. $\square$

4. Convergence Theorems for Countably Infinite Family of m -Accretive Mappings

For the rest of this paper, $\{\alpha_n\}_{n=1}^{\infty}$ and $\{\sigma_{i,n}\}_{n=1}^{\infty}$ are in $(0, 1)$ satisfying the following additional conditions:

- (i) $\lim_{n \rightarrow \infty} \alpha_n = 0$,
- (ii) $\sum_{n=1}^{\infty} \alpha_n = \infty$,
- (iii) $\lim_{n \rightarrow \infty} \sum_{i=1}^{\infty} |\sigma_{i,n+1} - \sigma_{i,n}| = 0$.

Theorem 4.1. Let K be a nonempty closed convex subset of a uniformly convex real Banach space E with uniformly Gâteaux differentiable norm. Let $A_i : K \rightarrow E, i = 1, 2, \dots$, be a countably infinite family of m -accretive mappings such that $\bigcap_{i=1}^{\infty} A_i^{-1}(0) \neq \emptyset$. For fixed $\delta \in [\gamma_1, \gamma_2]$, for some $\gamma_1, \gamma_2 \in (0, 1)$, $u \in K$, let $\{x_n\}_{n=1}^{\infty}$ be generated by

$$x_1 \in K, \quad x_{n+1} = \alpha_n u + \sum_{i=1}^{\infty} \sigma_{i,n} ((1 - \delta) x_n + \delta J_{A_i} x_n), \quad n \geq 1, \quad (4.1)$$

then $\lim_{n \rightarrow \infty} \sum_{i=1}^{\infty} \sigma_{i,n} ((1 - \delta) x_n + \delta J_{A_i} x_n - x_n) = 0$. Furthermore, if $\{\alpha_n\}_{n \geq 1}$ is such that

$$\lim_{n \rightarrow \infty} \frac{\sum_{i=1}^{\infty} \sigma_{i,n} ((1 - \delta) x_n + \delta J_{A_i} x_n - x_n)}{\alpha_n} = 0, \quad (4.2)$$

then $\{x_n\}_{n=1}^{\infty}$ converges strongly to a common zero of $\{A_i\}_{i=1}^{\infty}$.

Proof. Using mathematical induction, it is easy to see that for $x^* \in \bigcap_{i=1}^{\infty} A_i^{-1}(0)$ fixed

$$\|x_n - x^*\| \leq \max\{\|u - x^*\|, \|x_1 - x^*\|\}, \quad \forall n \geq 1. \quad (4.3)$$

Hence, $\{x_n\}_{n=1}^{\infty}$ is bounded and so $\{J_{A_i}x_n\}_{n=1}^{\infty}$ is also bounded.

Now, define the sequences $\{\beta_n\}_{n=1}^{\infty}$ and $\{y_n\}_{n=1}^{\infty}$ by $\beta_n := (1 - \delta)\alpha_n + \delta$ and $y_n := (x_{n+1} - x_n + \beta_n x_n) / \beta_n$. Then,

$$y_n = \frac{\alpha_n u + \delta \sum_{i \geq 1} \sigma_{i,n} ((1 - \delta)x_n + \delta J_{A_i} x_n)}{\beta_n}. \quad (4.4)$$

Observe that $\{y_n\}_{n=1}^{\infty}$ is bounded and that

$$\begin{aligned} \|y_{n+1} - y_n\| - \|x_{n+1} - x_n\| &\leq \left| \frac{\alpha_{n+1}}{1 - \beta_{n+1}} - \frac{\alpha_n}{1 - \beta_n} \right| \|u\| \\ &\quad + \left| \frac{\delta(1 - \alpha_{n+1})}{\beta_{n+1}} - 1 \right| \|x_{n+1} - x_n\| \\ &\quad + \frac{\delta M}{\beta_{n+1}\beta_n} \sum_{i=1}^{\infty} |\sigma_{i,n+1} - \sigma_{i,n}| + \frac{\delta M}{\beta_{n+1}\beta_n} |\beta_{n+1} - \beta_n|, \end{aligned} \quad (4.5)$$

for some $M > 0$. Thus,

$$\limsup_{n \rightarrow \infty} (\|y_{n+1} - y_n\| - \|x_{n+1} - x_n\|) \leq 0. \quad (4.6)$$

Hence, by Lemma 2.4, we have $\lim_{n \rightarrow \infty} \|y_n - x_n\| = 0$. Consequently, we have

$$\lim_{n \rightarrow \infty} \|x_{n+1} - x_n\| = \lim_{n \rightarrow \infty} \beta_n \|y_n - x_n\| = 0. \quad (4.7)$$

From (4.1), we have that

$$x_{n+1} - x_n = \alpha_n(u - x_n) + \sum_{i=1}^{\infty} \sigma_{i,n} (((1 - \delta)x_n + \delta J_{A_i} x_n) - x_n), \quad (4.8)$$

which implies that $\|\sum_{i=1}^{\infty} \sigma_{i,n} ((1 - \delta)x_n + \delta J_{A_i} x_n - x_n)\| \leq \|x_{n+1} - x_n\| + \alpha_n \|u - x_n\|$ and thus

$$\lim_{n \rightarrow \infty} \left\| \sum_{i=1}^{\infty} \sigma_{i,n} (((1 - \delta)x_n + \delta J_{A_i} x_n) - x_n) \right\| = 0. \quad (4.9)$$

Let $\{z_n\}_{n=1}^\infty$ be a sequence satisfying (3.6). Then, by Theorem 3.5, $z_n \rightarrow z^* \in \bigcap_{i=1}^\infty A_i^{-1}(0)$. Using Lemma 2.1, we have that

$$\begin{aligned} \|z_n - x_n\|^2 &\leq \left\| \sum_{i=1}^\infty \sigma_{i,n} ((1-\delta)z_n + \delta J_{A_i} z_n) - ((1-\delta)x_n + \delta J_{A_i} x_n) + ((1-\delta)x_n + \delta J_{A_i} x_n) - x_n \right\|^2 \\ &\quad + 2\alpha_n \langle u - x_n, j(z_n - x_n) \rangle \\ &\leq \left((1-\alpha_n) \|z_n - x_n\| + \left\| \sum_{i=1}^\infty \sigma_{i,n} ((1-\delta)x_n + \delta J_{A_i} x_n) - x_n \right\| \right)^2 \\ &\quad + 2\alpha_n \langle u - z_n, j(z_n - x_n) \rangle. \end{aligned} \quad (4.10)$$

This implies that

$$\begin{aligned} &\langle u - z_n, j(x_n - z_n) \rangle \\ &\leq \frac{\alpha_n}{2} \|z_n - x_n\|^2 + \frac{(1-\alpha_n) \|z_n - x_n\| \cdot \left\| \sum_{i=1}^\infty \sigma_{i,n} ((1-\delta)x_n + \delta J_{A_i} x_n) - x_n \right\|}{\alpha_n} \\ &\quad + \frac{\left\| \sum_{i=1}^\infty \sigma_{i,n} ((1-\delta)x_n + \delta J_{A_i} x_n) - x_n \right\|^2}{2\alpha_n} \end{aligned} \quad (4.11)$$

and hence,

$$\limsup_{n \rightarrow \infty} \langle u - z_n, j(x_n - z_n) \rangle \leq 0. \quad (4.12)$$

Moreover, we have that

$$\begin{aligned} \langle u - z_n, j(x_n - z_n) \rangle &= \langle u - z^*, j(x_n - z^*) \rangle + \langle u - z^*, j(x_n - z_n) - j(x_n - z^*) \rangle \\ &\quad + \langle z^* - z_n, j(x_n - z_n) \rangle. \end{aligned} \quad (4.13)$$

Using the boundedness of $\{x_n\}_{n=1}^\infty$, we have

$$\langle z^* - z_n, j(x_n - z_n) \rangle \rightarrow 0, \quad \text{as } n \rightarrow \infty. \quad (4.14)$$

Also, since j is norm-to-weak* uniformly continuous on bounded subsets of E , we have that

$$\langle u - z^*, j(x_n - z_n) - j(x_n - z^*) \rangle \rightarrow 0, \quad \text{as } n \rightarrow \infty. \quad (4.15)$$

From (4.12) and (4.13), we obtain that

$$\limsup_{n \rightarrow \infty} \langle u - z^*, j(x_n - z^*) \rangle \leq 0. \quad (4.16)$$

Finally, using Lemma 2.1 on (4.1), we get

$$\begin{aligned} \|x_{n+1} - z^*\|^2 &\leq \left\| \sum_{i=1}^{\infty} \sigma_{i,n+1} ((1-\delta)x_n + \delta J_{A_i} x_n) - z^* \right\|^2 \\ &\quad + 2\alpha_n \langle u - z^*, j(x_{n+1} - z^*) \rangle \\ &\leq (1 - \alpha_n) \|x_n - z^*\|^2 + 2\alpha_n \langle u - z, j(x_{n+1} - z^*) \rangle. \end{aligned} \quad (4.17)$$

Using (4.16) and Lemma 2.3 in (4.17), we get that $\{x_n\}_{n=1}^{\infty}$ converges strongly to common zero of the family $\{A_i\}_{i=1}^{\infty}$ of m -accretive operators. $\square$

Remark 4.2. If K is replaced with E in Theorems 3.5, 3.6, 3.7, and 4.1, then by Lemma 2.5, the assumption that A_i is m -accretive for each $i \geq 1$ could be replaced with A_i is continuous for each $i \geq 1$.

Hence, we have the following theorem.

Theorem 4.3. Let E be a uniformly convex real Banach space with uniformly Gâteaux differentiable norm. Let $A_i : E \rightarrow E$, $i = 1, 2, \dots$, be a countably infinite family of continuous accretive operators such that $\bigcap_{i=1}^{\infty} A_i^{-1}(0) \neq \emptyset$. For arbitrary but fixed $\delta \in (0, 1)$, $u \in K$, let $\{x_n\}_{n=1}^{\infty}$ be generated by $x_1 \in K$,

$$x_{n+1} = \alpha_n u + \sum_{i=1}^{\infty} \sigma_{i,n} ((1-\delta)x_n + \delta J_{A_i} x_n), \quad n \geq 1, \quad (4.18)$$

then, $\{x_n\}_{n=1}^{\infty}$ converges strongly to a common zero of $\{A_i\}_{i=1}^{\infty}$.

Proof. By Lemma 2.5, we have that A_i is m -accretive for each $i \geq 1$. Then, the rest of the proof follows from Theorem 4.1. $\square$

We also have the following theorems.

Theorem 4.4. Let K , E , A_i and $\{x_n\}_{n \geq 1}$ be as an Theorem 4.1. Suppose that at least one of J_{A_i} is demicompact, then $\{x_n\}_{n \geq 1}$ converges strongly to a common zero of A_i , $i = 1, 2, \dots$

Proof. The proof follows as in the proof of Theorem 4.1 but using Theorem 3.6. $\square$

Theorem 4.5. Let K , E , A_i and $\{x_n\}_{n \geq 1}$ be as an Theorem 4.1. Suppose that, in addition, E admits weakly sequential continuous duality mapping, then $\{x_n\}_{n \geq 1}$ converges strongly to a common zero of A_i , $i = 1, 2, \dots$

Proof. The proof follows as in the proof of Theorem 4.1 but using Theorem 3.7. $\square$

5. Convergence Theorems for Countably Infinite Family of Pseudocontractive Mappings

Theorem 5.1. Let K be a nonempty closed convex subset of a uniformly convex real Banach space E with uniformly Gâteaux differentiable norm. Let $T_i : K \rightarrow E$, $i = 1, 2, \dots$, be a countably infinite family of pseudocontractive mappings such that for each $i \geq 1$, $(I - T_i)$ is m -accretive and $\bigcap_{i=1}^{\infty} F(T_i) \neq \emptyset$. Let $J_{T_i} = (2I - T_i)^{-1}$, $i \geq 1$. For fixed $\delta \in [\gamma_1, \gamma_2]$, for some $\gamma_1, \gamma_2 \in (0, 1)$ and $u \in K$, let $\{x_n\}_{n=1}^{\infty}$ be generated by $x_1 \in K$:

$$x_{n+1} = \alpha_n u + \sum_{i=1}^{\infty} \sigma_{i,n} ((1 - \delta)x_n + \delta J_{T_i} x_n), \quad n \geq 1, \quad (5.1)$$

then $\{x_n\}_{n=1}^{\infty}$ converges strongly to a common fixed point of $\{T_i\}_{i=1}^{\infty}$.

Proof. Put $A_i := (I - T_i)$, $i \geq 1$. It is then obvious that $A_i^{-1}(0) = F(T_i)$, for all $i \in \mathbb{N}$ and hence $\bigcap_{i=1}^{\infty} A_i^{-1}(0) = \bigcap_{i=1}^{\infty} F(T_i)$. Furthermore, A_i is m -accretive for each $i \geq 1$. Thus, we obtain the conclusion from Theorem 4.1 with J_{A_i} in the definition of z_n replaced with J_{T_i} . $\square$

Theorem 5.2. Let E be a uniformly convex real Banach space with uniformly Gâteaux differentiable norm. For each $i \geq 1$, let $T_i : E \rightarrow E$ be a countably infinite family of continuous pseudocontractive mappings such that for each $i \geq 1$ and $\bigcap_{i=1}^{\infty} F(T_i) \neq \emptyset$. Let $J_{T_i} = (2I - T_i)^{-1}$, $i \geq 1$. For arbitrary but fixed $\delta \in [\gamma_1, \gamma_2]$, for some $\gamma_1, \gamma_2 \in (0, 1)$ and $u \in K$, let $\{x_n\}_{n=1}^{\infty}$ be generated by $x_1 \in K$:

$$x_{n+1} = \alpha_n u + \sum_{i=1}^{\infty} \sigma_{i,n} ((1 - \delta)x_n + \delta J_{T_i} x_n), \quad n \geq 1, \quad (5.2)$$

then $\{x_n\}_{n=1}^{\infty}$ converges strongly to a common fixed point of $\{T_i\}_{i=1}^{\infty}$.

Remark 5.3. Theorems similar to Theorems 4.4 and 4.5 could also be obtained for countably infinite family of pseudocontractive mappings.

Remark 5.4. Prototypes for our iteration parameters are

$$\alpha_n = \frac{1}{n+1}, \quad \sigma_{i,n} = \frac{n}{2^i(n+1)}, \quad \lambda_i = \frac{1}{2^i}. \quad (5.3)$$

Remark 5.5. The addition of bounded error terms in any of our recursion formulas leads to no further generalization.

Remark 5.6. If $f : K \rightarrow K$ is a contraction map and we replace u by $f(x_n)$ in the recursion formulas of our theorems, we obtain what some authors now call *viscosity* iteration process. We observe that all our theorems in this paper carry over trivially to the so-called viscosity process. One simply replaces u by $f(x_n)$, repeats the argument of this paper, using the fact that f is a contraction map.

Acknowledgments

The first author's research was supported by the Japanese Mori Fellowship of UNESCO at the Abdus Salam International Centre for Theoretical Physics, Trieste, Italy. The second author undertook this work when he was visiting the Abdus Salam International Centre for Theoretical Physics, Trieste, Italy.

References

- [1] C. E. Chidume, *Geometric Properties of Banach Spaces and Nonlinear Iterations*, vol. 1965 of *Lecture Notes in Mathematics*, Springer, London, UK, 2009.
- [2] Z. B. Xu and G. F. Roach, "Characteristic inequalities of uniformly convex and uniformly smooth Banach spaces," *Journal of Mathematical Analysis and Applications*, vol. 157, no. 1, pp. 189–210, 1991.
- [3] T. Kato, "Nonlinear semigroups and evolution equations," *Journal of the Mathematical Society of Japan*, vol. 19, pp. 508–520, 1967.
- [4] E. U. Ofoedu, "Iterative approximation of a common zero of a countably infinite family of m -accretive operators in Banach spaces," *Fixed Point Theory and Applications*, vol. 2008, Article ID 325792, 13 pages, 2008.
- [5] H. Zegeye and N. Shahzad, "Strong convergence theorems for a common zero for a finite family of m -accretive mappings," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 66, no. 5, pp. 1161–1169, 2007.
- [6] L. Hu and L. Liu, "A new iterative algorithm for common solutions of a finite family of accretive operators," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 70, no. 6, pp. 2344–2351, 2009.
- [7] C. H. Morales and J. S. Jung, "Convergence of paths for pseudocontractive mappings in Banach spaces," *Proceedings of the American Mathematical Society*, vol. 128, no. 11, pp. 3411–3419, 2000.
- [8] E. U. Ofoedu and Y. Shehu, "Iterative construction of a common fixed point of finite families of nonlinear mappings," *Journal of Applied Analysis*, vol. 16, no. 1, 2010, In press.
- [9] S. Reich, "Strong convergence theorems for resolvents of accretive operators in Banach spaces," *Journal of Mathematical Analysis and Applications*, vol. 75, no. 1, pp. 287–292, 1980.
- [10] C. Byrne, "A unified treatment of some iterative algorithms in signal processing and image reconstruction," *Inverse Problems*, vol. 20, no. 1, pp. 103–120, 2004.
- [11] L. C. Ceng, P. Cubiotti, and J. C. Yao, "Approximation of common fixed points of families of nonexpansive mappings," *Taiwanese Journal of Mathematics*, vol. 12, no. 2, pp. 487–500, 2008.
- [12] Y. C. Lin, N. C. Wong, and J. C. Yao, "Strong convergence theorems of Ishikawa iteration process with errors for fixed points of Lipschitz continuous mappings in Banach spaces," *Taiwanese Journal of Mathematics*, vol. 10, no. 2, pp. 543–552, 2006.
- [13] A. Petrușel and J.-C. Yao, "Viscosity approximation to common fixed points of families of nonexpansive mappings with generalized contractions mappings," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 69, no. 4, pp. 1100–1111, 2008.
- [14] J. W. Peng and J. C. Yao, "A new hybrid-extragradient method for generalized mixed equilibrium problems, fixed point problems and variational inequality problems," *Taiwanese Journal of Mathematics*, vol. 12, no. 6, pp. 1401–1433, 2008.
- [15] L. C. Zeng and J. C. Yao, "Strong convergence theorem by an extragradient method for fixed point problems and variational inequality problems," *Taiwanese Journal of Mathematics*, vol. 10, no. 5, pp. 1293–1303, 2006.
- [16] H. Brézis and P. L. Lion, "Product infinis de resolvents," *Israel Journal of Mathematics*, vol. 29, no. 4, pp. 329–345, 1978.
- [17] J. S. Jung and W. Takahashi, "Dual convergence theorems for the infinite products of resolvents in Banach spaces," *Kodai Mathematical Journal*, vol. 14, no. 3, pp. 358–365, 1991.
- [18] J. S. Jung and W. Takahashi, "On the asymptotic behavior of infinite products of resolvents in Banach spaces," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 20, no. 5, pp. 469–479, 1993.
- [19] S. Reich, "On infinite products of resolvents," *Atti della Accademia Nazionale dei Lincei*, vol. 63, no. 5, pp. 338–340, 1977.
- [20] R. T. Rockafellar, "Monotone operators and the proximal point algorithm," *SIAM Journal on Control and Optimization*, vol. 14, no. 5, pp. 877–898, 1976.

- [21] T. H. Kim and H. K. Xu, "Strong convergence of modified Mann iterations," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 61, no. 1-2, pp. 51–60, 2005.
- [22] X. Qin and Y. Su, "Approximation of a zero point of accretive operator in Banach spaces," *Journal of Mathematical Analysis and Applications*, vol. 329, no. 1, pp. 415–424, 2007.
- [23] R. Chen and Z. Zhu, "Viscosity approximation fixed points for nonexpansive and m -accretive operators," *Fixed Point Theory and Applications*, vol. 2006, Article ID 81325, 10 pages, 2006.
- [24] J. S. Jung, "Strong convergence of viscosity approximation methods for finding zeros of accretive operators in Banach spaces," *Nonlinear Analysis: Theory, Methods & Applications*. In press.
- [25] R. E. Bruck Jr., "Properties of fixed-point sets of nonexpansive mappings in Banach spaces," *Transactions of the American Mathematical Society*, vol. 179, pp. 251–262, 1973.
- [26] H. K. Xu, "Iterative algorithms for nonlinear operators," *Journal of the London Mathematical Society*, vol. 66, no. 1, pp. 1–17, 2002.
- [27] T. Suzuki, "Strong convergence of Krasnoselskii and Mann's type sequences for one-parameter non-expansive semigroups without Bochner integrals," *Journal of Mathematical Analysis and Applications*, vol. 305, no. 1, pp. 227–239, 2005.
- [28] I. Cioranescu, *Geometry of Banach Spaces, Duality Mappings and Nonlinear Problems*, vol. 62, Kluwer Academic Publishers, Dordrecht, The Netherlands, 1990.
- [29] C. E. Chidume and C. O. Chidume, "Iterative methods for common fixed points for a countable family of nonexpansive mappings in uniformly convex spaces," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 71, no. 10, pp. 4346–4356, 2008.
- [30] R. E. Megginson, *An Introduction to Banach Space Theory*, vol. 183, Springer, New York, NY, USA, 1998.
- [31] H. Y. Zhou, "Convergence theorems of common fixed points for a finite family of Lipschitz pseudocontractions in Banach spaces," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 68, no. 10, pp. 2977–2983, 2008.