Research Article

Some Identities on the q -Genocchi Polynomials of Higher-Order and q -Stirling Numbers by the Fermionic p -Adic Integral on $\mathbb{Z}_p$

**Seog-Hoon Rim, Jeong-Hee Jin, Eun-Jung Moon,
and Sun-Jung Lee**

Department of Mathematics, Kyungpook National University, Taegu 702-701, Republic of Korea

Correspondence should be addressed to Seog-Hoon Rim, shrim@knu.ac.kr

Received 25 September 2010; Accepted 8 November 2010

Academic Editor: H. Srivastava

Copyright © 2010 Seog-Hoon Rim et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

A systemic study of some families of q -Genocchi numbers and families of polynomials of Nörlund type is presented by using the multivariate fermionic p -adic integral on $\mathbb{Z}_p$. The study of these higher-order q -Genocchi numbers and polynomials yields an interesting q -analog of identities for Stirling numbers.

1. Introduction

Let p be a fixed odd prime number. Throughout this paper, $\mathbb{Z}_p$, $\mathbb{Q}_p$, $\mathbb{C}$, and $\mathbb{C}_p$ denote the ring of p -adic rational integers, the field of p -adic rational numbers, the complex number field, and the completion of the algebraic closure of $\mathbb{Q}_p$, respectively. Let $\mathbb{N}$ be the set of natural numbers and $\mathbb{Z}_+ = \mathbb{N} \cup \{0\}$. Let v_p be the normalized exponential valuation of $\mathbb{C}_p$ with $|p|_p = p^{-v_p(p)} = 1/p$.

When one talks of q -extension, q is variously considered as an indeterminate, a complex $q \in \mathbb{C}$, or a p -adic number $q \in \mathbb{C}_p$. If $q \in \mathbb{C}$, then one normally assumes $|q| < 1$. If $q \in \mathbb{C}_p$, then we assume $|q - 1|_p < 1$. In this paper, we use the following notation:

$$[x]_q = \frac{1 - q^x}{1 - q}, \quad [x]_{-q} = \frac{1 - (-q)^x}{1 + q}, \quad (1.1)$$

see [1–10]. Hence $\lim_{q \rightarrow 1} [x]_q = x$ for all $x \in \mathbb{Z}_p$.

The q -factorial is defined as $[n]_q! = [n]_q[n-1]_q \cdots [2]_q[1]_q$, and the Gaussian binomial coefficient is defined by the standard rule

$$\binom{n}{k}_q = \frac{[n]_q!}{[n-k]_q![k]_q!} = \frac{[n]_q[n-1]_q \cdots [n-k+1]_q!}{[k]_q!}, \quad (1.2)$$

(see [7, 9]). Note that $\lim_{q \rightarrow 1} \binom{n}{k}_q = \binom{n}{k} = n!/(n-k)!k! = n(n-1) \cdots (n-k+1)/k!$. It readily follows from (1.2) that

$$\binom{n+1}{k}_q = \binom{n}{k-1}_q + q^k \binom{n}{k}_q = q^{n-k+1} \binom{n}{k-1}_q + \binom{n}{k}_q, \quad (1.3)$$

(see [4, 7]).

The q -binomial formulas are known,

$$\begin{aligned} (b; q)_n &= (1-b)(1-bq) \cdots (1-bq^{n-1}) = \sum_{i=0}^n \binom{n}{i}_q q^{\binom{i}{2}} (-1)^i b^i, \\ \frac{1}{(b; q)_n} &= \frac{1}{(1-b)(1-bq) \cdots (1-bq^{n-1})} = \sum_{i=0}^{\infty} \binom{n+i-1}{i}_q b^i. \end{aligned} \quad (1.4)$$

We say that $f : \mathbb{Z}_p \rightarrow \mathbb{C}_p$ is uniformly differentiable function at a point $a \in \mathbb{Z}_p$, and we write $f \in UD(\mathbb{Z}_p)$, if the difference quotients $\Phi_f : \mathbb{Z}_p \times \mathbb{Z}_p \rightarrow \mathbb{C}_p$ such that $\Phi_f(x, y) = (f(x) - f(y))/(x - y)$ have a limit $f'(a)$ as $(x, y) \rightarrow (a, a)$. For $f \in UD(\mathbb{Z}_p)$, the q -deformed fermionic p -adic integral is defined as

$$I_{-q}(f) = \int_{\mathbb{Z}_p} f(x) d\mu_{-q}(x) = \lim_{N \rightarrow \infty} \frac{1}{[p^N]_{-q}} \sum_{x=0}^{p^N-1} f(x) (-q)^x, \quad (1.5)$$

(see [7, 9]). Note that

$$I_{-1}(f) = \lim_{q \rightarrow 1} I_{-q}(f) = \int_{\mathbb{Z}_p} f(x) d\mu_{-1}(x). \quad (1.6)$$

For $n \in \mathbb{N}$, write $f_n(x) = f(x+n)$. Then, we have

$$I_{-1}(f_n) = (-1)^n I_{-1}(f) + 2 \sum_{l=0}^{n-1} (-1)^{n-1-l} f(l). \quad (1.7)$$

Using (1.7), we can readily derive the Genocchi polynomials, $G_n(x)$, namely,

$$t \int_{\mathbb{Z}_p} e^{(x+y)t} d\mu_{-1}(y) = \frac{2t}{e^t + 1} e^{xt} = \sum_{n=0}^{\infty} G_n(x) \frac{t^n}{n!}, \quad (1.8)$$

(see [1–27]). Note that $G_n(0) = G_n$ are referred to as the n th Genocchi numbers. Let us now introduce the Genocchi polynomials of Nörlund type as follows:

$$t^r \underbrace{\int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p}}_{r \text{ times}} e^{(x+x_1+\cdots+x_r)t} d\mu_{-1}(x_1) \cdots d\mu_{-1}(x_r) = \left(\frac{2t}{e^t + 1} \right)^r e^{xt} = \sum_{n=0}^{\infty} G_n^{(r)}(x) \frac{t^n}{n!}, \quad (1.9)$$

$$\left(\frac{e^t + 1}{2t} \right)^r e^{xt} = \sum_{n=0}^{\infty} G_n^{(-r)}(x) \frac{t^n}{n!}, \quad (1.10)$$

(see [7, 9]). In the special case $x = 0$, $G_n^{(-r)}(0) = G_n^{(-r)}$, and $G_n^{(r)}(0) = G_n^{(r)}$ are referred to as the Genocchi numbers of Nörlund type. Let $(Eh)(x) = h(x+1)$ be the shift operator. Then, the q -difference operator Δ_q is defined as

$$\Delta_q^n = \prod_{i=1}^n (E - q^{i-1}I), \quad \text{where } (Ih)(x) = h(x), \quad (1.11)$$

(see [4, 7, 9]). It follows from (1.11) that

$$f(x) = \sum_{n \geq 0} \binom{x}{n}_q \Delta_q^n f(0), \quad (1.12)$$

where $\Delta_q^n f(0) = \sum_{k=0}^n \binom{n}{k}_q q^{\binom{k}{2}} f(n-k)$ (see [5, 6, 10]). The q -Stirling number of the second kind (as defined by Carlitz) is given by

$$S_2(n, k; q) = \frac{q^{-\binom{k}{2}}}{[k]_q!} \sum_{j=0}^k (-1)^j q^{\binom{j}{2}} \binom{k}{j}_q [k-j]_q^n, \quad (1.13)$$

(see [7, 10]). By (1.12) and (1.13), we see that

$$S_2(n, k; q) = \frac{q^{-\binom{k}{2}}}{[k]_q!} \Delta_q^k 0^n, \quad (1.14)$$

(see [6, 10]).

In this paper, the q -extensions of (1.9) are considered in several ways. Using these q -extensions, we derive some interesting identities and relations for Genocchi polynomials and

numbers of Nörlund type. The purpose of this paper is to present a systemic study of some families q -Genocchi numbers and polynomials of Nörlund type by using the multivariate fermionic p -adic integral on $\mathbb{Z}_p$.

2. q -Extensions of Genocchi Numbers and Polynomials of Nörlund Type

In this section, we assume that $q \in \mathbb{C}_p$ with $|1 - q|_p < 1$. We first consider the q -extensions of (1.8) given by the rule

$$\begin{aligned} \sum_{n=0}^{\infty} G_{n,q}(x) \frac{t^n}{n!} &= t \int_{\mathbb{Z}_p} e^{[x+y]_q t} d\mu_{-1}(y) \\ &= \sum_{n=0}^{\infty} \frac{2t}{(1-q)^n} \sum_{l=0}^n \left(\frac{\binom{n}{l} (-1)^l q^{lx}}{1+q^l} \right) \frac{t^n}{n!} = 2t \sum_{m=0}^{\infty} (-1)^m e^{[m+x]_q t}. \end{aligned} \quad (2.1)$$

Thus, we obtain the following lemma.

Lemma 2.1. *If $n \geq 0$, then*

$$\frac{G_{n+1,q}(x)}{n+1} = 2 \sum_{m=0}^{\infty} (-1)^m [m+x]_q^n = \frac{2}{(1-q)^n} \sum_{l=0}^n \frac{\binom{n}{l} (-1)^l q^{lx}}{1+q^l}. \quad (2.2)$$

By (1.14),

$$\begin{aligned} [x]_q^n &= \sum_{k=0}^n \binom{x}{k}_q [k]_q! S_2(k, n-k; q) q^{\binom{k}{2}} \\ &= \sum_{k=0}^n [x]_q [x-1]_q \cdots [x-k+1]_q \frac{q^{\binom{k}{2} - \binom{n-k}{2}}}{[n-k]_q!} \Delta_q^{n-k} 0^k \\ &= \sum_{k=0}^n \frac{q^{\binom{k}{2} - \binom{n-k}{2}}}{[n-k]_q!} \Delta_q^{n-k} 0^k \frac{1}{(1-q)^k} \sum_{l=0}^k \binom{k}{l}_q q^{\binom{l}{2}} (-1)^l q^{l(x-k+1)}. \end{aligned} \quad (2.3)$$

Thus, we have

$$\frac{G_{n+1,q}}{n+1} = \sum_{k=0}^n \frac{q^{\binom{k}{2}} S_2(k, n-k; q)}{(1-q)^k} \sum_{l=0}^k \binom{k}{l}_q q^{\binom{l}{2}} (-1)^l \sum_{m=0}^l \binom{l}{m} (q-1)^m \frac{G_{m+1,q}(1-k)}{m+1}, \quad (2.4)$$

and we obtain the following theorem.

Theorem 2.2. If $n \geq 0$, then

$$\frac{G_{n+1,q}}{n+1} = \sum_{k=0}^n \frac{q^{\binom{k}{2}} S_2(k, n-k; q)}{(1-q)^k} \sum_{l=0}^k \binom{k}{l}_q q^{\binom{l}{2}} (-1)^l \sum_{m=0}^l \binom{l}{m} (q-1)^m \frac{G_{m+1,q}(1-k)}{m+1}, \quad (2.5)$$

where $G_{n,q} = G_{n,q}(0)$ stand for the n th Genocchi numbers.

Consider a q -extensoin of (1.9) such that $G_{0,q}^{(r)}(x) = G_{1,q}^{(r)}(x) = \dots = G_{r-1,q}^{(r)}(x) = 0$ and

$$\begin{aligned} \frac{G_{n+r,q}^{(r)}(x)}{r! \binom{n+r}{r}} &= \int_{\mathbb{Z}_p} \dots \int_{\mathbb{Z}_p} [x + x_1 + \dots + x_r]_q^n d\mu_{-1}(x_1) \dots d\mu_{-1}(x_r) \\ &= \frac{2^r}{(1-q)^n} \sum_{l=0}^n \binom{n}{l} (-1)^l q^{lx} \left(\frac{1}{1+q^l} \right)^r = 2^r \sum_{m=0}^{\infty} \binom{m+r-1}{m} (-1)^m [m+x]_q^n. \end{aligned} \quad (2.6)$$

Let $F_q^{(r)}(t, x) = \sum_{n=0}^{\infty} G_{n,q}^{(r)}(x) (t^n/n!)$. Then,

$$F_q^{(r)}(t, x) = 2^r t^r \sum_{m=0}^{\infty} \binom{m+r-1}{m} (-1)^m e^{[m+x]_q t}. \quad (2.7)$$

In the special case $x = 0$, the numbers $G_{n,q}^{(r)}(0) = G_{n,q}^{(r)}$ are referred to as q -extension of the Genocchi numbers of order r . In the sense of the q -extension in (1.10), consider the q -extension of Genocchi polynomials of Nörlund type given by

$$G_q^{(r)}(t, x) = F_q^{(-r)}(t, x) = \frac{1}{2^r t^r} \sum_{m=0}^r \binom{r}{m} e^{[m+x]_q t} = \sum_{n=0}^{\infty} G_{n,q}^{(-r)}(x) \frac{t^n}{n!}. \quad (2.8)$$

By (2.8), $G_{0,q}^{(-r)}(x) = G_{1,q}^{(-r)}(x) = \dots = G_{r-1,q}^{(-r)}(x) = 0$ and $r! \binom{n}{r} G_{n-r,q}^{(r)}(x) = (1/2^r) \sum_{m=0}^r \binom{r}{m} [m+x]_q^n$. Therefore, we obtain the following theorem.

Theorem 2.3. For $r \in \mathbb{N}$, and, $n \geq 0$, write

$$2^r t^r \sum_{m=0}^{\infty} \binom{m+r-1}{m} (-1)^m e^{[m+x]_q t} = \sum_{n=0}^{\infty} G_{n,q}^{(r)}(x) \frac{t^n}{n!}. \quad (2.9)$$

Then,

$$\begin{aligned} \frac{G_{n+r,q}^{(r)}(x)}{r! \binom{n+r}{r}} &= \frac{2^r}{(1-q)^n} \sum_{l=0}^n \binom{n}{l} (-1)^l q^{lx} \left(\frac{1}{1+q^l} \right)^r = 2^r \sum_{m=0}^{\infty} \binom{m+r-1}{m} (-1)^m [m+x]_q^n, \\ r! \binom{n}{r} G_{n-r,q}^{(-r)}(x) &= \frac{1}{2^r (1-q)^n} \sum_{l=0}^n \binom{n}{l} (-1)^l q^{lx} (1+q^l)^r = \frac{1}{2^r} \sum_{m=0}^r \binom{r}{m} [m+x]_q^n. \end{aligned} \quad (2.10)$$

The numbers $G_{n,q}^{(-r)}(0) = G_{n,q}^{(-r)}$ are referred to as the q -extension of Genocchi numbers of Nörlund type. For $h \in \mathbb{Z}$ and $r \in \mathbb{N}$, introduce the extended higher-order q -Genocchi polynomials as follows:

$$\frac{G_{n+r,q}^{(h,r)}(x)}{r! \binom{n+r}{r}} = \int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} q^{\sum_{j=1}^r (h-j)x_j} [x + x_1 + \cdots + x_r]_q^n d\mu_{-1}(x_1) \cdots d\mu_{-1}(x_r). \quad (2.11)$$

Then,

$$\begin{aligned} \frac{G_{n+r,q}^{(h,r)}(x)}{r! \binom{n+r}{r}} &= \frac{2^r}{(1-q)^n} \sum_{l=0}^n \frac{\binom{n}{l} (-1)^l q^{lx}}{(-q^{h-1+l}; q^{-1})_r} = \frac{2^r}{(1-q)^n} \sum_{l=0}^n \frac{\binom{n}{l} (-1)^l q^{lx}}{(-q^{h-r+l}; q)_r} \\ &= 2^r \sum_{m=0}^{\infty} \binom{m+r-1}{m}_q (-1)^m q^{(h-r)m} [x+m]_q^n. \end{aligned} \quad (2.12)$$

Let $F_q^{(h,r)}(t, x) = \sum_{n=0}^{\infty} G_{n,q}^{(h,r)}(x) (t^n / n!)$. Then, we can readily see that

$$F_q^{(h,r)}(t, x) = 2^r t^r \sum_{m=0}^{\infty} \binom{m+r-1}{m}_q (-1)^m q^{(h-r)m} e^{[x+m]_q t}. \quad (2.13)$$

Therefore, we obtain the following theorem.

Theorem 2.4. For $h \in \mathbb{Z}$ and $n \geq 0$, let

$$2^r t^r \sum_{m=0}^{\infty} \binom{m+r-1}{m}_q (-1)^m q^{(h-r)m} e^{[x+m]_q t} = \sum_{n=0}^{\infty} G_{n,q}^{(h,r)}(x) \frac{t^n}{n!}. \quad (2.14)$$

Then,

$$\frac{G_{n+r,q}^{(h,r)}(x)}{r! \binom{n+r}{r}} = \frac{2^r}{(1-q)^n} \sum_{l=0}^n \frac{\binom{n}{l} (-1)^l q^{lx}}{(-q^{h-r+l}; q)_r} = 2^r \sum_{m=0}^{\infty} \binom{m+r-1}{m}_q (-1)^m q^{(h-r)m} [x+m]_q^n. \quad (2.15)$$

Let us now define the extended higher-order Nörlund type q -Genocchi polynomials as follows:

$$r! \binom{n}{r} G_{n-r,q}^{(h,-r)}(x) = \frac{1}{(1-q)^n} \sum_{l=0}^n \frac{\binom{n}{l} (-1)^l q^{lx}}{\int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} q^{l(x_1 + \cdots + x_r)} q^{\sum_{j=1}^r (h-j)x_j} d\mu_{-1}(x_1) \cdots d\mu_{-1}(x_r)}. \quad (2.16)$$

By (2.16),

$$\begin{aligned} r! \binom{n}{r} G_{n-r,q}^{(h,-r)}(x) &= \frac{1}{2^r (1-q)^n} \sum_{l=0}^n \binom{n}{l} (-1)^l q^{lx} (-q^{h-r+l}; q)_r \\ &= \frac{1}{2^r} \sum_{m=0}^r \binom{r}{m}_q q^{\binom{m}{2}} q^{(h-r)m} [m+x]_q^n. \end{aligned} \quad (2.17)$$

Let $F_q^{(h,-r)}(t, x) = \sum_{n=0}^{\infty} G_{n,q}^{(h,-r)}(x) (t^n / n!)$. Then, we have

$$F_q^{(h,-r)}(t, x) = \frac{1}{2^r t^r} \sum_{m=0}^r \binom{r}{m}_q q^{\binom{m}{2}} q^{(h-r)m} e^{[m+x]_q t}, \quad (2.18)$$

where, $G_{0,q}^{(h,-r)}(x) = G_{1,q}^{(h,-r)}(x) = \dots = G_{r-1,q}^{(h,-r)}(x) = 0$. Therefore, we obtain the following theorem.

Theorem 2.5. For $h \in \mathbb{Z}$, $n \geq 0$, and $r \in \mathbb{N}$, write

$$\frac{1}{2^r t^r} \sum_{m=0}^r \binom{r}{m}_q q^{\binom{m}{2}} q^{(h-r)m} e^{[m+x]_q t} = \sum_{n=0}^{\infty} G_{n,q}^{(h,-r)}(x) \frac{t^n}{n!}. \quad (2.19)$$

Then,

$$\begin{aligned} r! \binom{n}{r} G_{n-r,q}^{(h,-r)}(x) &= \frac{1}{2^r (1-q)^n} \sum_{l=0}^n \binom{n}{l} (-1)^l q^{lx} (-q^{h-r+l}; q)_r \\ &= \frac{1}{2^r} \sum_{m=0}^r \binom{r}{m}_q q^{\binom{m}{2}} q^{(h-r)m} [m+x]_q^n, \end{aligned} \quad (2.20)$$

where, $G_{0,q}^{(h,-r)}(x) = G_{1,q}^{(h,-r)}(x) = \dots = G_{r-1,q}^{(h,-r)}(x) = 0$.

For $h = r$,

$$\frac{G_{n+r,q}^{(r,r)}(x)}{r! \binom{n+r}{r}} = \frac{2^r}{(1-q)^n} \sum_{l=0}^n \frac{\binom{n}{l} (-1)^l q^{lx}}{(-q^l; q)_r} = 2^r \sum_{m=0}^{\infty} \binom{m+r-1}{m}_q (-1)^m [x+m]_q^n, \quad (2.21)$$

$$r! \binom{n}{r} G_{n-r,q}^{(r,-r)}(x) = \frac{1}{2^r (1-q)^n} \sum_{l=0}^n \binom{n}{l} (-1)^l q^{lx} (-q^l; q)_r = \frac{1}{2^r} \sum_{m=0}^r \binom{r}{m}_q q^{\binom{m}{2}} [m+x]_q^n. \quad (2.22)$$

It can readily be seen that

$$\begin{aligned}
 \frac{q^{mx}2^r}{(-q^{m-r};q)_r} &= \int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} q^{\sum_{j=1}^r (m-j)x_j + mx} d\mu_{-1}(x_1) \cdots d\mu_{-1}(x_r) \\
 &= \int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} ([x + x_1 + \cdots + x_r]_q (q-1) + 1)^m q^{-\sum_{j=1}^r jx_j} d\mu_{-1}(x_1) \cdots d\mu_{-1}(x_r) \\
 &= \sum_{l=0}^m \binom{m}{l} (q-1)^l \int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} [x + x_1 + \cdots + x_r]_q^l q^{-\sum_{j=1}^r jx_j} d\mu_{-1}(x_1) \cdots d\mu_{-1}(x_r) \\
 &= \sum_{l=0}^m \binom{m}{l} (q-1)^l \frac{G_{l+r,q}^{(0,r)}(x)}{r! \binom{l+r}{r}}.
 \end{aligned} \tag{2.23}$$

By (2.23), $q^{mx}2^r / (-q^{m-r};q)_r = \sum_{l=0}^m \binom{m}{l} (q-1)^l (G_{l+r,q}^{(0,r)}(x) / r! \binom{l+r}{r})$. As is known,

$$I_{-1}(f_1) + I_{-1}(f) = 2f(0), \quad \text{where } f_1(x) = f(x+1). \tag{2.24}$$

It follows from (2.24) that

$$\begin{aligned}
 q^{h-1} \int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} [x+1+x_1+\cdots+x_r]_q^n q^{\sum_{j=1}^r (h-j)x_j} d\mu_{-1}(x_1) \cdots d\mu_{-1}(x_r) \\
 = - \int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} [x+x_1+\cdots+x_r]_q^n q^{\sum_{j=1}^r (h-j)x_j} d\mu_{-1}(x_1) \cdots d\mu_{-1}(x_r) \\
 + 2 \int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} [x+x_2+\cdots+x_r]_q^n q^{\sum_{j=1}^{r-1} (h-1-j)x_{j+1}} d\mu_{-1}(x_2) \cdots d\mu_{-1}(x_r).
 \end{aligned} \tag{2.25}$$

By (2.25),

$$q^{h-1} \frac{G_{n+r,q}^{(h,r)}(x+1)}{n+r} + \frac{G_{n+r,q}^{(h,r)}(x)}{n+r} = 2G_{n+r-1,q}^{(h-1,r-1)}(x). \tag{2.26}$$

A simple manipulation shows that

$$\begin{aligned}
 q^x \int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} [x+x_1+\cdots+x_r]_q^n q^{\sum_{j=1}^r (h-j+1)x_j} d\mu_{-1}(x_1) \cdots d\mu_{-1}(x_r) \\
 = (q-1) \int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} [x+x_1+\cdots+x_r]_q^{n+1} q^{\sum_{j=1}^r (h-j)x_j} d\mu_{-1}(x_1) \cdots d\mu_{-1}(x_r) \\
 + \int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} [x+x_1+\cdots+x_r]_q^n q^{\sum_{j=1}^r (h-j)x_j} d\mu_{-1}(x_1) \cdots d\mu_{-1}(x_r).
 \end{aligned} \tag{2.27}$$

By (2.27), $q^x (G_{n+r,q}^{(h+1,r)}(x) / (n+1)) = (q-1) (G_{n+r+1,q}^{(h,r)}(x) / (n+r+1)) + (G_{n+r,q}^{(h,r)}(x) / (n+1))$.

Therefore, we obtain the following proposition.

Proposition 2.6. For $h \in \mathbb{Z}$, $r \in \mathbb{N}$ and $n \geq 0$, the following equations

$$\begin{aligned} q^{h-1} \frac{G_{n+r,q}^{(h,r)}(x+1)}{n+r} + \frac{G_{n+r,q}^{(h,r)}(x)}{n+r} &= 2G_{n+r-1,q}^{(h-1,r-1)}(x), \\ q^x \frac{G_{n+r,q}^{(h+1,r)}(x)}{n+1} &= (q-1) \frac{G_{n+r+1,q}^{(h,r)}(x)}{n+r+1} + \frac{G_{n+r,q}^{(h,r)}(x)}{n+1} \end{aligned} \quad (2.28)$$

hold. Moreover, $(q^{mx}2^r)/((-q^{m-r};q)_r) = \sum_{l=0}^m \binom{m}{l} (q-1)^l (G_{l+r,q}^{(0,r)}(x)/r! \binom{l+r}{r})$.

By (2.21),

$$\begin{aligned} \frac{G_{n+r,q^{-1}}^{(r,r)}(r-x)}{r! \binom{n+r}{r}} &= \frac{2^r}{(1-q^{-1})^n} \sum_{l=0}^n \frac{\binom{n}{l} (-1)^l q^{-l(r-x)}}{(-q^{-l}; q^{-1})_r} \\ &= (-1)^n q^{n+\binom{r}{2}} \frac{2^r}{(1-q)^n} \sum_{l=0}^n \frac{\binom{n}{l} (-1)^l q^x}{(-q^l; q)_r} = (-1)^n q^{n+\binom{r}{2}} \frac{G_{n+r,q}^{(r,r)}(x)}{r! \binom{n+r}{r}}. \end{aligned} \quad (2.29)$$

Hence,

$$\begin{aligned} \int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} [r-x+x_1+\cdots+x_r]_{q^{-1}}^n q^{-\sum_{j=1}^r (r-j)x_j} d\mu_{-1}(x_1) \cdots d\mu_{-1}(x_r) \\ = (-1)^n q^{n+\binom{r}{2}} \int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} [x+x_1+\cdots+x_r]_q^n q^{\sum_{j=1}^r (r-j)x_j} d\mu_{-1}(x_1) \cdots d\mu_{-1}(x_r). \end{aligned} \quad (2.30)$$

For $h=r$, $G_{n+r,q^{-1}}^{(r,r)}(0) = (-1)^n q^{n+\binom{r}{2}} G_{n+r,q}^{(r,r)}(r)$. It also follows from (2.26) that

$$q^{r-1} \frac{G_{n+r,q}^{(r,r)}(x+1)}{n+r} + \frac{G_{n+r,q}^{(r,r)}(x)}{n+r} = 2G_{n+r-1,q}^{(r-1,r-1)}(x). \quad (2.31)$$

The Stirling numbers of the first kind are defined as

$$\prod_{k=1}^n (1 + [k]_q z) = \sum_{k=0}^n S_1(n, k; q) z^k, \quad (2.32)$$

(see [6, 9]),

$$q^{\binom{m}{2}} \binom{r}{m}_q = \frac{q^{\binom{m}{2}} [r]_q \cdots [r-m+1]_q}{[m]_q!} = \frac{1}{[m]_q!} \prod_{k=0}^{m-1} ([r]_q - [k]_q). \quad (2.33)$$

It can readily be seen that

$$\prod_{k=0}^{n-1} (z - [k]_q) = z^n \prod_{k=0}^{n-1} \left(1 - \frac{[k]_q}{z} \right) = \sum_{k=0}^n S_1(n-1, k; q) (-1)^k z^{n-k}. \quad (2.34)$$

By (2.33) and (2.34),

$$\prod_{k=0}^{m-1} ([r]_q - [k]_q) = \sum_{k=0}^m S_1(m-1, k; q) (-1)^k [r]_q^{m-k}. \quad (2.35)$$

Formulas (2.22) and (2.35) imply the following assertion.

Proposition 2.7. For $r \in \mathbb{N}$ and $n \in \mathbb{Z}_+$,

$$r! \binom{n}{r} G_{n-r,q}^{(r,-r)}(x) = \frac{1}{2^r [m]_q!} \sum_{m=0}^r \sum_{k=0}^m S_1(m-1, k; q) (-1)^k [r]_q^{m-k} [m+x]_q^n. \quad (2.36)$$

The generalized Genocchi numbers and polynomials of Nörlund type are defined by

$$\frac{2^r t^r}{(e^{w_1 t} + 1)(e^{w_2 t} + 1) \cdots (e^{w_r t} + 1)} e^{xt} = \sum_{n=0}^{\infty} G_n^{(r)}(x | w_1, \dots, w_r) \frac{t^n}{n!}, \quad (2.37)$$

and $G_n^{(r)}(w_1, \dots, w_r) = G_n^{(r)}(0 | w_1, \dots, w_r)$. We can now also define a q -extension of (2.37) as follows. For $w_1, \dots, w_r \in \mathbb{Z}_p$ and $\delta_1, \dots, \delta_r \in \mathbb{Z}$, write

$$\frac{G_{n+r,q}^{(r)}(x | w_1, \dots, w_r; \delta_1, \dots, \delta_r)}{r! \binom{n+r}{r}} = \int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} [x_1 w_1 + \cdots + x_r w_r + x]_q^n d\mu_{-q^{\delta_1}}(x_1) \cdots d\mu_{-q^{\delta_r}}(x_r), \quad (2.38)$$

and $G_{n+r,q}^{(r)}(w_1, \dots, w_r; \delta_1, \dots, \delta_r) = G_{n+r,q}^{(r)}(0 | w_1, \dots, w_r; \delta_1, \dots, \delta_r)$. Thus,

$$\frac{G_{n+r,q}^{(r)}(x | w_1, \dots, w_r; \delta_1, \dots, \delta_r)}{r! \binom{n+r}{r}} = \frac{[2]_{q^{\delta_1}} \cdots [2]_{q^{\delta_r}}}{(1-q)^n} \sum_{l=0}^n \frac{\binom{n}{l} (-1)^l q^{lx}}{(1+q^{\delta_1+l w_1}) \cdots (1+q^{\delta_r+l w_r})}. \quad (2.39)$$

Another q -extension of Nörlund type generalized Genocchi numbers and polynomials is also of interest, namely,

$$\begin{aligned} & \frac{G_{n+r,q}^{*(r)}(x | w_1, \dots, w_r; \delta_1, \dots, \delta_r)}{r! \binom{n+r}{r}} \\ &= \int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} [x_1 w_1 + \cdots + x_r w_r + x]_q^n q^{\delta_1 x_1 + \cdots + \delta_r x_r} d\mu_{-1}(x_1) \cdots d\mu_{-1}(x_r), \end{aligned} \quad (2.40)$$

and $G_{n+r,q}^{*(r)}(w_1, \dots, w_r; \delta_1, \dots, \delta_r) = G_{n+r,q}^{*(r)}(0 \mid w_1, \dots, w_r; \delta_1, \dots, \delta_r)$. By (2.40),

$$\frac{G_{n+r,q}^{*(r)}(x \mid w_1, \dots, w_r; \delta_1, \dots, \delta_r)}{r! \binom{n+r}{r}} = \frac{2^r}{(1-q)^n} \sum_{l=0}^n \frac{\binom{n}{l} (-1)^l q^{lx}}{(1+q^{\delta_1+lw_1}) \cdots (1+q^{\delta_r+lw_r})}. \quad (2.41)$$

3. Further Remarks

For $h = 0$, consider the following polynomials $G_{n+r,q}^{(0,r)}(x)/r! \binom{n+r}{r}$ and $r! \binom{n}{r} G_{n+r,q}^{(0,-r)}(x)$:

$$\begin{aligned} \frac{G_{n+r,q}^{(0,r)}(x)}{r! \binom{n+r}{r}} &= \int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} [x + x_1 + \cdots + x_r]_q q^{-\sum_{j=1}^r jx_j} d\mu_{-1}(x_1) \cdots d\mu_{-1}(x_r), \\ r! \binom{n}{r} G_{n+r,q}^{(0,-r)}(x) &= \frac{1}{(1-q)^n} \sum_{l=0}^n \frac{\binom{n}{l} (-1)^l q^{lx}}{\int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} q^{l(x_1+\cdots+x_r)} q^{-\sum_{j=1}^r jx_j} d\mu_{-1}(x_1) \cdots d\mu_{-1}(x_r)}. \end{aligned} \quad (3.1)$$

Then,

$$\begin{aligned} \frac{G_{n+r,q}^{(0,r)}(x)}{r! \binom{n+r}{r}} &= \frac{2^r}{(1-q)^n} \sum_{l=0}^n \frac{\binom{n}{l} (-1)^l q^{lx}}{(-q^{l-r}; q)_r} = 2^r \sum_{m=0}^{\infty} \binom{m+r-1}{m}_q q^{-rm} (-1)^m [x+m]_q^n \\ r! \binom{n}{r} G_{n+r,q}^{(0,-r)}(x) &= \frac{1}{2^r (1-q)^n} \sum_{l=0}^n \binom{n}{l} (-1)^l q^{lx} (-q^{l-r}; q)_r = \frac{1}{2^r} \sum_{m=0}^r \binom{r}{m}_q q^{\binom{m}{2}} q^{-rm} [m+x]_q^n. \end{aligned} \quad (3.2)$$

Let $F_q^{(0,r)}(t, x) = \sum_{n=0}^{\infty} G_{n,q}^{(0,r)}(x) (t^n/n!)$ and let $F_q^{(0,-r)}(t, x) = \sum_{n=0}^{\infty} G_{n,q}^{(0,-r)}(x) (t^n/n!)$. Then,

$$\begin{aligned} F_q^{(0,r)}(t, x) &= 2^r t^r \sum_{m=0}^{\infty} \binom{m+r-1}{m}_q q^{-rm} (-1)^m e^{[x+m]_q t}, \\ F_q^{(0,-r)}(t, x) &= \frac{1}{2^r t^r} \sum_{m=0}^r \binom{r}{m}_q q^{\binom{m}{2}} q^{-rm} e^{[m+x]_q t}. \end{aligned} \quad (3.3)$$

Consider the following polynomials:

$$\frac{G_{n+1,q}^{(h,1)}(x)}{n+1} = \int_{\mathbb{Z}_p} q^{x_1(h-1)} [x + x_1]_q^n d\mu_{-1}(x_1) = \frac{2}{(1-q)^n} \sum_{l=0}^n \frac{\binom{n}{l} (-1)^l q^{lx}}{1+q^{l+h-1}}. \quad (3.4)$$

A simple calculation of the fermionic p -adic invariant integral on $\mathbb{Z}_p$ show that

$$\begin{aligned} q^x \int_{\mathbb{Z}_p} [x + x_1]_q^n q^{x_1(h-1)} d\mu_{-1}(x_1) \\ = (q-1) \int_{\mathbb{Z}_p} [x + x_1]_q^{n+1} q^{x_1(h-2)} d\mu_{-1}(x_1) + \int_{\mathbb{Z}_p} [x + x_1]_q^n q^{x_1(h-2)} d\mu_{-1}(x_1). \end{aligned} \quad (3.5)$$

By (3.5), $q^x G_{n+1,q}^{(h,1)}(x) = (q-1)(G_{n+2,q}^{(h-1,1)}(x)/2(n+2)) + G_{n+1,q}^{(h-1,1)}(x)$. It can readily be proved that

$$\int_{\mathbb{Z}_p} [x + x_1]_q^n q^{x_1(h-1)} d\mu_{-1}(x_1) = \sum_{j=0}^n \binom{n}{j} [x]_q^{n-j} q^{jx} \int_{\mathbb{Z}_p} [x_1]_q^j q^{x_1(h-1)} d\mu_{-1}(x_1). \quad (3.6)$$

By (3.6), $G_{n+1,q}^{(h,1)}(x)/(n+1) = \sum_{j=0}^n \binom{n}{j} [x]_q^{n-j} q^{jx} (G_{j+1,q}^{(h,1)}(1)/(j+1))$. Using (2.24), we can also prove that

$$\int_{\mathbb{Z}_p} [x + x_1 + 1]_q^n q^{(x_1+1)(h-1)} d\mu_{-1}(x_1) + \int_{\mathbb{Z}_p} [x + x_1]_q^n q^{x_1(h-1)} d\mu_{-1}(x_1) = 2[x]_q^n. \quad (3.7)$$

Thus, $q^{h-1}(G_{n+1,q}^{(h,1)}(x)/(n+1)) + (G_{n+1,q}^{(h,1)}(x)/(n+1)) = 2[x]_q^n$. For $x = 0$, we have $q^{h-1}(G_{n+1,q}^{(h,1)}(1)/(n+1)) + (G_{n+1,q}^{(h,1)}(1)/(n+1)) = 2\delta_{n,0}$, where $\delta_{n,0}$ is the Kronecker delta.

It is easy to see that $G_{1,q}^{(h,1)} = \int_{\mathbb{Z}_p} q^{x_1(h-1)} d\mu_{-1}(x_1) = 2/(1 + q^{h-1}) = 2/([2]_{q^{h-1}})$. By (3.4),

$$\begin{aligned} \frac{G_{n+1,q^{-1}}^{(h,1)}(1-x)}{n+1} &= \int_{\mathbb{Z}_p} [1-x+x_1]_{q^{-1}}^n q^{-x_1(h-1)} d\mu_{-1}(x_1) \\ &= (-1)^n q^{n+h-1} \frac{2}{(1-q)^n} \sum_{l=0}^n \frac{\binom{n}{l} (-1)^l q^{lx}}{1+q^{l+h-1}} \\ &= (-1)^n q^{n+h-1} \frac{G_{n+1,q}^{(h,1)}(x)}{n+1}. \end{aligned} \quad (3.8)$$

In particular, if $x = 1$, then $G_{n+1,q^{-1}}^{(h,1)}(0)/(n+1) = (-1)^n q^{n+h-1} (G_{n+1,q}^{(h,1)}(1)/(n+1)) = (-1)^{n-1} q^n (G_{n+1,q}^{(h,1)}(1)/(n+1))$ for $n \geq 1$.

Recently, Kim has studied p -adic fermionic integral on $\mathbb{Z}_p$ connected with the problems of mathematical physics (see [6, 10, 11]), and our result are closely related to his results. In the future, we will try to study p -adic stochastic problems associated with our theorems. For example, p -adic q -Bernstein polynomials seem to be closely related to our results (see [6, 14, 20]).

References

- [1] I. N. Cangul, H. Ozden, and Y. Simsek, "A new approach to q -Genocchi numbers and their interpolation functions," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 71, no. 12, pp. e793–e799, 2009.
- [2] J. Choi, P. J. Anderson, and H. M. Srivastava, "Some q -extensions of the Apostol-Bernoulli and the Apostol-Euler polynomials of order n , and the multiple Hurwitz zeta function," *Applied Mathematics and Computation*, vol. 199, no. 2, pp. 723–737, 2008.
- [3] L. Comtet, *Advanced Combinatorics*, D. Reidel, Dordrecht, The Netherlands, 1974.
- [4] E. Y. Deeba and D. M. Rodriguez, "Stirling's series and Bernoulli numbers," *The American Mathematical Monthly*, vol. 98, no. 5, pp. 423–426, 1991.
- [5] M. Cenkci, M. Can, and V. Kurt, " p -adic interpolation functions and kummer-type congruences for q -twisted euler numbers," *Advanced Studies in Contemporary Mathematics*, vol. 9, no. 2, pp. 203–216, 2004.
- [6] T. Kim, S. D. Kim, and D.-W. Park, "On uniform differentiability and q -Mahler expansions," *Advanced Studies in Contemporary Mathematics*, vol. 4, no. 1, pp. 35–41, 2001.
- [7] T. Kim, "Some identities on the q -Euler polynomials of higher order and q -Stirling numbers by the fermionic p -adic integral on $\mathbb{Z}_p$," *Russian Journal of Mathematical Physics*, vol. 16, no. 4, pp. 484–491, 2009.
- [8] T. Kim, "The modified q -Euler numbers and polynomials," *Advanced Studies in Contemporary Mathematics*, vol. 16, no. 2, pp. 161–170, 2008.
- [9] T. Kim, "Note on the Euler q -zeta functions," *Journal of Number Theory*, vol. 129, no. 7, pp. 1798–1804, 2009.
- [10] T. Kim, " q -Volkenborn integration," *Russian Journal of Mathematical Physics*, vol. 9, no. 2, pp. 288–299, 2002.
- [11] T. Kim, "Euler numbers and polynomials associated with zeta functions," *Abstract and Applied Analysis*, Article ID 581582, 11 pages, 2008.
- [12] T. Kim, "A note on p -adic q -integral on $\mathbb{Z}_p$ associated with q -Euler numbers," *Advanced Studies in Contemporary Mathematics*, vol. 15, no. 2, pp. 133–137, 2007.
- [13] T. Kim, L.-C. Jang, and H. Yi, "A note on the modified q -bernstein polynomials," *Discrete Dynamics in Nature and Society*, vol. 2010, Article ID 706483, 12 pages, 2010.
- [14] T. Kim, J. Choi, and Y.-H. Kim, "Some identities on the q -Bernstein polynomials, q -Stirling numbers and q -Bernoulli numbers," *Advanced Studies in Contemporary Mathematics*, vol. 20, no. 3, pp. 335–341, 2010.
- [15] T. Kim, "Power series and asymptotic series associated with the q -analog of the two-variable p -adic L -function," *Russian Journal of Mathematical Physics*, vol. 12, no. 2, pp. 186–196, 2005.
- [16] T. Kim, "Non-Archimedean q -integrals associated with multiple Changhee q -Bernoulli polynomials," *Russian Journal of Mathematical Physics*, vol. 10, no. 1, pp. 91–98, 2003.
- [17] T. Kim, " q -Euler numbers and polynomials associated with p -adic q -integrals," *Journal of Nonlinear Mathematical Physics*, vol. 14, no. 1, pp. 15–27, 2007.
- [18] Q.-M. Luo, " q -extensions for the Apostol-Genocchi polynomials," *General Mathematics*, vol. 17, no. 2, pp. 113–125, 2009.
- [19] Q.-M. Luo, "Fourier expansions and integral representations for Genocchi polynomials," *Journal of Integer Sequences*, vol. 12, no. 1, article 09.1.4, 2009.
- [20] Q.-M. Luo, "Some results for the q -Bernoulli and q -Euler polynomials," *Journal of Mathematical Analysis and Applications*, vol. 363, no. 1, pp. 7–18, 2010.
- [21] H. Ozden, Y. Simsek, S.-H. Rim, and I. N. Cangul, "A note on p -adic q -Euler measure," *Advanced Studies in Contemporary Mathematics*, vol. 14, no. 2, pp. 233–239, 2007.
- [22] K. i Shiratani and S. Yamamoto, "On a p -adic interpolation function for the Euler numbers and its derivatives," *Memoirs of the Faculty of Science, Kyushu University A*, vol. 39, no. 1, pp. 113–125, 1985.
- [23] Y. Simsek and M. Acikgoz, "A new generating function of (q) Bernstein-type polynomials and their interpolation function," *Abstract and Applied Analysis*, Article ID 769095, 12 pages, 2010.
- [24] Y. Simsek, "On p -Adic Twisted q - L -functions related to generalized twisted bernoulli numbers," *Russian Journal of Mathematical Physics*, vol. 13, no. 3, pp. 340–348, 2006.

- [25] Y. Simsek, "Theorems on twisted L -function and twisted Bernoulli numbers," *Advanced Studies in Contemporary Mathematics*, vol. 11, no. 2, pp. 205–218, 2005.
- [26] Y. Simsek, " q -Dedekind type sums related to q -zeta function and basic L -series," *Journal of Mathematical Analysis and Applications*, vol. 318, no. 1, pp. 333–351, 2006.
- [27] H. J. H. Tuentler, "A symmetry of power sum polynomials and Bernoulli numbers," *The American Mathematical Monthly*, vol. 108, no. 3, pp. 258–261, 2001.