Research Article

On a New Summation Formula for ${}_2\psi_2$ Basic Bilateral Hypergeometric Series and Its Applications

D. D. Somashekara, K. Narasimha Murthy, and S. L. Shalini

Department of Studies in Mathematics, University of Mysore, Manasagangotri, Mysore 570006, India

Correspondence should be addressed to K. Narasimha Murthy, simhamurth@yahoo.com

Received 7 December 2010; Accepted 24 January 2011

Academic Editor: J. Dydak

Copyright © 2011 D. D. Somashekara et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We have obtained a new summation formula for ${}_2\psi_2$ bilateral basic hypergeometric series by the method of parameter augmentation and demonstrated its various uses leading to some development of q -gamma, and q -beta function identities.

1. Introduction

The summation formulae for hypergeometric series form a very interesting and useful component of the theory of (basic) hypergeometric series. The q -binomial theorem of Cauchy [1] is perhaps the first identity in the class of the summation formulae, which can be stated as

$$\sum_{k=0}^{\infty} \frac{(a)_k}{(q)_k} z^k = \frac{(az)_{\infty}}{(z)_{\infty}}, \quad |z| < 1, \quad |q| < 1, \quad (1.1)$$

where

$$\begin{aligned} (a)_{\infty} &:= (a; q)_{\infty} := \prod_{k=0}^{\infty} (1 - aq^k), \\ (a)_k &:= (a; q)_k := \frac{(a)_{\infty}}{(aq^k)_{\infty}}, \quad k \text{ is an integer.} \end{aligned} \quad (1.2)$$

For more details about the q -binomial theorem and about the identities which fall in this sequel, one may refer to the book by Gasper and Rahman [2]. Another famous identity in the sequel is the Ramanujan's ${}_1\psi_1$ summation formula [3]

$$\sum_{k=-\infty}^{\infty} \frac{(a)_k}{(b)_k} z^k = \frac{(az)_{\infty} (q)_{\infty} (q/az)_{\infty} (b/a)_{\infty}}{(z)_{\infty} (b)_{\infty} (b/az)_{\infty} (q/a)_{\infty}}, \quad \left| \frac{b}{a} \right| < |z| < 1, \quad |q| < 1. \quad (1.3)$$

There are a number of proofs of the ${}_1\psi_1$ summation formula (1.3) in the literature. For more details, one refers to the book by Berndt [4] and a recent paper of Johnson [5].

In this paper, we derive a new summation formula for ${}_2\psi_2$ basic bilateral hypergeometric series using the ${}_1\psi_1$ summation formula (1.3) by the method of parameter augmentation. We then use the formula to derive the q -analogue of Gauss summation formula and to obtain a number of q -eta function, q -gamma, and q -beta function identities, which complement the works of Bhargava and Somashekara [6], Bhargava et al. [7], Somashekara and Mamta [8], Srivastava [9], and Bhargava and Adiga [10].

First, we recall that q -difference operator and the q -shift operator are defined by

$$D_q f(a) = \frac{f(a) - f(aq)}{a}, \quad \zeta(f(a)) = f(aq), \quad (1.4)$$

respectively. In [11], Chen and Liu have constructed an operator θ as

$$\theta = \zeta^{-1} D_q, \quad (1.5)$$

and thereby they defined the operator $E(b\theta)$ as

$$E(b\theta) = \sum_{k=0}^{\infty} \frac{(b\theta)^k q^{k(k-1)/2}}{(q; q)_k}. \quad (1.6)$$

Then, we have the following operator identities [12, Theorem 1]:

$$\begin{aligned} E(b\theta) \{ (at; q)_{\infty} \} &= (at, bt; q)_{\infty}, \\ E(b\theta) \{ (as, at; q)_{\infty} \} &= \frac{(as, at, bs, bt; q)_{\infty}}{(abst/q; q)_{\infty}}, \quad \left| \frac{abst}{q} \right| < 1. \end{aligned} \quad (1.7)$$

Further, the Dedekind etafunction is defined by

$$\eta(\tau) := e^{\pi i \tau / 12} \prod_{k=1}^{\infty} (1 - e^{2\pi i k \tau}) := q^{1/24} (q; q)_{\infty}, \quad (1.8)$$

where $q = e^{2\pi i \tau}$, and $\text{Im}(\tau) > 0$.

Jackson [13] defined the q -analogue of the gamma function by

$$\Gamma_q(x) = \frac{(q; q)_{\infty}}{(q^x; q)_{\infty}} (1 - q)^{1-x}, \quad 0 < q < 1. \quad (1.9)$$

In his paper on the q -gamma and q -beta function, Askey [14] has obtained q -analogues of several classical results about the gamma function. Further, he has given the definition for q -beta function as

$$B_q(x, y) = (1 - q) \sum_{n=0}^{\infty} q^{nx} \frac{(q^{n+1})_{\infty}}{(q^{x+y})_{\infty}}. \quad (1.10)$$

In fact, he has shown that

$$B_q(x, y) = \frac{\Gamma_q(x) \Gamma_q(y)}{\Gamma_q(x + y)}. \quad (1.11)$$

In Section 2, we prove our main result. In Section 3, we deduce the well-known q -analogue of the Gauss summation formula and some etafunction, q -gamma, and q -beta function identities.

2. Main Result

Theorem 2.1. *If $0 < |z| < 1$, $|q| < 1$, then*

$$\sum_{k=-\infty}^{\infty} \frac{(a)_k (bc/azq)_k}{(b)_k (c)_k} z^k = \frac{(az)_{\infty} (q)_{\infty} (q/az)_{\infty} (b/a)_{\infty} (c/a)_{\infty} (bc/azq)_{\infty}}{(z)_{\infty} (b)_{\infty} (c)_{\infty} (b/az)_{\infty} (c/az)_{\infty} (q/a)_{\infty}}. \quad (2.1)$$

Proof. Ramanujan's ${}_1\psi_1$ summation formula (1.3) can be written as

$$\sum_{k=0}^{\infty} \frac{(a)_k}{(b)_k} z^k + \sum_{k=1}^{\infty} \frac{(q/b)_k}{(q/a)_k} \left(\frac{b}{az} \right)^k = \frac{(az)_{\infty} (q)_{\infty} (q/az)_{\infty} (b/a)_{\infty}}{(z)_{\infty} (b)_{\infty} (b/az)_{\infty} (q/a)_{\infty}}. \quad (2.2)$$

This is the same as

$$\begin{aligned} & \sum_{k=0}^{\infty} (a)_k z^k \left\{ (bq^k)_{\infty} \left(\frac{b}{az} \right)_{\infty} \right\} + \sum_{k=1}^{\infty} \frac{(-1)^k q^{k(k+1)/2}}{(q/a)_k} \left(\frac{1}{az} \right)^k \left\{ (bq^{-k})_{\infty} \left(\frac{b}{az} \right)_{\infty} \right\} \\ &= \frac{(az)_{\infty} (q)_{\infty} (q/az)_{\infty}}{(z)_{\infty} (q/a)_{\infty}} \left\{ \left(\frac{b}{a} \right)_{\infty} \right\}. \end{aligned} \quad (2.3)$$

On applying $E(c\theta)$ to both sides with respect to b , we obtain

$$\begin{aligned} & \sum_{k=0}^{\infty} (a)_k z^k \left\{ \frac{(bq^k)_{\infty} (b/az)_{\infty} (cq^k)_{\infty} (c/az)_{\infty}}{(bcq^k/azq)_{\infty}} \right\} \\ &+ \sum_{k=1}^{\infty} \frac{(-1)^k q^{k(k+1)/2}}{(q/a)_k} \left(\frac{1}{az} \right)^k \left\{ \frac{(bq^{-k})_{\infty} (b/az)_{\infty} (cq^{-k})_{\infty} (c/az)_{\infty}}{(bcq^{-k}/azq)_{\infty}} \right\} \\ &= \frac{(az)_{\infty} (q)_{\infty} (q/az)_{\infty}}{(z)_{\infty} (q/a)_{\infty}} \left\{ \left(\frac{b}{a} \right)_{\infty} \left(\frac{c}{a} \right)_{\infty} \right\}. \end{aligned} \quad (2.4)$$

Multiplying (2.4) throughout by $\{(bc/azq)_{\infty}/(b)_{\infty}(c)_{\infty}\}$, we obtain

$$\begin{aligned} & \sum_{k=0}^{\infty} \frac{(a)_k (bc/azq)_k}{(b)_k (c)_k} z^k + \sum_{k=1}^{\infty} \frac{(a)_{-k} (bc/azq)_{-k}}{(b)_{-k} (c)_{-k}} z^{-k} \\ &= \frac{(az)_{\infty} (q)_{\infty} (q/az)_{\infty} (b/a)_{\infty} (c/a)_{\infty} (bc/azq)_{\infty}}{(z)_{\infty} (b)_{\infty} (c)_{\infty} (b/az)_{\infty} (c/az)_{\infty} (q/a)_{\infty}}, \end{aligned} \quad (2.5)$$

which yields (2.1). □

3. Some Applications of the Main Identity

The following identity is the well-known q -analogue of the Gauss summation formula.

Corollary 3.1 (see [15]). *If $|q| < 1$, $|\gamma/\alpha\beta| < 1$, then*

$$\sum_{k=0}^{\infty} \frac{(\alpha)_k (\beta)_k}{(q)_k (\gamma)_k} \left(\frac{\gamma}{\alpha\beta} \right)^k = \frac{(\gamma/\alpha)_{\infty} (\gamma/\beta)_{\infty}}{(\gamma)_{\infty} (\gamma/\alpha\beta)_{\infty}}. \quad (3.1)$$

Proof. Putting $a = \alpha$, $b = q$, $c = \gamma$, and $z = \gamma/\alpha\beta$ in (2.1), we obtain (3.1). □

Corollary 3.2. *If $|q| < 1$, then*

$$\sum_{k=-\infty}^{\infty} \frac{(q^2; q^4)_k}{(1 - q^{4k+1})(q^3; q^4)_k} q^k = q^{-1/8} \frac{\eta^7(2\tau)}{\eta^3(\tau)\eta^2(4\tau)}, \quad (3.2)$$

$$\sum_{k=-\infty}^{\infty} \frac{(-q^2; q^2)_k}{(1 + q^{2k+1})^2 (-q; q^2)_k} q^k = 2q^{-3/4} \frac{\eta^6(4\tau)}{\eta^3(2\tau)}, \quad (3.3)$$

$$\sum_{k=-\infty}^{\infty} \frac{(-q; q^2)_k (-q^2; q^2)_k}{(1 - q^{2k+1})^2 (q; q^2)_n^2} q^k = 2q^{-3/4} \frac{\eta^8(4\tau)}{\eta^7(2\tau)}, \quad (3.4)$$

$$\sum_{k=-\infty}^{\infty} \frac{(1+q)(-q; q^2)_k (-q^2; q^2)_k}{(1 - q^{2k+2})(q^2; q^2)_k^2} q^k = \frac{\eta(2\tau)}{\eta^2(\tau)}, \quad (3.5)$$

$$\sum_{k=-\infty}^{\infty} \frac{(q^2; q^6)_k}{(1 - q^{6k+1})(q^5; q^6)_k} q^{3k} = q^{-1/4} \frac{\eta^4(2\tau)}{\eta^2(\tau)}, \quad (3.6)$$

$$\sum_{k=-\infty}^{\infty} \frac{(1+q^2)^2 (-q^2; q^2)_{k-1} (-q^2; q^2)_{k+1}}{(1 - q^{2k+2})^2 (q^2; q^2)_k^2} q^k = \frac{\eta^2(4\tau)}{\eta^4(2\tau)}. \quad (3.7)$$

Proof. Putting $a = z = q^{1/4}$, $b = q^{3/4}$, $c = q^{5/4}$, and then changing q to q^4 in (2.1), we obtain

$$\sum_{k=-\infty}^{\infty} \frac{(q; q^4)_k (q^2; q^4)_k}{(q; q^4)_{k+1} (q^3; q^4)_k} q^k = \frac{(q^2; q^4)_{\infty}^4 (q^4; q^4)_{\infty}^2}{(q; q^4)_{\infty}^3 (q^3; q^4)_{\infty}^3}. \quad (3.8)$$

Simplifying the right hand side and then using (1.8), we obtain (3.2).

Similarly, putting $a = -q^{1/2}$, $z = q^{1/2}$, $b = -q^{3/2}$, $c = -q^{5/2}$, and then changing q to q^2 , we obtain (3.3). Putting $a = -q^{1/2}$, $z = q^{1/2}$, $b = c = q^{3/2}$, and then changing q to q^2 , we obtain (3.4). Putting $a = -q^{1/2}$, $z = q^{1/2}$, $b = q$, $c = q^2$, and then changing q to q^2 , we obtain (3.5). Putting $a = q^{1/6}$, $z = q^{1/2}$, $b = q^{7/6}$, $c = q^{5/6}$, and then changing q to q^6 , we obtain (3.6). Finally, putting $a = -1$, $z = q$, $b = c = q^2$, and then changing q to q^2 , we obtain (3.7). $\square$

Corollary 3.3. *If $0 < q < 1$, $0 < x, y < 1$, and $0 < x + y < 1$, then*

$$B_q^3(x, y) = \frac{\Gamma_q(1-x+y)\Gamma_q(x-y)(1-q)^2}{(1-q^y)^2} \sum_{k=-\infty}^{\infty} \frac{(q^{1-x})_k (q^{x+y})_k}{(q^{1+y})_k^2} q^{ky}. \quad (3.9)$$

Proof. Putting $a = q^{1-x}$, $z = q^y$, and $b = c = q^{1+y}$ in (2.1), we get

$$\sum_{k=-\infty}^{\infty} \frac{(q^{1-x})_k (q^{x+y})_k}{(q^{1+y})_k^2} q^{ky} = \frac{(q^{1-x+y})_{\infty} (q)_{\infty} (q^{x-y})_{\infty}}{(q^y)_{\infty} (q^{1+y})_{\infty} (q^{1+y})_{\infty}} \left(\frac{(q^{x+y})_{\infty}}{(q^x)_{\infty}} \right)^3. \quad (3.10)$$

On using (1.9), (1.10), and (1.11), we obtain (3.9). $\square$

Corollary 3.4. *If $0 < q < 1$, $0 < x, y < 1$, and $1 < x + y < 2$, then*

$$B_q^2(x, y) = \frac{\Gamma_q(y-x)\Gamma_q(x-y+1)\Gamma_q(x+y-1)}{\Gamma_q(y)\Gamma_q(1+x)} \sum_{k=-\infty}^{\infty} \frac{(q^{-x})_k (q^{x+y-1})_k}{(q^y)_k^2} q^{ky}. \quad (3.11)$$

Proof. Putting $a = q^{-x}$, $z = b = c = q^y$, in (2.1), and then using (1.9), (1.10), and (1.11), we obtain (3.11). $\square$

Corollary 3.5. *If $0 < x, y < 1$, and $0 < x + y < 1$, then*

$$B^3(x, y) = \frac{\Gamma(1-x+y)\Gamma(x-y)}{y^2} \left[\sum_{k=0}^{\infty} \frac{(1-x)_k (x+y)_k}{(1+y)_k^2} + \sum_{k=1}^{\infty} \frac{(-y)_k^2}{(x)_k (1-x-y)_k} \right]. \quad (3.12)$$

Proof. Letting $q \rightarrow 1$ in (3.9), we obtain (3.12). $\square$

Corollary 3.6. *If $0 < x, y < 1$, and $1 < x + y < 2$, then*

$$B^2(x, y) = \frac{\Gamma(y-x)\Gamma(x-y+1)\Gamma(x+y-1)}{\Gamma(y)\Gamma(1+x)} \times \left[\sum_{k=0}^{\infty} \frac{(-x)_k (x+y-1)_k}{(y)_k^2} + \sum_{k=1}^{\infty} \frac{(1-y)_k^2}{(1+x)_k (2-x-y)_k} \right]. \quad (3.13)$$

Proof. Letting $q \rightarrow 1$ in (3.11), we obtain (3.13). $\square$

Corollary 3.7. *If $0 < q < 1$, $0 < x, y < 1$, and $0 < x + y < 1$, then*

$$B_q(x, y) = \Gamma_q(y)\Gamma_q(1-y) \sum_{k=0}^{\infty} \frac{(q^{1-x-y})_k (q^y)_k}{(q)_k^2} q^{kx}. \quad (3.14)$$

Proof. Putting $a = q^{1-x-y}$, $z = q^x$, and $b = c = q$ in (2.1), we obtain (3.14). $\square$

Acknowledgment

The authors would like to thank the referees for their valuable suggestions which considerably improved the quality of the paper.

References

- [1] A.-L. Cauchy, "Mémoire sur les fonctions dont plusieurs valeurs sont liées entre elles par une équation linéaire, et sur diverses transformations de produits composés d'un nombre indéfini de facteurs," *Comptes Rendus de l'Académie des Sciences, Paris*, vol. 17, pp. 42-50, 1893, Oeuvres de Cauchy, 1re érie, T. VIII, Gauthier-Villars, Paris, France.

- [2] G. Gasper and M. Rahman, *Basic Hypergeometric Series*, vol. 96 of *Encyclopedia of Mathematics and Its Applications*, Cambridge University Press, Cambridge, UK, 2nd edition, 2004.
- [3] S. Ramanujan, *Notebooks*, vol. 2, Tata Institute of Fundamental Research, Bombay, India, 1957.
- [4] B. C. Berndt, *Ramanujan's Notebooks. Part III*, Springer, New York, NY, USA, 1991.
- [5] W. P. Johnson, "How Cauchy missed Ramanujan's ${}_1\psi_1$ summation," *American Mathematical Monthly*, vol. 111, no. 9, pp. 791–800, 2004.
- [6] S. Bhargava and D. D. Somashekara, "Some eta-function identities deducible from Ramanujan's ${}_1\psi_1$ summation," *Journal of Mathematical Analysis and Applications*, vol. 176, no. 2, pp. 554–560, 1993.
- [7] S. Bhargava, D. D. Somashekara, and S. N. Fathima, "A new transformation for basic bilateral hypergeometric series," *Mathematics Student*, vol. 74, no. 1–4, pp. 225–230, 2005.
- [8] D. D. Somashekara and D. Mamta, "On some identities of Ramanujan found in his lost notebook," *Advanced Studies in Contemporary Mathematics*, vol. 16, no. 2, pp. 171–180, 2008.
- [9] H. M. Srivastava, "Some convolution identities based upon Ramanujan's bilateral sum," *Bulletin of the Australian Mathematical Society*, vol. 49, pp. 433–427, 1994.
- [10] S. Bhargava and C. Adiga, "A basic bilateral series summation formula and its applications," *Integral Transforms and Special Functions*, vol. 2, no. 3, pp. 165–184, 1994.
- [11] W. Y. C. Chen and Z.-G. Liu, "Parameter augmentation for basic hypergeometric series. I," in *Mathematical Essays in Honor of Gian-Carlo Rota (Cambridge, MA, 1996)*, B. E. Sagan and R. P. Stanley, Eds., vol. 161 of *Progress in Mathematics*, pp. 111–129, Birkhäuser, Boston, Mass, USA, 1998.
- [12] Z.-G. Liu, "Some operator identities and q -series transformation formulas," *Discrete Mathematics*, vol. 265, no. 1–3, pp. 119–139, 2003.
- [13] F. H. Jackson, "On q -definite integrals," *Pure and Applied Mathematics Quarterly*, vol. 41, pp. 193–203, 1910.
- [14] R. Askey, "The q -gamma and q -beta functions," *Applicable Analysis*, vol. 8, no. 2, pp. 125–141, 1978.
- [15] E. Heine, "Untersuchungen über die Reihe $1 + [(1 - q^\alpha)(1 - q^\beta)] / [(1 - q)(1 - q^\gamma)]X + [(1 - q^\alpha)(1 - q^{\alpha+1})(1 - q^\beta)'(1 - q^{\beta+1})] / [(1 - q)(1 - q^2)(1 - q^\gamma)(1 - q^{\gamma+1})]X^2 + \dots$," *Journal für die Reine und Angewandte Mathematik*, vol. 34, pp. 285–328, 1847.