Research Article

Fekete-Szegő Problem for a New Class of Analytic Functions

Deepak Bansal

*Department of Mathematics, College of Engineering and Technology,
 Bikaner 334004, Rajasthan, India*

Correspondence should be addressed to Deepak Bansal, deepakbansal.79@yahoo.com

Received 2 December 2010; Revised 8 March 2011; Accepted 15 June 2011

Academic Editor: Attila Gilányi

Copyright © 2011 Deepak Bansal. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We consider the Fekete-Szegő problem with complex parameter μ for the class $R_V^r(\phi)$ of analytic functions.

1. Introduction and Preliminaries

Let $\mathcal{A}$ denote the class of functions of the form

$$f(z) = z + \sum_{k=2}^{\infty} a_k z^k, \quad (1.1)$$

which are analytic in the open unit disk $\mathbb{U} = \{z : z \in \mathbb{C} \text{ and } |z| < 1\}$ and $\mathcal{S}$ denote the subclass of $\mathcal{A}$ that are univalent in $\mathbb{U}$. A function $f(z)$ in $\mathcal{A}$ is said to be in class $\mathcal{S}^*$ of starlike functions of order zero in $\mathbb{U}$, if $\Re(zf'(z)/f(z)) > 0$ for $z \in \mathbb{U}$. Let $\mathcal{K}$ denote the class of all functions $f \in \mathcal{A}$ that are convex. Further, f is convex if and only if $zf'(z)$ is star-like. A function $f \in \mathcal{A}$ is said to be close-to-convex with respect to a fixed star-like function $g \in \mathcal{S}^*$ if and only if $\Re(zf'(z)/g(z)) > 0$ for $z \in \mathbb{U}$. Let $\mathcal{C}$ denote of all such close-to-convex functions [1].

Fekete and Szegő proved a noticeable result that the estimate

$$|a_3 - \lambda a_2^2| \leq 1 + 2 \exp\left(\frac{-2\lambda}{1-\lambda}\right) \quad (1.2)$$

holds for any normalized univalent function $f(z)$ of the form (1.1) in the open unit disk $\mathbb{U}$ and for $0 \leq \lambda \leq 1$. This inequality is sharp for each λ (see [2]). The coefficient functional

$$\phi_\lambda(f) = a_3 - \lambda a_2^2 = \frac{1}{6} \left(f'''(0) - \frac{3\lambda}{2} [f''(0)]^2 \right), \quad (1.3)$$

on normalized analytic functions f in the unit disk represents various geometric quantities, for example, when $\lambda = 1$, $\phi_\lambda(f) = a_3 - a_2^2$, becomes $S_f(0)/6$, where S_f denote the Schwarzian derivative $(f'''/f')' - (f''/f')^2/2$ of locally univalent functions f in $\mathbb{U}$. In literature, there exists a large number of results about inequalities for $\phi_\lambda(f)$ corresponding to various subclasses of $\mathcal{S}$. The problem of maximising the absolute value of the functional $\phi_\lambda(f)$ is called the Fekete-Szegő problem; see [2]. In [3], Koepf solved the Fekete-Szegő problem for close-to-convex functions and the largest real number λ for which $\phi_\lambda(f)$ is maximised by the Koebe function $z/(1-z)^2$ is $\lambda = 1/3$, and later in [4] (see also [5]), this result was generalized for functions that are close-to-convex of order β .

Let $\phi(z)$ be an analytic function with positive real part on $\mathbb{U}$ with $\phi(0) = 1$, $\phi'(0) > 0$ which maps the unit disk $\mathbb{U}$ onto a star-like region with respect to 1 which is symmetric with respect to the real axis. Let $\mathcal{S}^*(\phi)$ be the class of functions in $f \in \mathcal{S}$ for which

$$\frac{zf'(z)}{f(z)} < \phi(z) \quad (z \in \mathbb{U}), \quad (1.4)$$

and $\mathcal{C}(\phi)$ be the class of functions in $f \in \mathcal{S}$ for which

$$1 + \frac{zf''(z)}{f'(z)} < \phi(z) \quad (z \in \mathbb{U}), \quad (1.5)$$

where $<$ denotes the subordination between analytic functions. These classes were introduced and studied by Ma and Minda [6]. They have obtained the Fekete-Szegő inequality for the functions in the class $\mathcal{C}(\phi)$.

Motivated by the class $R_\lambda^\tau(\beta)$ in paper [7], we introduce the following class.

Definition 1.1. Let $0 \leq \gamma \leq 1$, $\tau \in \mathbb{C} \setminus \{0\}$. A function $f \in \mathcal{A}$ is in the class $R_\gamma^\tau(\phi)$, if

$$1 + \frac{1}{\tau} (f'(z) + \gamma z f''(z) - 1) < \phi(z) \quad (z \in \mathbb{U}), \quad (1.6)$$

where $\phi(z)$ is defined the same as above.

If we set

$$\phi(z) = \frac{1 + Az}{1 + Bz} \quad (-1 \leq B < A \leq 1; z \in \mathbb{U}), \quad (1.7)$$

in (1.6), we get

$$R_\gamma^\tau \left(\frac{1 + Az}{1 + Bz} \right) = R_\gamma^\tau(A, B) = \left\{ f \in \mathcal{A} : \left| \frac{f'(z) + \gamma z f''(z) - 1}{\tau(A - B) - B(f'(z) + \gamma z f''(z) - 1)} \right| < 1 \right\}, \quad (1.8)$$

which is again a new class. We list few particular cases of this class discussed in the literature

- (1) $R_Y^\tau(1 - 2\beta, -1) = R_Y^\tau(\beta)$ for $0 \leq \beta < 1$, $\tau \in \mathbb{C} \setminus \{0\}$ was discussed recently by Swaminathan [7].
- (2) The class $R_Y^\tau(1 - 2\beta, -1)$ for $\tau = e^{i\eta} \cos \eta$, where $-\pi/2 < \eta < \pi/2$ is considered in [8] (see also [9]).
- (3) The class $R_1^\tau(0, -1)$ with $\tau = e^{i\eta} \cos \eta$ was considered in [10] with reference to the univalence of partial sums.
- (4) $f \in R_Y^{e^{i\eta} \cos \eta}(1 - 2\beta, -1)$ whenever $zf'(z) \in P_Y^\tau(\beta)$, the class considered in [11].

For geometric aspects of these classes, see the corresponding references. The class $R_Y^\tau(A, B)$ is new as the author Swaminathan [7] has introduced class $R_Y^\tau(\beta)$ which is subclass of the class $R_Y^\tau(A, B)$, in his recent paper. To prove our main result, we need the following lemma.

Lemma 1.2 (see [12, 13]). *If $p(z) = 1 + c_1z + c_2z^2 + c_3z^3 + \dots$ ($z \in \mathbb{U}$) is a function with positive real part, then for any complex number μ ,*

$$|c_2 - \mu c_1^2| \leq 2 \max\{1, |2\mu - 1|\}, \quad (1.9)$$

and the result is sharp for the functions given by

$$p(z) = \frac{1+z^2}{1-z^2}, \quad p(z) = \frac{1+z}{1-z} \quad (z \in \mathbb{U}). \quad (1.10)$$

2. Fekete-Szegő Problem

Our main result is the following theorem.

Theorem 2.1. *Let $\phi(z) = 1 + B_1z + B_2z^2 + B_3z^3 + \dots$, where $\phi(z) \in \mathcal{A}$ with $\phi'(0) > 0$. If $f(z)$ given by (1.1) belongs to $R_Y^\tau(\phi)$ ($0 \leq \gamma \leq 1$, $\tau \in \mathbb{C} \setminus \{0\}$, $z \in \mathbb{U}$), then for any complex number μ*

$$|a_3 - \mu a_2^2| \leq \frac{B_1|\tau|}{3(1+2\gamma)} \max \left\{ 1, \left| \frac{B_2}{B_1} - \frac{3\tau\mu B_1(1+2\gamma)}{4(1+\gamma)^2} \right| \right\}. \quad (2.1)$$

The result is sharp.

Proof. If $f(z) \in R_Y^\tau(\phi)$, then there exists a Schwarz function $w(z)$ analytic in $\mathbb{U}$ with $w(0) = 0$ and $|w(z)| < 1$ in $\mathbb{U}$ such that

$$1 + \frac{1}{\tau} (f'(z) + \gamma z f''(z) - 1) = \phi(w(z)) \quad (z \in \mathbb{U}). \quad (2.2)$$

Define the function $p_1(z)$ by

$$p_1(z) = \frac{1+w(z)}{1-w(z)} = 1 + c_1z + c_2z^2 + \dots. \quad (2.3)$$

Since $w(z)$ is a Schwarz function, we see that $\Re p_1(z) > 0$ and $p_1(0) = 1$. Define the function $p(z)$ by

$$p(z) = 1 + \frac{1}{\tau} (f'(z) + \gamma z f''(z) - 1) = 1 + b_1 z + b_2 z^2 + \dots \quad (2.4)$$

In view of (2.2), (2.3), (2.4), we have

$$\begin{aligned} p(z) &= \phi \left(\frac{p_1(z) - 1}{p_1(z) + 1} \right) = \phi \left(\frac{c_1 z + c_2 z^2 + \dots}{2 + c_1 z + c_2 z^2 + \dots} \right) \\ &= \phi \left(\frac{1}{2} c_1 z + \frac{1}{2} \left(c_2 - \frac{c_1^2}{2} \right) z^2 + \dots \right) \\ &= 1 + B_1 \frac{1}{2} c_1 z + B_1 \frac{1}{2} \left(c_2 - \frac{c_1^2}{2} \right) z^2 + B_2 \frac{1}{4} c_1^2 z^2 + \dots \end{aligned} \quad (2.5)$$

Thus,

$$b_1 = \frac{1}{2} B_1 c_1; \quad b_2 = \frac{1}{2} B_1 \left(c_2 - \frac{c_1^2}{2} \right) + \frac{1}{4} B_2 c_1^2. \quad (2.6)$$

From (2.4), we obtain

$$a_2 = \frac{B_1 c_1 \tau}{4(1 + \gamma)}; \quad a_3 = \frac{\tau}{6(1 + 2\gamma)} \left[B_1 \left(c_2 - \frac{c_1^2}{2} \right) + \frac{1}{2} B_2 c_1^2 \right]. \quad (2.7)$$

Therefore, we have

$$a_3 - \mu a_2^2 = \frac{B_1 \tau}{6(1 + 2\gamma)} \left(c_2 - \nu c_1^2 \right), \quad (2.8)$$

where

$$\nu = \frac{1}{2} \left(1 - \frac{B_2}{B_1} + \frac{3\tau\mu B_1(1 + 2\gamma)}{4(1 + \gamma)^2} \right). \quad (2.9)$$

Our result now is followed by an application of Lemma 1.2. Also, by the application of Lemma 1.2 equality in (2.1) is obtained when

$$p_1(z) = \frac{1 + z^2}{1 - z^2} \quad \text{or} \quad p_1(z) = \frac{1 + z}{1 - z} \quad (2.10)$$

but

$$p(z) = 1 + \frac{1}{\tau} (f'(z) + \gamma z f''(z) - 1) = \phi\left(\frac{p_1(z)-1}{p_1(z)+1}\right). \quad (2.11)$$

Putting value of $p_1(z)$ we get the desired results. $\square$

For class $R_\gamma^\tau(A, B)$,

$$\begin{aligned} \phi(z) &= \frac{1 + Az}{1 + Bz} = (1 + Az)(1 + Bz)^{-1} \quad (z \in \mathbb{U}) \\ &= 1 + (A - B)z - (AB - B^2)z^2 + \dots \end{aligned} \quad (2.12)$$

Thus, putting $B_1 = A - B$ and $B_2 = -B(A - B)$ in Theorem 2.1, we get the following corollary.

Corollary 2.2. *If $f(z)$ given by (1.1) belongs to $R_\gamma^\tau(A, B)$, then*

$$\left| a_3 - \mu a_2^2 \right| \leq \frac{(A - B)|\tau|}{3(1 + 2\gamma)} \max \left\{ 1, \left| B + \frac{3\tau\mu(A - B)(1 + 2\gamma)}{4(1 + \gamma)^2} \right| \right\}. \quad (2.13)$$

References

- [1] P. L. Duren, *Univalent Functions*, vol. 259 of *Fundamental Principles of Mathematical Sciences*, Springer, Berlin, Germany, 1983.
- [2] M. Fekete and G. Szegő, "Eine bemerkung uber ungerade schlichten funktionene," *Journal of London Mathematical Society*, vol. 8, pp. 85–89, 1993.
- [3] W. Koepf, "On the Fekete-Szegő problem for close-to-convex functions," *Proceedings of the American Mathematical Society*, vol. 101, no. 1, pp. 89–95, 1987.
- [4] W. Koepf, "On the Fekete-Szegő problem for close-to-convex functions. II," *Archiv der Mathematik*, vol. 49, no. 5, pp. 420–433, 1987.
- [5] R. R. London, "Fekete-Szegő inequalities for close-to-convex functions," *Proceedings of the American Mathematical Society*, vol. 117, no. 4, pp. 947–950, 1993.
- [6] W. C. Ma and D. Minda, "A unified treatment of some special classes of univalent functions," in *Proceedings of the Conference on Complex Analysis*, Z. Li, F. Ren, L. Lang, and S. Zhang, Eds., pp. 157–169, International Press, Cambridge, Mass, USA.
- [7] A. Swaminathan, "Sufficient conditions for hypergeometric functions to be in a certain class of analytic functions," *Computers & Mathematics with Applications*, vol. 59, no. 4, pp. 1578–1583, 2010.
- [8] S. Ponnusamy and F. Ronning, "Integral transforms of a class of analytic functions," *Complex Variables and Elliptic Equations*, vol. 53, no. 5, pp. 423–434, 2008.
- [9] S. Ponnusamy, "Neighborhoods and Carathéodory functions," *Journal of Analysis*, vol. 4, pp. 41–51, 1996.
- [10] J. L. Li, "On some classes of analytic functions," *Mathematica Japonica*, vol. 40, no. 3, pp. 523–529, 1994.
- [11] A. Swaminathan, "Certain sufficiency conditions on Gaussian hypergeometric functions," *Journal of Inequalities in Pure and Applied Mathematics*, vol. 5, no. 4, article 83, 6 pages, 2004.
- [12] F. R. Keogh and E. P. Merkes, "A coefficient inequality for certain classes of analytic functions," *Proceedings of the American Mathematical Society*, vol. 20, pp. 8–12, 1969.
- [13] R. J. Libera and E. J. Zlotkiewicz, "Coefficient bounds for the inverse of a function with derivative in ρ ," *Proceedings of the American Mathematical Society*, vol. 87, no. 2, pp. 251–257, 1983.