Research Article

A Construction of Mirror Q -Algebras

Keum Sook So

Department of Mathematics, Hallym University, Chuncheon 200-702, Republic of Korea

Correspondence should be addressed to Keum Sook So, kssso@hallym.ac.kr

Received 5 November 2010; Accepted 29 November 2010

Academic Editor: Hee Sik Kim

Copyright © 2011 Keum Sook So. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We investigate how to construct mirror Q -algebras of a Q -algebra, and we obtain the necessary conditions for $M(X)$ to be a Q -algebra.

1. Introduction

Imai and Iséki introduced two classes of abstract algebras: BCK -algebras and BCI -algebras [1, 2]. It is known that the class of BCK -algebras is a proper subclass of the class of BCI -algebras. We refer the reader for useful textbooks for BCK/BCI -algebra to [3–5]. Neggers et al. [6] introduced the notion of Q -algebras which is a generalization of $BCK/BCI/BCH$ -algebras, obtained several properties, and discussed quadratic Q -algebras. Ahn and Kim [7] introduced the notion of QS -algebras, and Ahn et al. [8] studied positive implicativeity in Q -algebras and discussed some relations between $R - (L-)$ maps and positive implicativeity. Neggers and Kim introduced the notion of d -algebras which is another useful generalization of BCK -algebras and then investigated several relations between d -algebras and BCK -algebras as well as several other relations between d -algebras and oriented digraphs [9]. After that some further aspects were studied [10–13]. Allen et al. [14] introduced the notion of mirror image of given algebras and obtained some interesting properties: a mirror algebra of a d -algebra is also a d -algebra, and a mirror algebra of an implicative BCK -algebra is a left L -up algebra.

In this paper we introduce the notion of mirror algebras to Q -algebras, and we investigate how to construct mirror Q -algebras from a Q -algebra; and we also obtain the necessary conditions for $M(X)$ to be a Q -algebra.

2. Q -Algebras and Related Algebras

A Q -algebra [6] is a nonempty set X with a constant 0 and a binary operation “ $*$ ” satisfying the following axioms:

- (I) $x * x = 0$,
- (II) $0 * x = 0$,
- (III) $(x * y) * z = (x * z) * y$ for all $x, y, z \in X$.

For brevity we also call X a Q -algebra. In X we can define a binary relation “ $\leq$ ” by $x \leq y$ if and only if $x * y = 0$.

Example 2.1 (see [6]). Let $X : \{0, 1, 2, 3\}$ be a set with the following table:

$*$	0	1	2	3
0	0	0	0	0
1	0	0	3	2
2	2	0	0	0
3	3	3	3	0

(2.1)

Then $(X, *, 0)$ is a Q -algebra, which is not a $BCK/BCI/BCH$ -algebra.

Ahn and Kim [7] introduced the notion of QS -algebras. They showed that the G -part of an associative QS -algebra is a group in which every element is an involution. A Q -algebra X is said to be a QS -algebra if it satisfies the following condition:

- (IV) $(x * y) * (x * z) = z * y$, for all $x, y, z \in X$.

Proposition 2.2 (see [6]). *If $(X, *, 0)$ is a Q -algebra, then*

- (V) $(x * (x * y)) * y = 0$, for all $x, y \in X$.

It was proved that every BCH -algebra is a Q -algebra and every Q -algebra satisfying some additional conditions is a BCI -algebra.

Neggers and Kim [15] introduced the notion of B -algebras which is related to several classes of algebras of interest such as $BCH/BCI/BCK$ -algebras and which seems to have rather nice properties without being excessively complicated otherwise. And they demonstrated some interesting connections between B -algebras and groups.

Example 2.3. Let $X := \{0, 1, 2, \dots, \omega\}$ be a set. Define a binary operation “ $*$ ” on X by

$$x * y := \begin{cases} 0, & x \leq y, \\ \omega, & y < x, x \neq 0, \\ x, & y < x, y = 0. \end{cases} \quad (2.2)$$

Then $(X, *, 0)$ is a Q -algebra, but not a B -algebra, since $(3 * \omega) * 0 = 0$, $3 * (0 * (0 * \omega)) = 3$.

Example 2.4. Let $X := \{0, 1, \dots, 5\}$ be a set with the following table:

$*$	0	1	2	3	4	5
0	0	2	1	3	4	5
1	1	0	2	4	5	3
2	2	1	0	5	3	4
3	3	4	5	0	2	1
4	4	5	3	1	0	2
5	5	3	4	2	1	0

(2.3)

Then $(X, *, 0)$ is a B -algebra, but not a Q -algebra, since $(5 * 3) * 1 = 1$, $(5 * 1) * 3 = 0$.

Example 2.5. Let X be the set of all real numbers except for a negative integer $-n$. Define a binary operation $*$ on X by

$$x * y := \frac{n(x - y)}{n + y} \quad (2.4)$$

for any $x, y \in X$. Then $(X, *, 0)$ is both a Q -algebra and B -algebra.

If we consider several families of abstract algebras including the well-known BCK -algebras and several larger classes including the class of d -algebras which is a generalization of BCK -algebras, then it is usually difficult and often impossible to obtain a complementation operation and the associated “de Morgan’s laws.” In the sense of this point of view it is natural to construct a “mirror image” of a given algebra which when adjoined to the original algebra permits a natural complementation to take place. The class of BCK -algebras is not closed under this operation but the class of d -algebras is, thus explaining why it may be better to work with this class rather than the class of BCK -algebras. Allen et al. [14] introduced the notion of mirror algebras of a given algebra.

Let $(X, *, 0)$ be an algebra. Let $M(X) := X \times \{0, 1\}$, and define a binary operation “ $*$ ” on $M(X)$ as follows:

$$\begin{aligned}
 (x, 0) * (y, 0) &:= (x * y, 0), \\
 (x, 1) * (y, 1) &:= (y * x, 0), \\
 (x, 0) * (y, 1) &:= (x * (x * y), 0), \\
 (x, 1) * (y, 0) &:= \begin{cases} (y, 1) & \text{when } x * y = 0, \\ (x, 1) & \text{when } x * y \neq 0. \end{cases}
 \end{aligned} \quad (2.5)$$

Then we say that $M(X) := (M(X), *, (0, 0))$ is a *left mirror algebra* of the algebra $(X, *, 0)$. Similarly, if we define

$$(x, *) * (y, 1) := (y * (y * x), 0), \quad (2.6)$$

then $M(X) := (M(X), *, (0, 0))$ is a *right mirror algebra* of the algebra $(X, *, 0)$.

It was shown [14] that the mirror algebra of a d (resp., d -BH)-algebra is also a d (resp., d -BH)-algebra, but the mirror algebra of a BCK-algebra need not be a BCK-algebra.

3. A Construction of Mirror Q -Algebras

In [14] Allen et al. defined (left, right) mirror algebras of an algebra, but it is not known how to construct mirror algebras of any given algebra. In this paper, we investigate a construction of a mirror algebra in Q -algebras.

Let $(X, *, 0)$ be a Q -algebra, and let $M(X) := X \times \{0, 1\}$. Define a binary operation " $\oplus$ " on $M(X)$ by

$$(M1) \quad (x, 0) \oplus (y, 0) = (x * y, 0),$$

$$(M2) \quad (x, 1) \oplus (y, 1) = (y * x, 0),$$

$$(M3) \quad (x, 0) \oplus (y, 1) = (\alpha(x, y), 0),$$

$$(M4) \quad (x, 1) \oplus (y, 0) = (\beta(x, y), 1),$$

where $\alpha, \beta : X \times X \rightarrow X$ are mappings.

Consider condition (I). If we let $x = y$ in (1) and (2), then (I) holds trivially. Consider condition (II). For any $(x, 0) \in M(X)$, we have $(x, 0) \oplus (0, 0) = (x * 0, 0) = (x, 0)$. For any $(x, 1) \in M(X)$, we have $(x, 1) = (x, 1) \oplus (0, 0) = (\beta(x, 0), 1)$, which shows that the required condition is $\beta(x, 0) = x$. Consider condition (III). There are 8 cases to check that condition (III) holds.

Case 1 $((x, 0), (y, 0), (z, 0))$. It holds trivially.

Case 2 $((x, 0), (y, 1), (z, 0))$. Since $((x, 0) \oplus (y, 1)) \oplus (z, 0) = (\alpha(x, y), 0) \oplus (z, 0) = (\alpha(x, y) * z, 0)$ and $((x, 0) \oplus (z, 0)) \oplus (y, 1) = (x * z, 0) \oplus (y, 1) = (\alpha(x * z, y), 0)$, we obtain the requirement that $\alpha(x, y) * z = \alpha(x * z, y)$.

Case 3 $((x, 0), (y, 0), (z, 1))$. It is the same as Case 2.

Case 4 $((x, 0), (y, 1), (z, 1))$. Since $((x, 0) \oplus (y, 1)) \oplus (z, 1) = (\alpha(x, y), 0) \oplus (z, 1) = (\alpha(\alpha(x, y), z), 0)$ and $((x, 0) \oplus (z, 1)) \oplus (y, 1) = (\alpha(x, z), 0) \oplus (y, 1) = (\alpha(\alpha(x, z), y), 0)$, we obtain the requirement that $\alpha(\alpha(x, y), z) = \alpha(\alpha(x, z), y)$.

Case 5 $((x, 1), (y, 0), (z, 0))$. Since $((x, 1) \oplus (y, 0)) \oplus (z, 0) = (\beta(x, y), 1) \oplus (z, 0) = (\beta(\beta(x, y), z), 0)$ and $((x, 1) \oplus (z, 0)) \oplus (y, 0) = (\beta(x, z), 1) \oplus (y, 0) = (\beta(\beta(x, z), y), 0)$, we obtain the requirement that $\beta(\beta(x, y), z) = \beta(\beta(x, z), y)$.

Case 6 $((x, 1), (y, 0), (z, 1))$. Since $((x, 1) \oplus (y, 0)) \oplus (z, 1) = (\beta(x, y), 1) \oplus (z, 1) = (z * \beta(x, y), 0)$ and $((x, 1) \oplus (z, 1)) \oplus (y, 0) = (z * x, 0) \oplus (y, 0) = ((z * x) * y, 0)$, we obtain the requirement that $z * \beta(x, y) = (z * x) * y$.

Case 7 $((x, 1), (y, 1), (z, 0))$. It is the same as Case 6.

Case 8 $((x, 1), (y, 1), (z, 1))$. Since $((x, 1) \oplus (y, 1)) \oplus (z, 1) = (\beta(x, y), 0) \oplus (z, 1) = (\alpha(\beta(x, y), z), 0)$ and $((x, 1) \oplus (z, 1)) \oplus (y, 1) = (\alpha(\beta(x, z), y), 0)$ by exchanging y with z , we obtain the requirement that $\alpha(\beta(x, y), z) = \alpha(\beta(x, z), y)$. If we summarize this discussion, we obtain the following theorem.

Theorem 3.1. Let $(X, *, 0)$ be a Q -algebra, and let $M(X) := X \times \{0, 1\}$ be a set with a binary operation “ $\oplus$ ” on $M(X)$ with (M1) $\sim$ (M4). Then the necessary conditions for $(M(X), \oplus, (0, 0))$ to be a Q -algebra are the following:

- (i) $\beta(x, 0) = x$,
- (ii) $\alpha(\alpha(x, y), z) = \alpha(\alpha(x, z), y)$,
- (iii) $\alpha(x, y) * z = \alpha(x * z, y)$,
- (iv) $\beta(\beta(x, y), z) = \beta(\beta(x, z), y)$,
- (v) $z * \beta(x, y) = (z * x) * y$,
- (vi) $\alpha(\beta(x, y), z) = \alpha(\beta(x, z), y)$

for any $x, y, z \in X$.

Remark 3.2. By condition (M1), if we identify $(x, 0) \equiv x$ for any $x \in X$, then X is a subalgebra of $M(X)$. By applying Theorem 3.1, we obtain many (mirror) Q -algebras: $X \subseteq M(X) \subseteq M(M(X)) = M^2(X) \subseteq M^3(X) \subseteq M^4(X) \subseteq \dots$.

Example 3.3. Let Z be the set of all integers. Then $(Z, -, 0)$ is a Q -algebra where “ $-$ ” is the usual subtraction in Z . If we define mappings $\alpha, \beta : Z \times Z \rightarrow Z$ by $\alpha(x, y) = \beta(x, y) = x + y$ for any $x, y \in Z$, then the mirror algebra $(M(Z), \oplus, (0, 0))$ is also a Q -algebra, that is, $(x, 0) \oplus (y, 0) = (x + y, 0)$, $(x, 1) \oplus (y, 1) = (y - x, 0)$, $(x, 0) \oplus (y, 1) = (x + y, 0)$, and $(x, 1) \oplus (y, 0) = (x + y, 1)$.

Example 3.4. Let $X := \{0, 1\}$ be a set with the following table:

$$\begin{array}{c|cc} * & 0 & 1 \\ \hline 0 & 0 & 0 \\ 1 & 1 & 0 \end{array} \quad (3.1)$$

Then $(X, *, 0)$ is a Q -algebra. Using the same method we obtain its mirror algebra as follows: $M(X) = \{0, \alpha, \beta, \gamma\}$ with the following table:

$$\begin{array}{c|cccc} \oplus & 0 & \alpha & \beta & \gamma \\ \hline 0 & 0 & 0 & 0 & 0 \\ \alpha & \alpha & 0 & \alpha & 0 \\ \beta & \beta & \beta & 0 & 0 \\ \gamma & \gamma & \beta & \alpha & 0 \end{array} \quad (3.2)$$

where $0 := (0, 0)$, $\alpha := (0, 1)$, $\beta := (1, 0)$, and $\gamma := (1, 1)$. It is easy to see that $(M(X), \oplus, 0)$ is a Q -algebra.

Problems

- (1) Find necessary conditions for $M(X)$ to be a QS -algebra if $(X, *, 0)$ is a QS -algebra.
- (2) Given a homomorphism $f : X \rightarrow Y$ of Q -algebras, construct a homomorphism $\hat{f} : M(X) \rightarrow M(Y)$ of Q -algebras which is an extension of f .
- (3) Given Q -algebras X, Y , are the mirror algebras $M(M(X \times Y))$ and $M(X) \times M(Y)$ isomorphic?

References

- [1] K. Iséki, "On BCI-algebras," *Mathematics Seminar Notes*, vol. 8, no. 1, pp. 125–130, 1980.
- [2] K. Iséki and S. Tanaka, "An introduction to the theory of BCK-algebras," *Mathematica Japonica*, vol. 23, no. 1, pp. 1–26, 1978.
- [3] A. Iorgulescu, *Algebras of Logic as BCK Algebras*, Editura ASE, Bucharest, Romania, 2008.
- [4] J. Meng and Y. B. Jun, *BCK-Algebras*, Kyung Moon Sa, Seoul, Republic of Korea, 1994.
- [5] H. Yisheng, *BCI-Algebras*, Science Press, Beijing, China, 2006.
- [6] J. Neggers, S. S. Ahn, and H. S. Kim, "On Q -algebras," *International Journal of Mathematics and Mathematical Sciences*, vol. 27, no. 12, pp. 749–757, 2001.
- [7] S. S. Ahn and H. S. Kim, "On QS -algebras," *The Journal of Chungcheong Mathematical Society*, vol. 12, pp. 33–41, 1999.
- [8] S. S. Ahn, H. S. Kim, and H. D. Lee, " R -maps and L -maps in Q -algebras," *International Journal of Pure and Applied Mathematics*, vol. 12, no. 4, pp. 419–425, 2004.
- [9] J. Neggers and H. S. Kim, "On d -algebras," *Mathematica Slovaca*, vol. 49, no. 1, pp. 19–26, 1999.
- [10] P. J. Allen, H. S. Kim, and J. Neggers, "Companion d -algebras," *Mathematica Slovaca*, vol. 57, no. 2, pp. 93–106, 2007.
- [11] P. J. Allen, H. S. Kim, and J. Neggers, "Deformations of d/BCK -algebras," to appear in *Bulletin of the Korean Mathematical Society*.
- [12] J. Neggers, A. Dvure enskij, and H. S. Kim, "On d -fuzzy functions in d -algebras," *Foundations of Physics*, vol. 30, no. 10, pp. 1807–1816, 2000.
- [13] J. Neggers, Y. B. Jun, and H. S. Kim, "On d -ideals in d -algebras," *Mathematica Slovaca*, vol. 49, no. 3, pp. 243–251, 1999.
- [14] P. J. Allen, J. Neggers, and H. S. Kim, " L -up and mirror algebras," *Scientiae Mathematicae Japonicae*, vol. 59, no. 3, pp. 605–612, 2004.
- [15] J. Neggers and H. S. Kim, "On B -algebras," *Matematichki Vesnik*, vol. 54, no. 1-2, pp. 21–29, 2002.