Research Article

Fuzzy Stability of a Quadratic-Additive Functional Equation

Sun Sook Jin and Yang Hi Lee

*Department of Mathematics Education, Gongju National University of Education,
Gongju 314-711, Republic of Korea*

Correspondence should be addressed to Yang Hi Lee, yanghi2@hanmail.net

Received 27 April 2011; Accepted 25 August 2011

Academic Editor: Naseer Shahzad

Copyright © 2011 S. S. Jin and Y. H. Lee. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We investigate a fuzzy version of stability for the functional equation $f(x + y + z + w) + 2f(x) + 2f(y) + 2f(z) + 2f(w) - f(x + y) - f(x + z) - f(x + w) - f(y + z) - f(y + w) - f(z + w) = 0$.

1. Introduction

A classical question in the theory of functional equations is “when is it true that a mapping, which approximately satisfies a functional equation, must be somehow close to an exact solution of the equation?” Such a problem, called *a stability problem of the functional equation*, was formulated by Ulam [1] in 1940. In the next year, Hyers [2] gave a partial solution of Ulam’s problem for the case of approximate additive mappings. Subsequently, his result was generalized by Aoki [3] for additive mappings, and by Rassias [4] for linear mappings, to considering the stability problem with the unbounded Cauchy differences. During the last decades, the stability problems of functional equations have been extensively investigated by a number of mathematicians, see [5–16].

In 1984, Katsaras [17] defined a fuzzy norm on a linear space to construct a fuzzy structure on the space. Since then, some mathematicians have introduced several types of fuzzy norm in different points of view. In particular, Bag and Samanta [18], following Cheng and Mordeson [19], gave an idea of a fuzzy norm in such a manner that the corresponding fuzzy metric is of the Kramosil and Michálek type [20]. In 2008, Mirmostafaei and Moslehian [21] introduced for the first time the notion of fuzzy Hyers-Ulam-Rassias stability. They obtained a fuzzy version of stability for *the Cauchy functional equation*

$$f(x + y) - f(x) - f(y) = 0 \quad (1.1)$$

whose solution is called *an additive mapping*. In the same year, they [22] proved a fuzzy version of stability for *the quadratic functional equation*

$$f(x+y) + f(x-y) - 2f(x) - 2f(y) = 0 \quad (1.2)$$

whose solution is called *a quadratic mapping*. Now we consider the *quadratic-additive functional equation*

$$\begin{aligned} f(x+y+z+w) + 2f(x) + 2f(y) + 2f(z) + 2f(w) - f(x+y) \\ - f(x+z) - f(x+w) - f(y+z) - f(y+w) - f(z+w) = 0 \end{aligned} \quad (1.3)$$

whose solution is called *a quadratic-additive mapping*. In [23], Chang et al. obtained a stability of the quadratic-additive functional equation by taking and composing an additive mapping A and a quadratic mapping Q to prove the existence of a quadratic-additive mapping F which is close to the given mapping f . In their processing, A is approximate to the odd part $f((x) - f(-x))/2$ of f and Q is close to the even part $(f(x) + f(-x))/2$ of it, respectively.

In this paper, we get a general stability result of the quadratic-additive functional equation in the fuzzy normed linear space. To do it, we introduce a Cauchy sequence $\{J_n f(x)\}$ starting from a given mapping f , which converges to the desired mapping F in the fuzzy sense. As we mentioned before, in previous studies of stability problem of (1.3), Chang et al. attempted to get stability theorems by handling the odd and even part of f , respectively. According to our proposal in this paper, we can take the desired approximate solution F at once.

2. Fuzzy Stability of the Quadratic-Additive Functional Equation

We use the definition of a fuzzy normed space given in [18] to exhibit a reasonable fuzzy version of stability for the quadratic-additive functional equation in the fuzzy normed linear space.

Definition 2.1 (see [18]). Let X be a real linear space. A mapping $N : X \times \mathbb{R} \rightarrow [0, 1]$ (the so-called fuzzy subset) is said to be *a fuzzy norm on X* if for all $x, y \in X$ and all $s, t \in \mathbb{R}$,

- (N1) $N(x, c) = 0$ for $c \leq 0$,
- (N2) $x = 0$ if and only if $N(x, c) = 1$ for all $c > 0$,
- (N3) $N(cx, t) = N(x, t/|c|)$ if $c \neq 0$,
- (N4) $N(x+y, s+t) \geq \min\{N(x, s), N(y, t)\}$,
- (N5) $N(x, \cdot)$ is a nondecreasing mapping on $\mathbb{R}$ and $\lim_{t \rightarrow \infty} N(x, t) = 1$.

The pair (X, N) is called *a fuzzy normed linear space*. Let (X, N) be a fuzzy normed linear space. Let $\{x_n\}$ be a sequence in X . Then $\{x_n\}$ is said to be convergent if there exists $x \in X$ such that $\lim_{n \rightarrow \infty} N(x_n - x, t) = 1$ for all $t > 0$. In this case, x is called *the limit of the sequence $\{x_n\}$* , and we denote it by $N - \lim_{n \rightarrow \infty} x_n = x$. A sequence $\{x_n\}$ in X is called *Cauchy* if for each $\varepsilon > 0$ and each $t > 0$ there exists n_0 such that for all $n \geq n_0$ and all $p > 0$ we have $N(x_{n+p} - x_n, t) > 1 - \varepsilon$. It is known that every convergent sequence in a fuzzy normed space

is Cauchy. If each Cauchy sequence is convergent, then the fuzzy norm is said to be *complete* and the fuzzy normed space is called a *fuzzy Banach space*.

Let (X, N) be a fuzzy normed space and (Y, N') a fuzzy Banach space. For a given mapping $f : X \rightarrow Y$, we use the abbreviation

$$Df(x, y, z, w) := f(x + y + z + w) + 2f(x) + 2f(y) + 2f(z) + 2f(w) - f(x + y) - f(x + z) - f(x + w) - f(y + z) - f(y + w) - f(z + w) \quad (2.1)$$

for all $x, y, z, w \in X$. For given $q > 0$, the mapping f is called a *fuzzy q -almost quadratic-additive mapping* if

$$N'(Df(x, y, z, w), s + t + u + v) \geq \min\{N(x, s^q), N(y, t^q), N(z, u^q), N(w, v^q)\} \quad (2.2)$$

for all $x, y, z, w \in X$ and $s, t, u, v \in (0, \infty)$. Now we get the general stability result in the fuzzy normed linear setting.

Theorem 2.2. *Let q be a positive real number with $q \neq 1/2, 1$. And let f be a fuzzy q -almost quadratic-additive mapping from a fuzzy normed space (X, N) into a fuzzy Banach space (Y, N') . Then there is a unique quadratic-additive mapping $F : X \rightarrow Y$ such that*

$$N'(F(x) - f(x), t) \geq \sup_{t' < t} N\left(x, \frac{(|4 - 2^p||2 - 2^p|t')^q}{(2(|4 - 2^p| + |2 - 2^p|))^q}\right) \quad (2.3)$$

for all $x \in X$ and $t > 0$, where $p = 1/q$.

Proof. It follows from (2.2) and (N4) that

$$N'(f(0), t) = N'(Df(0, 0, 0, 0), 3t) \geq N\left(0, \left(\frac{3t}{4}\right)^q\right) = 1 \quad (2.4)$$

for all $t \in (0, \infty)$. By (N2), we have $f(0) = 0$. We will prove the theorem in three cases: $q > 1$, $1/2 < q < 1$, and $0 < q < 1/2$.

Case 1. Let $q > 1$ and let $J_n f : X \rightarrow Y$ be a mapping defined by

$$J_n f(x) = \frac{f(2^n x) + f(-2^n x)}{2 \cdot 4^n} + \frac{f(2^n x) - f(-2^n x)}{2^{n+1}} \quad (2.5)$$

for all $x \in X$. Notice that $J_0 f(x) = f(x)$ and

$$J_j f(x) - J_{j+1} f(x) = \frac{Df(2^j x, -2^j x, 2^j x, -2^j x)}{2 \cdot 4^{j+1}} + \frac{Df(2^j x, 2^j x, 2^j x, -2^j x)}{2^{j+3}} - \frac{Df(-2^j x, -2^j x, -2^j x, 2^j x)}{2^{j+3}} \quad (2.6)$$

for all $x \in X$ and $j \geq 0$. Together with (N3), (N4), and (2.2), this equation implies that if $n + m > m \geq 0$, then

$$\begin{aligned}
 & N' \left(J_m f(x) - J_{n+m} f(x), \sum_{j=m}^{n+m-1} \left(\frac{1}{2} \left(\frac{2^p}{4} \right)^j + \left(\frac{2^p}{2} \right)^j \right) t^p \right) \\
 & \geq N' \left(\sum_{j=m}^{n+m-1} (J_j f(x) - J_{j+1} f(x)), \sum_{j=m}^{n+m-1} \frac{(1 + 2^{j+1}) 2^{jp}}{2 \cdot 4^j} t^p \right) \\
 & \geq \min \bigcup_{j=m}^{n+m-1} \left\{ N' \left(J_j f(x) - J_{j+1} f(x), \frac{(1 + 2^{j+1}) 2^{jp}}{2 \cdot 4^j} t^p \right) \right\} \\
 & \geq \min \bigcup_{j=m}^{n+m-1} \left\{ \min \left\{ N' \left(\frac{Df(2^j x, -2^j x, 2^j x, -2^j x)}{2 \cdot 4^{j+1}}, \frac{2^{jp} t^p}{2 \cdot 4^j} \right), \right. \right. \\
 & \quad N' \left(\frac{Df(2^j x, 2^j x, 2^j x, -2^j x)}{2^{j+3}}, \frac{2^{jp} t^p}{2^{j+1}} \right), \\
 & \quad \left. \left. N' \left(-\frac{Df(-2^j x, -2^j x, -2^j x, 2^j x)}{2^{j+3}}, \frac{2^{jp} t^p}{2^{j+1}} \right) \right\} \right\} \\
 & \geq \min \bigcup_{j=m}^{n+m-1} \left\{ N(2^j x, 2^j t) \right\} = N(x, t)
 \end{aligned} \tag{2.7}$$

for all $x \in X$ and $t > 0$. Let $\varepsilon > 0$ be given. Since $\lim_{t \rightarrow \infty} N(x, t) = 1$, there is $t_0 > 0$ such that

$$N(x, t_0) \geq 1 - \varepsilon. \tag{2.8}$$

We observe that for some $\tilde{t} > t_0$, the series $\sum_{j=0}^{\infty} (1 + 2^{j+1}) 2^{jp} / 2 \cdot 4^j \tilde{t}^p$ converges for $p = (1/q) < 1$. It guarantees that, for an arbitrary given $c > 0$, there exists $n_0 \geq 0$ such that

$$\sum_{j=m}^{n+m-1} \frac{(1 + 2^{j+1}) 2^{jp}}{2 \cdot 4^j} \tilde{t}^p < c \tag{2.9}$$

for each $m \geq n_0$ and $n > 0$. By (N5) and (2.7), we have

$$\begin{aligned}
 N'(J_m f(x) - J_{n+m} f(x), c) & \geq N' \left(J_m f(x) - J_{n+m} f(x), \sum_{j=m}^{n+m-1} \frac{(1 + 2^{j+1}) 2^{jp}}{2 \cdot 4^j} \tilde{t}^p \right) \\
 & \geq N(x, \tilde{t}) \geq N(x, t_0) \geq 1 - \varepsilon
 \end{aligned} \tag{2.10}$$

for all $x \in X$. Hence $\{J_n f(x)\}$ is a Cauchy sequence in the fuzzy Banach space (Y, N') , and so we can define a mapping $F : X \rightarrow Y$ by

$$F(x) := N' - \lim_{n \rightarrow \infty} J_n f(x) \quad (2.11)$$

for all $x \in X$. Moreover, if we put $m = 0$ in (2.7), we have

$$N'(f(x) - J_n f(x), t) \geq N\left(x, \frac{t^q}{\left(\sum_{j=0}^{n-1} \left((1/2)(2^p/4)^j + (2^p/2)^j\right)\right)^q}\right) \quad (2.12)$$

for all $x \in X$. Next we will show that F is a desired quadratic-additive mapping. Using (N4), we have

$$\begin{aligned} & N'(DF(x, y, z, w), t) \\ & \geq \min \left\{ N'\left(F(x + y + z + w) - J_n f(x + y + z + w), \frac{t}{30}\right), \right. \\ & \quad N'\left(2F(x) - 2J_n f(x), \frac{t}{15}\right), N'\left(2F(y) - 2J_n f(y), \frac{t}{15}\right), \\ & \quad N'\left(2F(z) - 2J_n f(z), \frac{t}{15}\right), N'\left(2F(w) - 2J_n f(w), \frac{t}{15}\right), \\ & \quad N'\left(F(x + y) - J_n f(x + y), \frac{t}{30}\right), N'\left(F(x + z) - J_n f(x + z), \frac{t}{30}\right), \\ & \quad N'\left(F(x + w) - J_n f(x + w), \frac{t}{30}\right), N'\left(F(y + z) - J_n f(y + z), \frac{t}{30}\right), \\ & \quad N'\left(F(y + w) - J_n f(y + w), \frac{t}{30}\right), N'\left(F(z + w) - J_n f(z + w), \frac{t}{30}\right), \\ & \quad \left. N'\left(DJ_n f(x, y, z, w), \frac{t}{2}\right) \right\} \quad (2.13) \end{aligned}$$

for all $x, y, z, w \in X$ and $n \in \mathbb{N}$. The first eleven terms on the right-hand side of (2.13) tend to 1 as $n \rightarrow \infty$ by the definition of F and (N2), and the last term holds:

$$\begin{aligned} N'\left(DJ_n f(x, y, z, w), \frac{t}{2}\right) & \geq \min \left\{ N'\left(\frac{Df(2^n x, 2^n y, 2^n z, 2^n w)}{2 \cdot 4^n}, \frac{t}{8}\right), \right. \\ & \quad N'\left(\frac{Df(-2^n x, -2^n y, -2^n z, -2^n w)}{2 \cdot 4^n}, \frac{t}{8}\right), \\ & \quad N'\left(\frac{Df(2^n x, 2^n y, 2^n z, 2^n w)}{2 \cdot 2^n}, \frac{t}{8}\right), \\ & \quad \left. N'\left(\frac{Df(-2^n x, -2^n y, -2^n z, -2^n w)}{2 \cdot 2^n}, \frac{t}{8}\right) \right\} \quad (2.14) \end{aligned}$$

for all $x, y, z, w \in X$. By (N3) and (2.2), we obtain

$$\begin{aligned}
 N' \left(\frac{Df(\pm 2^n x, \pm 2^n y, \pm 2^n z, \pm 2^n w)}{2 \cdot 4^n}, \frac{t}{8} \right) &= N' \left(Df(\pm 2^n x, \pm 2^n y, \pm 2^n z, \pm 2^n w), \frac{4^n t}{4} \right) \\
 &\geq \min \left\{ N \left(2^n x, \left(\frac{4^n t}{16} \right)^q \right), N \left(2^n y, \left(\frac{4^n t}{16} \right)^q \right), \right. \\
 &\quad \left. N \left(2^n z, \left(\frac{4^n t}{16} \right)^q \right), N \left(2^n w, \left(\frac{4^n t}{16} \right)^q \right) \right\} \\
 &\geq \min \left\{ N \left(x, \frac{2^{(2q-1)n}}{2^{4q}} t^q \right), N \left(y, \frac{2^{(2q-1)n}}{2^{4q}} t^q \right), \right. \\
 &\quad \left. N \left(z, \frac{2^{(2q-1)n}}{2^{4q}} t^q \right), N \left(w, \frac{2^{(2q-1)n}}{2^{4q}} t^q \right) \right\}, \\
 N' \left(\frac{Df(\pm 2^n x, \pm 2^n y, \pm 2^n z, \pm 2^n w)}{2 \cdot 2^n}, \frac{t}{8} \right) &\geq \min \left\{ N \left(x, \frac{2^{(q-1)n}}{2^{4q}} t^q \right), N \left(y, \frac{2^{(q-1)n}}{2^{4q}} t^q \right), \right. \\
 &\quad \left. N \left(z, \frac{2^{(q-1)n}}{2^{4q}} t^q \right), N \left(w, \frac{2^{(q-1)n}}{2^{4q}} t^q \right) \right\} \tag{2.15}
 \end{aligned}$$

for all $x, y, z, w \in X$ and $n \in \mathbb{N}$. Since $q > 1$, together with (N5), we can deduce that the last term of (2.13) also tends to 1 as $n \rightarrow \infty$. It follows from (2.13) that

$$N'(DF(x, y, z, w), t) = 1 \tag{2.16}$$

for each $x, y, z, w \in X$ and $t > 0$. By (N2), this means that $DF(x, y, z, w) = 0$ for all $x, y, z, w \in X$.

Now we approximate the difference between f and F in a fuzzy sense. For an arbitrary fixed $x \in X$ and $t > 0$, choose $0 < \varepsilon < 1$ and $0 < t' < t$. Since F is the limit of $\{J_n f(x)\}$, there is $n \in \mathbb{N}$ such that $N'(F(x) - J_n f(x), t - t') \geq 1 - \varepsilon$. By (2.12), we have

$$\begin{aligned}
 N'(F(x) - f(x), t) &\geq \min \{ N'(F(x) - J_n f(x), t - t'), N'(J_n f(x) - f(x), t') \} \\
 &\geq \min \left\{ 1 - \varepsilon, N \left(x, \frac{t'^q}{\left(\sum_{j=0}^{n-1} \left((1/2)(2^p/4)^j + (2^p/2)^j \right) \right)^q} \right) \right\} \\
 &\geq \min \left\{ 1 - \varepsilon, N \left(x, \frac{(4 - 2^p)^q (2 - 2^p)^q t'^q}{(2(4 - 2^p + 2 - 2^p))^q} \right) \right\} \tag{2.17}
 \end{aligned}$$

since $\sum_{j=0}^{n-1} ((1/2)(2^p/4)^j + (2^p/2)^j) \leq 2(4 - 2^p + 2 - 2^p)/((4 - 2^p)(2 - 2^p))$. Because $0 < \varepsilon < 1$ is arbitrary, we get inequality (2.3) in this case. Finally, to prove the uniqueness of F , let $F' : X \rightarrow Y$ be another quadratic-additive mapping satisfying (2.3). Then by (2.6), we get

$$\begin{aligned} F(x) - J_n F(x) &= \sum_{j=0}^{n-1} (J_j F(x) - J_{j+1} F(x)) = 0, \\ F'(x) - J_n F'(x) &= \sum_{j=0}^{n-1} (J_j F'(x) - J_{j+1} F'(x)) = 0 \end{aligned} \quad (2.18)$$

for all $x \in X$ and $n \in \mathbb{N}$. Together with (N4) and (2.3), this implies that

$$\begin{aligned} &N'(F(x) - F'(x), t) \\ &= N'(J_n F(x) - J_n F'(x), t) \\ &\geq \min \left\{ N' \left(J_n F(x) - J_n f(x), \frac{t}{2} \right), N' \left(J_n f(x) - J_n F'(x), \frac{t}{2} \right) \right\} \\ &\geq \min \left\{ N' \left(\frac{(F - f)(2^n x)}{2 \cdot 4^n}, \frac{t}{8} \right), N' \left(\frac{(f - F')(2^n x)}{2 \cdot 4^n}, \frac{t}{8} \right), \right. \\ &\quad \left. N' \left(\frac{(F - f)(-2^n x)}{2 \cdot 4^n}, \frac{t}{8} \right), N' \left(\frac{(f - F')(-2^n x)}{2 \cdot 4^n}, \frac{t}{8} \right), \right. \\ &\quad \left. N' \left(\frac{(F - f)(2^n x)}{2 \cdot 2^n}, \frac{t}{8} \right), N' \left(\frac{(f - F')(2^n x)}{2 \cdot 2^n}, \frac{t}{8} \right), \right. \\ &\quad \left. N' \left(\frac{(F - f)(-2^n x)}{2 \cdot 2^n}, \frac{t}{8} \right), N' \left(\frac{(f - F')(-2^n x)}{2 \cdot 2^n}, \frac{t}{8} \right) \right\} \\ &\geq \sup_{t' < t} N \left(x, \frac{2^{(q-1)n-2q}(4-2^p)^q(2-2^p)^q t'^q}{(4(3-2^p))^q} \right) \end{aligned} \quad (2.19)$$

for all $x \in X$ and $n \in \mathbb{N}$. Observe that, for $q = (1/p) > 1$, the last term of the above inequality tends to 1 as $n \rightarrow \infty$ by (N5). This implies that $N'(F(x) - F'(x), t) = 1$, and so we get

$$F(x) = F'(x) \quad (2.20)$$

for all $x \in X$ by (N2).

Case 2. Let $(1/2) < q < 1$ and let $J_n f : X \rightarrow Y$ be a mapping defined by

$$J_n f(x) = \frac{1}{2} \left(4^{-n} (f(2^n x) + f(-2^n x)) + 2^n \left(f\left(\frac{x}{2^n}\right) - f\left(-\frac{x}{2^n}\right) \right) \right) \quad (2.21)$$

for all $x \in X$. Then we also have $J_0 f(x) = f(x)$ and

$$\begin{aligned} J_j f(x) - J_{j+1} f(x) &= \frac{Df(2^j x, -2^j x, 2^j x, -2^j x)}{2 \cdot 4^{j+1}} - 2^{j-2} Df\left(\frac{x}{2^{j+1}}, \frac{x}{2^{j+1}}, \frac{x}{2^{j+1}}, -\frac{x}{2^{j+1}}\right) \\ &\quad + 2^{j-2} Df\left(-\frac{x}{2^{j+1}}, -\frac{x}{2^{j+1}}, -\frac{x}{2^{j+1}}, \frac{x}{2^{j+1}}\right) \end{aligned} \quad (2.22)$$

for all $x \in X$ and $j \geq 0$. If $n + m > m \geq 0$, then we have

$$\begin{aligned} &N' \left(J_m f(x) - J_{n+m} f(x), \sum_{j=m}^{n+m-1} \left(\frac{1}{2} \left(\frac{2^p}{4} \right)^j + \left(\frac{2}{2^p} \right)^{j+1} \right) t^p \right) \\ &\geq \min \bigcup_{j=m}^{n+m-1} \left\{ \min \left\{ N' \left(\frac{Df(2^j x, -2^j x, 2^j x, -2^j x)}{2 \cdot 4^{j+1}}, \frac{2^j t^p}{2 \cdot 4^j} \right), \right. \right. \\ &\quad \left. N' \left(-2^{j-2} Df\left(\frac{x}{2^{j+1}}, \frac{x}{2^{j+1}}, \frac{x}{2^{j+1}}, -\frac{x}{2^{j+1}}\right), \frac{2^j t^p}{2^{(j+1)p}} \right), \right. \\ &\quad \left. \left. N' \left(2^{j-2} Df\left(-\frac{x}{2^{j+1}}, -\frac{x}{2^{j+1}}, -\frac{x}{2^{j+1}}, \frac{x}{2^{j+1}}\right), \frac{2^j t^p}{2^{(j+1)p}} \right) \right\} \right\} \\ &\geq \min \bigcup_{j=m}^{n+m-1} \left\{ N(2^j x, 2^j t), N\left(\frac{x}{2^{j+1}}, \frac{t}{2^{j+1}}\right) \right\} = N(x, t) \end{aligned} \quad (2.23)$$

for all $x \in X$ and $t > 0$. In a similar argument following (2.7) of the previous case, we can define the limit $F(x) := N' - \lim_{n \rightarrow \infty} J_n f(x)$ of the Cauchy sequence $\{J_n f(x)\}$ in the Banach fuzzy space Y . Moreover, putting $m = 0$ in the above inequality, we have

$$N'(f(x) - J_n f(x), t) \geq N \left(x, \frac{t^q}{\left(\sum_{j=0}^{n-1} \left((1/2)(2^p/4)^j + (2/2^p)^{j+1} \right) \right)^q} \right) \quad (2.24)$$

for each $x \in X$ and $t > 0$. To prove that F is a quadratic-additive mapping, we have enough to show that the last term of (2.13) in Case 1 tends to 1 as $n \rightarrow \infty$. By (N3) and (2.2), we get

$$\begin{aligned}
 & N' \left(DJ_n f(x, y, z, w), \frac{t}{2} \right) \\
 & \geq \min \left\{ N' \left(\frac{Df(2^n x, 2^n y, 2^n z, 2^n w)}{2 \cdot 4^n}, \frac{t}{8} \right), N' \left(\frac{Df(-2^n x, -2^n y, -2^n z, -2^n w)}{2 \cdot 4^n}, \frac{t}{8} \right), \right. \\
 & \quad \left. N' \left(2^{n-1} Df \left(\frac{x}{2^n}, \frac{y}{2^n}, \frac{z}{2^n}, \frac{w}{2^n} \right), \frac{t}{8} \right), N' \left(2^{n-1} Df \left(\frac{-x}{2^n}, \frac{-y}{2^n}, \frac{-z}{2^n}, \frac{-w}{2^n} \right), \frac{t}{8} \right) \right\} \\
 & \geq \min \left\{ N \left(x, 2^{(2q-1)n-4q} t^q \right), N \left(y, 2^{(2q-1)n-4q} t^q \right), N \left(z, 2^{(2q-1)n-4q} t^q \right), \right. \\
 & \quad \left. N \left(w, 2^{(2q-1)n-4q} t^q \right), N \left(x, 2^{(1-q)n-4q} t^q \right), N \left(y, 2^{(1-q)n-4q} t^q \right), \right. \\
 & \quad \left. N \left(z, 2^{(1-q)n-4q} t^q \right), N \left(w, 2^{(1-q)n-4q} t^q \right) \right\}
 \end{aligned} \tag{2.25}$$

for all $x, y, z, w \in X$ and $t > 0$. Observe that all the terms on the right-hand side of the above inequality tend to 1 as $n \rightarrow \infty$, since $1/2 < q < 1$. Hence, with arguments similar to those of (2.13)–(2.15), we can say that $DF(x, y, z, w) = 0$ for all $x, y, z, w \in X$. Recall that, in Case 1, inequality (2.3) follows from (2.12). By the same reasoning, we get (2.3) from (2.24) in this case.

Now to prove the uniqueness of F , let F' be another quadratic-additive mapping satisfying (2.3). Then, together with (N4), (2.3), and (2.18), we have

$$\begin{aligned}
 & N'(F(x) - F'(x), t) \\
 & = N'(J_n F(x) - J_n F'(x), t) \\
 & \geq \min \left\{ N' \left(J_n F(x) - J_n f(x), \frac{t}{2} \right), N' \left(J_n f(x) - J_n F'(x), \frac{t}{2} \right) \right\} \\
 & \geq \min \left\{ N' \left(\frac{(F - f)(2^n x)}{2 \cdot 4^n}, \frac{t}{8} \right), N' \left(\frac{(f - F')(2^n x)}{2 \cdot 4^n}, \frac{t}{8} \right), \right. \\
 & \quad \left. N' \left(\frac{(F - f)(-2^n x)}{2 \cdot 4^n}, \frac{t}{8} \right), N' \left(\frac{(f - F')(-2^n x)}{2 \cdot 4^n}, \frac{t}{8} \right), \right. \\
 & \quad \left. N' \left(2^{n-1} \left((F - f) \left(\frac{x}{2^n} \right) \right), \frac{t}{8} \right), N' \left(2^{n-1} \left((f - F') \left(\frac{x}{2^n} \right) \right), \frac{t}{8} \right), \right. \\
 & \quad \left. N' \left(2^{n-1} \left((F - f) \left(\frac{-x}{2^n} \right) \right), \frac{t}{8} \right), N' \left(2^{n-1} \left((f - F') \left(\frac{-x}{2^n} \right) \right), \frac{t}{8} \right) \right\}
 \end{aligned}$$

$$\geq \min \left\{ \sup_{t' < t} N \left(x, 2^{(2q-1)n-4q} (4-2^p)^q (2^p-2)^q t'^q \right), \right. \\ \left. \sup_{t' < t} N \left(x, 2^{(1-q)n-4q} (4-2^p)^q (2^p-2)^q t'^q \right) \right\} \quad (2.26)$$

for all $x \in X$ and $n \in \mathbb{N}$. Since $\lim_{n \rightarrow \infty} 2^{(2q-1)n-4q} = \lim_{n \rightarrow \infty} 2^{(1-q)n-4q} = \infty$ in this case, both terms on the right-hand side of the above inequality tend to 1 as $n \rightarrow \infty$ by (N5). This implies that $N'(F(x) - F'(x), t) = 1$ and so $F(x) = F'(x)$ for all $x \in X$ by (N2).

Case 3. Finally, we take $0 < q < (1/2)$ and define $J_n f : X \rightarrow Y$ by

$$J_n f(x) = \frac{1}{2} \left(4^n (f(2^{-n}x) + f(-2^{-n}x)) + 2^n \left(f\left(\frac{x}{2^n}\right) - f\left(-\frac{x}{2^n}\right) \right) \right) \quad (2.27)$$

for all $x \in X$. Then we have $J_0 f(x) = f(x)$ and

$$J_j f(x) - J_{j+1} f(x) = -\frac{4^j}{2} Df \left(\frac{x}{2^{j+1}}, -\frac{x}{2^{j+1}}, \frac{x}{2^{j+1}}, -\frac{x}{2^{j+1}} \right) \\ - 2^{j-2} Df \left(\frac{x}{2^{j+1}}, \frac{x}{2^{j+1}}, \frac{x}{2^{j+1}}, -\frac{x}{2^{j+1}} \right) \\ + 2^{j-2} Df \left(-\frac{x}{2^{j+1}}, -\frac{x}{2^{j+1}}, -\frac{x}{2^{j+1}}, \frac{x}{2^{j+1}} \right) \quad (2.28)$$

which implies that if $n + m > m \geq 0$ then

$$N' \left(J_m f(x) - J_{n+m} f(x), \frac{2}{2^p} \sum_{j=m}^{n+m-1} \left(\left(\frac{4}{2^p} \right)^j + \left(\frac{2}{2^p} \right)^j \right) t^p \right) \\ \geq \min_{j=m}^{n+m-1} \left\{ \min \left\{ N' \left(-\frac{4^j}{2} Df \left(\frac{x}{2^{j+1}}, -\frac{x}{2^{j+1}}, \frac{x}{2^{j+1}}, -\frac{x}{2^{j+1}} \right), \frac{2 \cdot 4^j t^p}{2^{(j+1)p}} \right), \right. \right. \\ N' \left(-2^{j-2} Df \left(\frac{x}{2^{j+1}}, \frac{x}{2^{j+1}}, \frac{x}{2^{j+1}}, -\frac{x}{2^{j+1}} \right), \frac{2^j t^p}{2^{(j+1)p}} \right), \\ \left. \left. N' \left(2^{j-2} Df \left(-\frac{x}{2^{j+1}}, -\frac{x}{2^{j+1}}, -\frac{x}{2^{j+1}}, \frac{x}{2^{j+1}} \right), \frac{2^j t^p}{2^{(j+1)p}} \right) \right\} \right\} \\ \geq \min_{j=m}^{n+m-1} \left\{ N \left(\frac{x}{2^{j+1}}, \frac{t}{2^{j+1}} \right) \right\} = N(x, t) \quad (2.29)$$

for all $x \in X$ and $t > 0$. Similar to the previous cases, it leads us to define the mapping $F : X \rightarrow Y$ by $F(x) := N' - \lim_{n \rightarrow \infty} J_n f(x)$. Putting $m = 0$ in the above inequality, we have

$$N'(f(x) - J_n f(x), t) \geq N\left(x, \frac{t^q}{\left(\sum_{j=0}^{n-1} \left((1/2)(4/2^p)^{j+1} + (2/2^p)^{j+1}\right)\right)^q}\right) \quad (2.30)$$

for all $x \in X$ and $t > 0$. Notice that

$$\begin{aligned} & N'\left(DJ_n f(x, y, z, w), \frac{t}{2}\right) \\ & \geq \min\left\{N'\left(\frac{4^n}{2} Df\left(\frac{x}{2^n}, \frac{y}{2^n}, \frac{z}{2^n}, \frac{w}{2^n}\right), \frac{t}{8}\right), N'\left(\frac{4^n}{2} Df\left(\frac{-x}{2^n}, \frac{-y}{2^n}, \frac{-z}{2^n}, \frac{-w}{2^n}\right), \frac{t}{8}\right), \right. \\ & \quad \left. N'\left(2^{n-1} Df\left(\frac{x}{2^n}, \frac{y}{2^n}, \frac{z}{2^n}, \frac{w}{2^n}\right), \frac{t}{8}\right), N'\left(2^{n-1} Df\left(\frac{-x}{2^n}, \frac{-y}{2^n}, \frac{-z}{2^n}, \frac{-w}{2^n}\right), \frac{t}{8}\right)\right\} \\ & \geq \min\left\{N\left(x, 2^{(1-2q)n-4q} t^q\right), N\left(y, 2^{(1-2q)n-4q} t^q\right), N\left(z, 2^{(1-2q)n-4q} t^q\right), \right. \\ & \quad \left. N\left(w, 2^{(1-2q)n-4q} t^q\right), N\left(x, 2^{(1-q)n-4q} t^q\right), N\left(y, 2^{(1-q)n-4q} t^q\right), \right. \\ & \quad \left. N\left(z, 2^{(1-q)n-4q} t^q\right), N\left(w, 2^{(1-q)n-4q} t^q\right)\right\} \end{aligned} \quad (2.31)$$

for each $x, y, z, w \in X$ and $t > 0$. Since $0 < q < (1/2)$, all terms on the right-hand side tend to 1 as $n \rightarrow \infty$, which implies that the last term of (2.13) tends to 1 as $n \rightarrow \infty$. Therefore, we can say that $DF \equiv 0$. Moreover, using arguments similar to those of (2.13)–(2.15) in Case 1, we get inequality (2.3) from (2.30) in this case. To prove the uniqueness of F , let $F' : X \rightarrow Y$ be another quadratic-additive mapping satisfying (2.3). Then by (2.18), we get

$$\begin{aligned} & N'(F(x) - F'(x), t) \\ & \geq \min\left\{N'\left(J_n F(x) - J_n f(x), \frac{t}{2}\right), N'\left(J_n f(x) - J_n F'(x), \frac{t}{2}\right)\right\} \\ & \geq \min\left\{N'\left(\frac{4^n}{2} \left((F - f)\left(\frac{x}{2^n}\right)\right), \frac{t}{8}\right), N'\left(\frac{4^n}{2} \left((f - F')\left(\frac{x}{2^n}\right)\right), \frac{t}{8}\right), \right. \\ & \quad \left. N'\left(\frac{4^n}{2} \left((F - f)\left(-\frac{x}{2^n}\right)\right), \frac{t}{8}\right), N'\left(\frac{4^n}{2} \left((f - F')\left(-\frac{x}{2^n}\right)\right), \frac{t}{8}\right), \right. \\ & \quad \left. N'\left(2^{n-1} \left((F - f)\left(\frac{x}{2^n}\right)\right), \frac{t}{8}\right), N'\left(2^{n-1} \left((f - F')\left(\frac{x}{2^n}\right)\right), \frac{t}{8}\right), \right. \\ & \quad \left. N'\left(2^{n-1} \left((F - f)\left(-\frac{x}{2^n}\right)\right), \frac{t}{8}\right), N'\left(2^{n-1} \left((f - F')\left(-\frac{x}{2^n}\right)\right), \frac{t}{8}\right)\right\} \\ & \geq \sup_{t' < t} N\left(x, \frac{2^{(1-2q)n-4q} (2^p - 4)(2^p - 2)t'^q}{(2^p - 3)^q}\right) \end{aligned} \quad (2.32)$$

for all $x \in X$ and $n \in \mathbb{N}$. Observe that, for $0 < q < (1/2)$, the last term tends to 1 as $n \rightarrow \infty$ by (N5). This implies that $N'(F(x) - F'(x), t) = 1$ and $F(x) = F'(x)$ for all $x \in X$ by (N2). $\square$

Remark 2.3. Consider a mapping $f : X \rightarrow Y$ satisfying (2.2) for all $x, y, z, w \in X$ and a real number $q < 0$. Take any $t > 0$. If we choose a real number s with $0 < 4s < t$, then

$$\begin{aligned} N'(Df(x, y, z, w), t) &\geq N'(Df(x, y, z, w), 4s) \\ &\geq \min\{N(x, s^q), N(y, s^q), N(z, s^q), N(w, s^q)\} \end{aligned} \quad (2.33)$$

for all $x, y, z, w \in X$. Since $q < 0$, we have $\lim_{s \rightarrow 0^+} s^q = \infty$. This implies that

$$\lim_{s \rightarrow 0^+} N(x, s^q) = \lim_{s \rightarrow 0^+} N(y, s^q) = \lim_{s \rightarrow 0^+} N(z, s^q) = \lim_{s \rightarrow 0^+} N(w, s^q) = 1, \quad (2.34)$$

and so

$$N'(Df(x, y, z, w), t) = 1 \quad (2.35)$$

for all $x, y, z, w \in X$ and $t > 0$. By (N2), we are allowed to get $Df(x, y, z, w) = 0$ for all $x, y, z, w \in X$. In other words, f is itself a quadratic-additive mapping if f is a fuzzy q -almost quadratic-additive mapping for the case $q < 0$.

Corollary 2.4. *Let f be an even mapping satisfying all of the conditions of Theorem 2.2. Then there is a unique quadratic mapping $F : X \rightarrow Y$ such that*

$$N'(F(x) - f(x), t) \geq \sup_{t' < t} N\left(x, \left(\frac{|4 - 2^p|t'}{2}\right)^q\right) \quad (2.36)$$

for all $x \in X$ and $t > 0$, where $p = 1/q$.

Proof. Let $J_n f$ be defined as in Theorem 2.2. Since f is an even mapping, we obtain

$$J_n f(x) = \begin{cases} \frac{f(2^n x) + f(-2^n x)}{2 \cdot 4^n} & \text{if } q > \frac{1}{2}, \\ \frac{1}{2}(4^n(f(2^{-n}x) + f(-2^{-n}x))) & \text{if } 0 < q < \frac{1}{2} \end{cases} \quad (2.37)$$

for all $x \in X$. Notice that $J_0 f(x) = f(x)$ and

$$J_j f(x) - J_{j+1} f(x) = \begin{cases} \frac{Df(2^j x, -2^j x, 2^j x, -2^j x)}{2 \cdot 4^{j+1}} & \text{if } q > \frac{1}{2}, \\ -\frac{4^j}{2} Df\left(\frac{x}{2^{j+1}}, -\frac{x}{2^{j+1}}, \frac{x}{2^{j+1}}, -\frac{x}{2^{j+1}}\right) & \text{if } 0 < q < \frac{1}{2} \end{cases} \quad (2.38)$$

for all $x \in X$ and $j \in \mathbb{N} \cup \{0\}$. From these, using the similar method in Theorem 2.2, we obtain the quadratic-additive mapping F satisfying (2.36). Notice that $F(x) := N' - \lim_{n \rightarrow \infty} J_n f(x)$ for all $x \in X$, F is even, $F(2x) = 4F(x)$, and $DF(x, y, z, w) = 0$ for all $x, y, z, w \in X$. Hence, we get

$$F(x+y) + F(x-y) - 2F(x) - 2F(y) = -DF(x, y, x, -x) = 0 \quad (2.39)$$

for all $x, y \in X$. This means that F is a quadratic mapping. $\square$

Corollary 2.5. *Let f be an odd mapping satisfying all of the conditions of Theorem 2.2. Then there is a unique additive mapping $F : X \rightarrow Y$ such that*

$$N'(F(x) - f(x), t) \geq \sup_{t' < t} N\left(x, \left(\frac{|2 - 2^p|t'}{2}\right)^q\right) \quad (2.40)$$

for all $x \in X$ and $t > 0$, where $p = 1/q$.

Proof. Let $J_n f$ be defined as in Theorem 2.2. Since f is an odd mapping, we obtain

$$J_n f(x) = \begin{cases} \frac{f(2^n x) + f(-2^n x)}{2^{n+1}} & \text{if } q > 1, \\ 2^{n-1}(f(2^{-n} x) + f(-2^{-n} x)) & \text{if } 0 < q < 1 \end{cases} \quad (2.41)$$

for all $x \in X$. Notice that $J_0 f(x) = f(x)$ and

$$\begin{aligned} J_j f(x) - J_{j+1} f(x) &= \begin{cases} \frac{Df(2^j x, 2^j x, 2^j x, -2^j x)}{2^{j+3}} - \frac{Df(-2^j x, -2^j x, -2^j x, 2^j x)}{2^{j+3}} & \text{if } q > 1, \\ 2^{j-2} \left(Df\left(-\frac{x}{2^{j+1}}, -\frac{x}{2^{j+1}}, -\frac{x}{2^{j+1}}, \frac{x}{2^{j+1}}\right) - Df\left(\frac{x}{2^{j+1}}, \frac{x}{2^{j+1}}, \frac{x}{2^{j+1}}, -\frac{x}{2^{j+1}}\right) \right) & \text{if } 0 < q < 1 \end{cases} \end{aligned} \quad (2.42)$$

for all $x \in X$ and $j \in \mathbb{N} \cup \{0\}$. From these, using the similar method in Theorem 2.2, we obtain the quadratic-additive mapping F satisfying (2.40). Notice that $F(x) := N' - \lim_{n \rightarrow \infty} J_n f(x)$ for all $x \in X$, F is odd, $F(2x) = 2F(x)$, and $DF(x, y, z, w) = 0$ for all $x, y, z, w \in X$. Hence, we get

$$F(x+y) - F(x) - F(y) = -DF\left(\frac{x-y}{2}, \frac{-x-y}{2}, \frac{x-y}{2}, \frac{-x+y}{2}\right) = 0 \quad (2.43)$$

for all $x, y \in X$. This means that F is an additive mapping. $\square$

We can use Theorem 2.2 to get a classical result in the framework of normed spaces. Let $(X, \|\cdot\|)$ be a normed linear space. Then we can define a fuzzy norm N_X on X by

$$N_X(x, t) = \begin{cases} 0 & \text{if } t \leq \|x\|, \\ 1 & \text{if } t > \|x\|, \end{cases} \quad (2.44)$$

where $x \in X$ and $t \in \mathbb{R}$, see [22]. Suppose that $f : X \rightarrow Y$ is a mapping into a Banach space $(Y, \|\cdot\|)$ such that

$$\|Df(x, y, z, w)\| \leq \|x\|^p + \|y\|^p + \|z\|^p + \|w\|^p \quad (2.45)$$

for all $x, y, z, w \in X$, where $p > 0$ and $p \neq 1, 2$. Let N_Y be a fuzzy norm on Y . Then we get

$$N_Y(Df(x, y, z, w), s + t + u + v) = \begin{cases} 0 & \text{if } s + t + u + v \leq \|Df(x, y, z, w)\|, \\ 1 & \text{if } s + t + u + v > \|Df(x, y, z, w)\| \end{cases} \quad (2.46)$$

for all $x, y, z, w \in X$ and $s, t, u, v \in \mathbb{R}$. Consider the case

$$N_Y(Df(x, y, z, w), s + t + u + v) = 0. \quad (2.47)$$

This implies that

$$\|x\|^p + \|y\|^p + \|z\|^p + \|w\|^p \geq \|Df(x, y, z, w)\| \geq s + t + u + v, \quad (2.48)$$

and so $\|x\|^p \geq s$, $\|y\|^p \geq t$, $\|z\|^p \geq u$, or $\|w\|^p \geq v$ in this case. Hence, for $q = 1/p$, we have

$$\min\{N_X(x, s^q), N_X(y, t^q), N_X(z, u^q), N_X(w, v^q)\} = 0 \quad (2.49)$$

for all $x, y, z, w \in X$ and $s, t, u, v > 0$. Therefore, in every case, the inequality

$$N_Y(Df(x, y, z, w), s + t + u + v) \geq \min\{N_X(x, s^q), N_X(y, t^q), N_X(z, u^q), N_X(w, v^q)\} \quad (2.50)$$

holds. It means that f is a fuzzy q -almost quadratic-additive mapping, and by Theorem 2.2, we get the following stability result.

Corollary 2.6. *Let $(X, \|\cdot\|)$ be a normed linear space and let $(Y, \|\cdot\|)$ be a Banach space. If $f : X \rightarrow Y$ satisfies*

$$\|Df(x, y, z, w)\| \leq \|x\|^p + \|y\|^p + \|z\|^p + \|w\|^p \quad (2.51)$$

for all $x, y, z, w \in X$, where $p > 0$ and $p \neq 1, 2$, then there is a unique quadratic-additive mapping $F : X \rightarrow Y$ such that

$$|||F(x) - f(x)||| \leq \left(\frac{2}{|2 - 2^p|} + \frac{2}{|4 - 2^p|} \right) \|x\|^p \quad (2.52)$$

for all $x \in X$.

Lemma 2.7. If $F : X \rightarrow Y$ is a mapping such that

$$Df(x, y, z, w) = 0 \quad (2.53)$$

for all $x, y, z, w \in X \setminus \{0\}$ with $F(0) = 0$, then

$$DF(x, y, z, w) = 0 \quad (2.54)$$

for all $x, y, z, w \in X$.

Proof. We easily know that

$$\begin{aligned} DF(x, y, z, 0) &= DF(2x, y, z, -x) - DF(2x, y, -x, -x) + DF(2x, y, x, -x) \\ &\quad - DF(2x, z, -x, -x) + DF(2x, z, x, -x) - 2DF(2x, x, x, -x) \\ &\quad + DF(x, -x, -x, -x) = 0 \end{aligned} \quad (2.55)$$

for all $x, y, z \in X \setminus \{0\}$. Using a similar method, we have $DF(x, y, 0, w) = 0$, $DF(x, 0, z, w) = 0$, $DF(0, y, z, w) = 0$, $DF(x, y, 0, 0) = 0$, $DF(x, 0, z, 0) = 0$, $DF(x, 0, 0, w) = 0$, $DF(0, y, z, 0) = 0$, $DF(0, 0, z, w) = 0$, $DF(0, y, 0, w) = 0$, $DF(x, 0, 0, 0) = 0$, $DF(0, y, 0, 0) = 0$, $DF(0, 0, z, 0) = 0$, $DF(0, 0, 0, w) = 0$, and $DF(0, 0, 0, 0) = 0$ for all $x, y, z, w \in X \setminus \{0\}$, as we desired. $\square$

References

- [1] S. M. Ulam, *A Collection of Mathematical Problems*, Interscience, New York, NY, USA, 1968.
- [2] D. H. Hyers, "On the stability of the linear functional equation," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 27, pp. 222–224, 1941.
- [3] T. Aoki, "On the stability of the linear transformation in Banach spaces," *Journal of the Mathematical Society of Japan*, vol. 2, pp. 64–66, 1950.
- [4] T. M. Rassias, "On the stability of the linear mapping in Banach spaces," *Proceedings of the American Mathematical Society*, vol. 72, no. 2, pp. 297–300, 1978.
- [5] A. Adam and S. Czerwik, "On the stability of the quadratic functional equation in topological spaces," *Banach Journal of Mathematical Analysis*, vol. 1, no. 2, pp. 245–251, 2007.
- [6] P. Găvruta, "A generalization of the Hyers-Ulam-Rassias stability of approximately additive mappings," *Journal of Mathematical Analysis and Applications*, vol. 184, no. 3, pp. 431–436, 1994.
- [7] K.-W. Jun and Y.-H. Lee, "A generalization of the Hyers-Ulam-Rassias stability of the Pexiderized quadratic equations. II," *Kyungpook Mathematical Journal*, vol. 47, no. 1, pp. 91–103, 2007.
- [8] H.-M. Kim, "On the stability problem for a mixed type of quartic and quadratic functional equation," *Journal of Mathematical Analysis and Applications*, vol. 324, no. 1, pp. 358–372, 2006.
- [9] Y.-H. Lee, "On the Hyers-Ulam-Rassias stability of the generalized polynomial function of degree 2," *Journal of the Chungcheong Mathematical Society*, no. 22, pp. 201–209, 2009.

- [10] Y.-H. Lee, "On the stability of the monomial functional equation," *Bulletin of the Korean Mathematical Society*, vol. 45, no. 2, pp. 397–403, 2008.
- [11] Y.-H. Lee and K.-W. Jun, "A generalization of the Hyers-Ulam-Rassias stability of Jensen's equation," *Journal of Mathematical Analysis and Applications*, vol. 238, no. 1, pp. 305–315, 1999.
- [12] Y.-H. Lee and K.-W. Jun, "A generalization of the Hyers-Ulam-Rassias stability of the Pexider equation," *Journal of Mathematical Analysis and Applications*, vol. 246, no. 2, pp. 627–638, 2000.
- [13] Y.-H. Lee and K.-W. Jun, "On the stability of approximately additive mappings," *Proceedings of the American Mathematical Society*, vol. 128, no. 5, pp. 1361–1369, 2000.
- [14] M. Mirzavaziri and M. S. Moslehian, "A fixed point approach to stability of a quadratic equation," *Bulletin of the Brazilian Mathematical Society*, vol. 37, no. 3, pp. 361–376, 2006.
- [15] A. Najati and A. Rahimi, "A fixed point approach to the stability of a generalized cauchy functional equation," *Banach Journal of Mathematical Analysis*, vol. 2, no. 1, pp. 105–112, 2008.
- [16] F. Skof, "Local properties and approximation of operators," *Rendiconti del Seminario Matematico e Fisico di Milano*, vol. 53, pp. 113–129, 1983.
- [17] A. K. Katsaras, "Fuzzy topological vector spaces. II," *Fuzzy Sets and Systems*, vol. 12, no. 2, pp. 143–154, 1984.
- [18] T. Bag and S. K. Samanta, "Finite dimensional fuzzy normed linear spaces," *Journal of Fuzzy Mathematics*, vol. 11, no. 3, pp. 687–705, 2003.
- [19] S. C. Cheng and J. N. Mordeson, "Fuzzy linear operators and fuzzy normed linear spaces," *Bulletin of the Calcutta Mathematical Society*, vol. 86, no. 5, pp. 429–436, 1994.
- [20] I. Kramosil and J. Michálek, "Fuzzy metrics and statistical metric spaces," *Kybernetika*, vol. 11, no. 5, pp. 336–344, 1975.
- [21] A. K. Mirmostafae and M. S. Moslehian, "Fuzzy versions of Hyers-Ulam-Rassias theorem," *Fuzzy Sets and Systems*, vol. 159, no. 6, pp. 720–729, 2008.
- [22] A. K. Mirmostafae and M. S. Moslehian, "Fuzzy almost quadratic functions," *Results in Mathematics*, vol. 52, no. 1-2, pp. 161–177, 2008.
- [23] I.-S. Chang, E. H. Lee, and H.-M. Kim, "On Hyers-Ulam-Rassias stability of a quadratic functional equation," *Mathematical Inequalities and Applications*, vol. 6, no. 1, pp. 87–95, 2003.