Hindawi Publishing Corporation
International Journal of Mathematics and Mathematical Sciences
Volume 2011, Article ID 795934, 13 pages
doi:10.1155/2011/795934

Research Article

Fuzzy Filter Spectrum of a BCK Algebra

Xiao Long Xin, Wei Ji, and Xiu Juan Hua

Department of Mathematics, Northwest University, Xi'an 710127, China

Correspondence should be addressed to Xiao Long Xin, xlxin@nwu.edu.cn

Received 14 October 2010; Accepted 22 December 2010

Academic Editor: Young Bae Jun

Copyright © 2011 Xiao Long Xin et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The notion of fuzzy s-prime filters of a bounded BCK-algebra is introduced. We discuss the relation between fuzzy s-prime filters and fuzzy prime filters. By the fuzzy s-prime filters of a bounded commutative BCK-algebra X, we establish a fuzzy topological structure on X. We prove that the set of all fuzzy s-prime filters of a bounded commutative BCK-algebra forms a topological space. Moreover, we show that the set of all fuzzy s-prime filters of a bounded implicative BCK-algebra is a Hausdorff space.

1. Introduction

BCK-algebras are an important class of logical algebras introduced by Iséki in 1966 (see [1–3]). Since then, a great deal of the literature has been produced on the theory of BCK-algebras. In particular, emphasis seems to have been put on the ideal and filter theory of BCK-algebras (see [4]). The concept of fuzzy sets was introduced by Zadeh [5]. At present, these ideas have been applied to other algebraic structures such as semigroups, groups, rings, ideals, modules, vector spaces, and so on (see [6, 7]). In 1991, Ougen [8] applied the concept of fuzzy sets to BCK-algebras. For the general development of BCK-algebras the fuzzy ideal theory and fuzzy filter theory play important roles (see [9–12]). Meng [13] introduced the notion of BCK-filters and investigated some results. Jun et al. [9, 10] studied the fuzzification of BCK-filters. Meng [13] showed how to generate the BCK-filter by a subset of Alò, and Deeba [14] attempted to study the topological aspects of the BCK-structures. They initiated the study of various topologies on BCK-algebras analogous to which has already been studied on lattices. In [15], Jun et al. introduced the notion of topological BCI-algebras and found some elementary properties.

In this paper, the topological structure and fuzzy structure on BCK-algebras are investigated together. We introduce the concept of fuzzy s-prime filters and discuss some related properties. By the fuzzy s-prime filters, we establish a fuzzy topological structure on bounded commutative BCK-algebras and bounded implicative BCK-algebras, respectively.

2. Preliminaries

A nonempty set X with a constant 0 and a binary operation denoted by juxtaposition is called a BCK-algebra if for all $x, y, z \in X$ the following conditions hold:

(1) $((xy)(xz))(zy) = 0$,

(2) $(x(xy))y = 0$,

(3) $xx = 0$,

(4) $0x = 0$,

(5) $xy = 0$ and $yx = 0$ imply $x = y$.

A BCK-algebra can be (partially) ordered by $x \leq y$ if and only if $xy = 0$. This ordering is called BCK-ordering. The following statements are true in any BCK-algebra: for all x, y, z,

(6) $x0 = x$.

(7) $(xy)z = (xz)y$.

(8) $xy \leq x$.

(9) $(xy)z \leq (xz)(yz)$.

(10) $x \leq y$ implies $xz \leq yz$ and $zy \leq zx$.

A BCK-algebra X satisfying the identity $x(xy) = y(yx)$ is said to be commutative. If there is a special element 1 of a BCK-algebra X satisfying $x \leq 1$ for all $x \in X$, then 1 is called unit of X. A BCK-algebra with unit is said to be bounded. In a bounded BCK-algebra X, we denote $1x$ by x^* for every $x \in X$.

In a bounded BCK-algebra, we have

(11) $1^* = 0$ and $0^* = 1$.

(12) $y \leq x$ implies $x^* \leq y^*$.

(13) $x^*y^* \leq yx$.

Now, we review some fuzzy logic concepts. A fuzzy set in X is a function $\mu : X \to [0, 1]$. We use the notation X_μ for $\{x \in X \mid \mu(x) = \mu(1)\}$ and μ_t, called a level subset of μ, for $\{x \in X \mid \mu(x) \geq t\}$ where $t \in [0, 1]$.

In this paper, unless otherwise specified, X denotes a bounded BCK-algebra. A nonempty subset F of X is called a BCK-filter of X if

(F1) $1 \in F$,

(F2) $(x^*y^*)^* \in F$ and $y \in F$ imply $x \in F$ for all $x, y \in X$.

Note that the intersection of a family of BCK-filters is a BCK-filter. For convenience, we call a BCK-filter of X as a filter of X, and write $F <_{\mathrm{F}} X$.

Let μ be a fuzzy set in X. Then, μ is called a fuzzy filter of X if

(FF1) $\mu(1) \geq \mu(x)$,

(FF2) $\mu(x) \geq \min\{\mu(x^*y^*)^*, \mu(y)\}$, for all $x, y \in X$. In this case, we write $\mu <_{\mathrm{FF}} X$.

Note that in a bounded commutative BCK-algebra, the identity $x^*y^* = yx$ holds, then (F2) and

(F3) $(yx)^* \in F$ and $y \in F$ imply $x \in F$ for all x, y in X coincide, and (FF2) and

(FF3) $\mu(x) \geq \min\{\mu(yx)^*, \mu(y)\}$ coincide.

A proper filter F of X is said to be prime, denoted by $F <_{\mathrm{PF}} X$, if, for any $x, y \in X$, $x \vee y \in F$ implies $x \in F$ or $y \in F$.

A nonconstant fuzzy filter μ of X is said to be prime, denoted by $\mu <_{\mathrm{FPF}} X$, if $\mu(x \vee y) \leq \max\{\mu(x), \mu(y)\}$ for all $x, y \in X$.

For any fuzzy sets μ and ν in X, we denote

$$\begin{aligned}
&\mu \subseteq \nu \Longleftrightarrow \mu(x) \leq \nu(x), \quad \forall x, y \text{ in } X,\\
&\mu \cap \nu(x) = \min\{\mu(x), \nu(x)\}, \quad \forall x \in X,\\
&\mu \cup \nu(x) = \max\{\mu(x), \nu(x)\}, \quad \forall x \in X,\\
&\bigcap_{\alpha\in\Omega} \mu_\alpha(x) = \inf_{\alpha\in\Omega} \mu_\alpha(x), \quad \forall x \in X,\\
&\bigcup_{\alpha\in\Omega} \mu_\alpha(x) = \sup_{\alpha\in\Omega} \mu_\alpha(x), \quad \forall x \in X,\\
&\mu\eta(x) = \sup_{x=y\vee z} \{\min\{\mu(y), \eta(z)\}\}.
\end{aligned} \tag{2.1}$$

Lemma 2.1. *Let $\{\eta_\alpha \mid \alpha \in \Omega\}$ be a family of fuzzy filters of X. Then, $\bigcap_{\alpha\in\Omega} \eta_\alpha$ is a fuzzy filter of X.*

Proof. Let $x \in X$. For any $\alpha \in \Omega$, $\eta_\alpha(1) \geq \eta_\alpha(x)$ since $\eta_\alpha <_{\mathrm{FF}} X$. Then, $\inf_{\alpha\in\Omega}\eta_\alpha(1) \geq \inf_{\alpha\in\Omega}\eta_\alpha(x)$ and so, $\bigcap_{\alpha\in\Omega} \eta_\alpha(1) \geq \bigcap_{\alpha\in\Omega} \eta_\alpha(x)$. (FF1) holds.

Moreover, for any $\varepsilon > 0$, there exists $\alpha(\varepsilon) \in \Omega$ such that

$$\begin{aligned}
\bigcap_{\alpha\in\Omega} \eta_\alpha(x) + \varepsilon &= \inf_{\alpha\in\Omega} \eta_\alpha(x) + \varepsilon\\
&\geq \eta_{\alpha(\varepsilon)}(x)\\
&\geq \min\{\eta_{\alpha(\varepsilon)}((x^*y^*)^*), \eta_{\alpha(\varepsilon)}(y)\}\\
&\geq \min\left\{\inf_{\alpha\in\Omega} \eta_\alpha(x^*y^*)^*, \inf_{\alpha\in\Omega} \eta_\alpha(y)\right\}\\
&= \min\left\{\bigcap_{\alpha\in\Omega} \eta_\alpha(x^*y^*)^*, \bigcap_{\alpha\in\Omega} \eta_\alpha(y)\right\}.
\end{aligned} \tag{2.2}$$

Since ε is arbitrary, we get $\bigcap_{\alpha\in\Omega} \eta_\alpha(x) \geq \min\{\bigcap_{\alpha\in\Omega} \eta_\alpha(x^*y^*)^*, \bigcap_{\alpha\in\Omega} \eta_\alpha(y)\}$. So, (FF2) holds.

Therefore, $\bigcap_{\alpha\in\Omega} \eta_\alpha$ is a fuzzy filter of X. □

Lemma 2.2 (see, [16]). *Let μ be a fuzzy filter of X. For any $x, y \in X$, if $x \leq y$, then $\mu(x) \leq \mu(y)$.*

Definition 2.3. Let μ be a fuzzy subset of X. Then the fuzzy filter generated by μ, which is denoted by $\langle\mu\rangle$, is defined as

$$\langle\mu\rangle = \bigcap\{\eta : \mu \subseteq \eta, \eta <_{\mathrm{FF}} X\}. \tag{2.3}$$

Obviously, we get $\mu \subseteq \langle\mu\rangle$, and if $\mu <_{\mathrm{FF}} X$, then $\mu = \langle\mu\rangle$.

Lemma 2.4. *If $\mu, \eta <_{FF} X$, then $\mu\eta = \mu \cap \eta$.*

Proof. Let $x \in X$, $x = a \vee b$ and μ, η be fuzzy filters. Then, by Lemma 2.2, $\mu(a) \leq \mu(a \vee b) = \mu(x)$ and $\eta(b) \leq \eta(a \vee b) = \eta(x)$. Hence, $\min\{\mu(a), \eta(b)\} \leq \mu \cap \eta(x)$.

Therefore, $\mu\eta \leq \mu \cap \eta(x)$, or equivalently $\mu\eta \subseteq \mu \cap \eta$.

Conversely, $\mu\eta(x) = \sup_{x=y\vee z}\{\min\{\mu(y), \eta(z)\}\} \geq \min\{\mu(x), \eta(x)\} = \mu \cap \eta(x)$. So $\mu\eta \supseteq \mu \cap \eta$.

Thus, $\mu\eta = \mu \cap \eta$. □

Corollary 2.5. *If $\mu, \eta <_{FF} X$, $\mu\eta <_{FF} X$.*

Lemma 2.6. *If $\eta <_{FF} X$, $\mu\eta \subseteq \eta$.*

Proof. Let $\eta <_{\mathrm{FF}} X$. If $x = y \vee z$, then from Lemma 2.2 we know $\eta(z) \leq \eta(x)$. Thus, $\mu\eta(x) = \sup_{x=y\vee z}\{\min\{\mu(y), \mu(z)\} \leq \sup_{x=y\vee z}\{\eta(z)\} \leq \eta(x)$. So, $\mu\eta \subseteq \eta$. □

3. Fuzzy Filter Spectrum

Definition 3.1. A nonconstant fuzzy filter μ of X is said to be s-prime if for all $\theta, \sigma <_{\mathrm{FF}} X$, $\theta\sigma \subseteq \mu$ implies $\theta \subseteq \mu$ or $\sigma \subseteq \mu$. In this case, we write $\mu <_{\mathrm{FSP}} X$.

In this paper, we give some notations in the following.

(i) $F(X) = \{\mu \mid \mu <_{\mathrm{FSP}} X\}$.

(ii) $V(\theta) = \{\mu \in F(X) \mid \theta \subseteq \mu\}$, where θ is a fuzzy subset of X.

(iii) $F(\theta) = F(X)\backslash V(\theta) = \{\mu \in F(X) \mid \theta \not\subseteq \mu\}$, where $F(X)\backslash V(\theta)$ is called the complement of $V(\theta)$ in $F(X)$.

Lemma 3.2. *If σ is a fuzzy subset of X, then $V(\langle\sigma\rangle) = V(\sigma)$. So $F(\sigma) = F(\langle\sigma\rangle)$.*

Proof. Let $\mu \in V(\sigma)$, then $\sigma \subseteq \mu$ and so $\langle\sigma\rangle \subseteq \mu$. Hence, $\mu \in V(\langle\sigma\rangle)$. Conversely, let $\mu \in V(\langle\sigma\rangle)$, then $\langle\sigma\rangle \subseteq \mu$. Note that $\sigma \subseteq \langle\sigma\rangle \subseteq \mu$, we get $\mu \in V(\sigma)$. Therefore, $V(\sigma) = V(\langle\sigma\rangle)$. □

Theorem 3.3. *Let $\zeta = \{F(\theta) \mid \theta <_{\mathrm{FF}} X\}$. Then the pair $(F(X), \zeta)$ is a topological space.*

Proof. Consider $\theta_0 = 0$ and $\theta_1 = 1$. Then $\theta_0, \theta_1 <_{\mathrm{FF}} X$, $F(\theta_0) = \emptyset$ and $F(\theta_1) = F(X)$. Thus, $F(X)$, $\emptyset \in \zeta$.

Then, we prove that ζ is closed under finite intersection.

Let η and θ be two fuzzy filters of X. We claim that $V(\theta) \cup V(\eta) = V(\theta\eta)$. Let $\tau \in V(\theta\eta)$. Then, $\theta\eta \subseteq \tau$. Since $\tau \in F(X)$, we have $\theta \subseteq \tau$ or $\sigma \subseteq \tau$. It follows that $\tau \in V(\theta) \cup V(\eta)$.

Conversely, let $\tau \in V(\theta) \cup V(\eta)$, then $\theta \subseteq \tau$ or $\eta \subseteq \tau$. By Lemma 2.6, $\theta\eta \subseteq \theta$ and $\theta\eta \subseteq \eta$. Thus, $\theta\eta \subseteq \tau$ and so $\tau \in V(\theta\eta)$. It follows that $V(\theta) \cup V(\eta) \subseteq V(\theta\eta)$.

Combining the above arguments we get $V(\theta) \cup V(\eta) = V(\theta\eta)$, or equivalently, $F(\theta) \cap F(\eta) = (F(X) \setminus V(\theta)) \cap (F(X) \setminus V(\eta)) = (F(X) \setminus (V(\theta) \cup V(\eta)) = (F(X) \setminus V(\theta\eta)) = F(\theta\eta)$. By Corollary 2.5, $\theta\eta <_{\mathrm{FF}} X$ and so $F(\theta) \cap F(\eta) = F(\theta\eta) \in \zeta$.

Finally, let $\{\theta_\alpha \mid \alpha \in \Omega\}$ be a family of fuzzy prime filters of X. We will prove that $\bigcap_{\alpha\in\Omega} V(\theta_\alpha) = V(\bigcup_{\alpha\in\Omega} \theta_\alpha)$.

Let $\mu \in \bigcap_{\alpha\in\Omega} V(\theta_\alpha)$, then for any $\alpha \in \Omega$, $\mu \in V(\theta_\alpha)$ and so $\theta_\alpha \subseteq \mu$. Hence, $\bigcup_{\alpha\in\Omega} \theta_\alpha \subseteq \mu$ and thus $\mu \in V(\bigcup_{\alpha\in\Omega} \theta_\alpha)$.

Conversely, let $\mu \in V(\bigcup_{\alpha\in\Omega} \theta_\alpha)$, then $\bigcup_{\alpha\in\Omega} \theta_\alpha \subseteq \mu$. Thus, for any $\alpha \in \Omega$, $\theta_\alpha \subseteq \bigcup_{\alpha\in\Omega} \theta_\alpha \subseteq \mu$. Hence, $\mu \in V(\theta_\alpha)$ for all $\alpha \in \Omega$ and so $\mu \in \bigcap_{\alpha\in\Omega} V(\theta_\alpha)$.

This shows that $\bigcap_{\alpha\in\Omega} V(\theta_\alpha) = V(\bigcup_{\alpha\in\Omega} \theta_\alpha)$.

By Lemma 3.2, we get $V(\bigcup_{\alpha\in\Omega} \theta_\alpha) = V(\langle \bigcup_{\alpha\in\Omega} \theta_\alpha \rangle)$ and so $\bigcap_{\alpha\in\Omega} V(\theta_\alpha) = V(\langle \bigcup_{\alpha\in\Omega} \theta_\alpha \rangle)$.

Furthermore, we get $\bigcup_{\alpha\in\Omega} F(\theta_\alpha) = \bigcup_{\alpha\in\Omega} (F(X) \setminus V(\theta_\alpha)) = F(X) \setminus \bigcap_{\alpha\in\Omega} V(\theta_\alpha) = F(X) \setminus V(\langle \bigcup_{\alpha\in\Omega} \theta_\alpha \rangle) = F(\langle \bigcup_{\alpha\in\Omega} \theta_\alpha \rangle) \in \zeta$.

It follows that $(F(X), \zeta)$ is a topological space. □

Theorem 3.4. *The collection*

$$ß = \{F(x_\beta) \mid x \in X, \beta \in (0,1]\} \tag{3.1}$$

of ζ is a base of ζ where $x_\beta <_{\mathrm{FF}} X$ is defined by

$$x_\beta(t) = \begin{cases} \beta, & t = x, \\ 0, & t \neq x. \end{cases} \tag{3.2}$$

Proof. By Lemma 3.2, for any $x \in X$, $\beta \in (0,1]$, $F(x_\beta) = F(\langle x_\beta \rangle)$ and so $F(x_\beta) \in \zeta$.

Now, we prove that ß is a base of ζ. It is sufficient to show that for all $F(\theta) \in \zeta$, and $\mu \in F(\theta)$, there exists $F(x_\beta) \in \beta$ such that $\mu \in F(x_\beta)$ and $F(x_\beta) \subseteq F(\theta)$.

Let $F(\theta) \in \zeta$ and $\mu \in F(\theta)$. Then, $\theta \not\subseteq \mu$ and so there exists $x \in X$ such that $\mu(x) < \theta(x)$. Let $\theta(x) = \beta$ and then $\mu \in F(x_\beta)$. Moreover, for any $\sigma \in V(\theta)$, $\sigma(x) \geq \theta(x) = \beta = x_\beta(x)$ and so $x_\beta \subseteq \sigma$. Thus $\sigma \in V(x_\beta)$. This means $V(\theta) \subseteq V(x_\beta)$. It follows that $F(x_\beta) \subseteq F(\theta)$.

Therefore, ß is a base of ζ. □

The topological space $(F(X), \zeta)$ is called fuzzy filter spectrum of X, denoted by FF-spec(X), or $F(X)$ for convenience.

Theorem 3.5. *FF-spec*(X) *is a T_0 space.*

Proof. Let $\mu, \eta \in F(X)$ and $\mu \neq \eta$. Then, $\mu \not\subseteq \eta$ or $\eta \not\subseteq \mu$.

If $\mu \not\subseteq \eta$, then, $\eta \notin V(\mu)$ but $\mu \in V(\mu)$. Moreover, $\eta \in F(\mu)$ but $\mu \notin F(\mu)$.

If $\eta \not\subseteq \mu$, similarly we can get $\mu \in F(\eta)$ but $\eta \notin F(\eta)$. It follows that FF-spec(X) is a T_0 space. □

Lemma 3.6 (see [9]). *Let μ be a fuzzy subset of X. Then, μ is a fuzzy filter of X if and only if μ_t is a filter of X for each $t \in [0,1]$ wherever $\mu_t \neq \emptyset$.*

Lemma 3.7. *A non-constant fuzzy subset μ of X is a fuzzy prime filter if and only if μ_t is a prime filter of X for each $t \in [0,1]$ whenever $\mu_t \neq \emptyset$.*

Proof. Let μ be a fuzzy prime filter and $t \in [0,1]$ such that $\mu_t \neq \emptyset$. Then by Lemma 3.6, μ_t is a filter of X.

Suppose $x \vee y \in \mu_t$. It follows that $\mu(x \vee y) \geq t$. Since μ is prime, we have $\max\{\mu(x), \mu(y)\} \geq \mu(x \vee y) \geq t$ and thus $\mu(x) \geq t$ or $\mu(y) \geq t$. It follows that $x \in \mu_t$ or $y \in \mu_t$. Therefore μ_t is a prime filter.

Conversely, suppose that for each $t \in [0,1]$, μ_t is a prime filter whenever $\mu_t \neq \emptyset$. If μ is not a fuzzy prime filter, then there exist $x, y \in X$ such that $\mu(x \vee y) > \max\{\mu(x), \mu(y)\}$. Take t satisfying $\mu(x \vee y) > t > \max\{\mu(x), \mu(y)\}$. Then $x \vee y \in \mu_t$. Since μ_t is a prime filter of X, then $x \vee y \in \mu_t$ implies $x \in \mu_t$ or $y \in \mu_t$. But on the other hand, $\mu(x) \leq \max\{\mu(x), \mu(y)\} < t$ and $\mu(y) \leq \max\{\mu(x), \mu(y)\} < t$ imply $x \notin \mu_t$ and $y \notin \mu_t$, a contradiction. It follows that μ is indeed a fuzzy prime filter. □

Lemma 3.8 (see [13]). *Let X be a bounded commutative BCK-algebra and F be a BCK-filter of X. Then, F is prime if and only if, for any filters A, B, $F = A \cap B$ implies $F = A$ or $F = B$.*

Theorem 3.9. *Let X be a bounded commutative BCK-algebra and μ be a fuzzy s-prime filter. Then for each $t \in [0,1]$, μ_t is a prime filter of X whenever $\mu_t \neq \emptyset$ and $\mu_t \neq X$.*

Proof. Let μ be a fuzzy s-prime filter and $t \in [0,1]$, $\mu_t \neq \emptyset$. Then by Lemma 3.6, μ_t is a filter.

Let A, B be two filters such that $\mu_t = A \cap B$. Define the fuzzy subset $\theta = t\chi_A$ and $\sigma = t\chi_B$. It is easy to see that θ and σ are fuzzy filters of X. Note that

$$\theta \cap \sigma(x) = \theta \cdot \sigma(x) = \begin{cases} t, & x \in A \cap B, \\ 0, & x \notin A \cap B. \end{cases} \tag{3.3}$$

Since $\mu_t = A \cap B$, then for any $x \in A \cap B = \mu_t$, $\mu(x) \geq t = \theta \cap \sigma(x)$ and so $\mu(x) \geq \theta \cap \sigma(x)$ for all $x \in X$. Thus $\mu \supseteq \theta \cap \sigma$. It follows from μ being a fuzzy s-prime filter that $\theta \subseteq \mu$ or $\sigma \subseteq \mu$. Without loss of generality let $\theta \subseteq \mu$. Then, for any $x \in A$, $\theta(x) = t\chi_A(x) = t \leq \mu(x)$ and so $x \in \mu(t)$. This means that $A \subseteq \mu_t$. But $\mu_t = A \cap B$ implies $\mu_t \subseteq A$ and thus $\mu_t = A$. Therefore, μ_t is a prime filter by Lemma 3.8. □

Theorem 3.10. *Let X be a bounded commutative BCK-algebra. If μ is a fuzzy s-prime filter, then it is a fuzzy prime filter.*

Proof. The proof follows from Lemma 3.7 and Theorem 3.9. □

In general, the converse of Theorem 3.10 is not true. Let us see the following example.

Example 3.11. Let $X = \{0,1\}$. Define the operation $*$ on X as follows: $0 * 0 = 0$, $0 * 1 = 0$, $1 * 0 = 1$ and $1 * 1 = 0$. It is easy to see that $\langle X; *, 0\rangle$ is a bounded commutative BCK-algebra. Define a fuzzy subset μ of X by $\mu(0) = 0$, $\mu(1) = 1/2$. Clearly μ is a fuzzy prime filter of X.

Moreover, we define the fuzzy filters σ and θ by $\sigma(x) = 1/2$ for all $x \in X$ and $\theta(1) = 1$, $\theta(0) = 0$. Then, we get $\theta\sigma \subseteq \mu$ but $\sigma \not\subseteq \mu$ and $\theta \not\subseteq \mu$. Therefore, μ is not a s-prime fuzzy filter.

Lemma 3.12. *F is a prime filter of X if and only if χ_F^α is a fuzzy s-prime filter, where $\alpha \in [0,1)$, and χ_F^α is defined by*

$$\chi_F^\alpha(x) = \begin{cases} 1, & x \in F \\ \alpha, & x \notin F. \end{cases} \tag{3.4}$$

Proof. Let F be a prime filter. Then, by Lemma 3.6, we can easily see that χ_F^α is a fuzzy filter.

Let θ, σ be two fuzzy prime filters such that $\theta\sigma \subseteq \chi_F^\alpha$, we will prove $\theta \subseteq \chi_F^\alpha$ or $\sigma \subseteq \chi_F^\alpha$. If it is not true, then there exist $x, y \in X \setminus F$ such that $\theta(x) > \alpha$ and $\sigma(y) > \alpha$. Since F is prime, then $x \vee y \notin F$. Note that $\theta\sigma \subseteq \chi_F^\alpha$, then

$$\alpha < \min\{\theta(x), \sigma(y)\} \leq \min\{\theta(x \vee y), \sigma(x \vee y)\} = \theta \cap \sigma(x \vee y) = \theta\sigma(x \vee y) \leq \chi_F^\alpha(x \vee y). \tag{3.5}$$

Thus, $x \vee y \in F$, a contradiction. It follows that $\theta \subseteq \chi_F^\alpha$ or $\sigma \subseteq \chi_F^\alpha$, and so χ_F^α is a fuzzy s-prime filter.

Conversely, let χ_F^α be a fuzzy s-prime filter. By Theorem 3.10, χ_F^α is also a fuzzy prime filter. Then, by Lemma 3.7, $(\chi_F)_t = F$ is a prime filter, where $\alpha \leq t \leq 1$. □

Corollary 3.13. *F is a prime filter of X if and only if χ_F is a fuzzy s-prime filter.*

Lemma 3.14 (see [13]). *Let X be a bounded implicative BCK-algebra, then $x \wedge x^* = 0$ and $x \vee x^* = 1$.*

Lemma 3.15. *Let μ be a fuzzy filter of a bounded commutative BCK-algebra X. Then, $\mu(0) = \min\{\mu(x), \mu(x^*)\}$ for all $x \in X$.*

Proof. Since μ is a fuzzy filter, we have $\mu(0) \geq \min\{\mu(x * 0)^*, \mu(x)\} = \min\{\mu(x^*), \mu(x)\}$ for all $x \in X$. On the other hand, $\mu(0) \leq \min\{\mu(x^*), \mu(x)\}$, since any fuzzy filter is order preserving. Thus, $\mu(0) = \min\{\mu(x), \mu(x^*)\}$. □

Lemma 3.16. *If μ is a fuzzy filter of a bounded BCK-algebra X, then $\mu_1 = \{x \in X \mid \mu(x) = \mu(1)\}$ is a filter of X and χ_{μ_1} is a fuzzy filter of X.*

Proof. Let μ be a fuzzy filter and take $t = \mu(1)$. Then, $\mu_t = \mu_1$ and so $\mu_t = \mu_1$ is a filter of X by Lemma 3.6. Clearly χ_{μ_1} is a fuzzy filter. □

Lemma 3.17. *Let X be a bounded commutative BCK-algebra and μ be a fuzzy s-prime filter of X. Then $\mu(1) = 1$.*

Proof. Suppose that $\mu(1) < 1$. Since μ is non-constant, there exists $a \in X$ such that $\mu(a) < \mu(1)$. Define fuzzy subset θ and σ of X by

$$\theta(x) = \begin{cases} 1, & \mu(x) = \mu(1), \\ 0, & \text{otherwise.} \end{cases} \tag{3.6}$$

and $\sigma(x) = \mu(1)$ for all $x \in X$. By Lemma 3.16, $\theta(x) = \chi_{\mu_1}$is a fuzzy filter and clearly σ is a fuzzy filter. Note that $\theta(1) = 1 > \mu(1)$ and $\sigma(a) = \mu(1) > \mu(a)$, we get $\theta\sigma \not\subseteq \mu$. But note that

for any $x, y \in X$

$$\begin{aligned}\min\{\theta(x), \sigma(x)\} &\leq \begin{cases}\sigma(y), & x \in \mu_1\\ 0, & x \notin \mu_1\end{cases}\\ &\leq \begin{cases}\mu(1), & x \in \mu_1\\ \mu(x), & x \notin \mu_1\end{cases}\\ &= \mu(x)\\ &\leq \mu(x \vee y).\end{aligned} \tag{3.7}$$

Thus, $\theta\sigma(x) = \sup_{x=y\vee z}\{\min\{\theta(y), \sigma(z)\}\} \leq \sup_{x=y\vee z}\{\mu(y \vee z)\} = \sup\{\mu(x)\} = \mu(x)$ for any $x \in X$, a contradiction. Therefore, $\mu(1) = 1$. □

Lemma 3.18. *Let X be a bounded implicative BCK-algebra and μ be a fuzzy s-prime filter of X. Then for any $x \in X$, $\mu(x) = 1$ or $\mu(x^*) = 1$.*

Proof. By Lemma 3.14, $x \vee x^* = 1$, for all $x \in X$. Since μ is a fuzzy s-prime filter, we get that μ_1 is a prime filter of X by Theorem 3.9. Hence, $x \vee x^* = 1 \in \mu_1$ implies $x \in \mu_1$ or $x^* \in \mu_1$. Therefore, $\mu(x) = \mu(1) = 1$ or $\mu(x^*) = \mu(1) = 1$ by Lemma 3.17. □

Theorem 3.19. *Let X be a bounded implicative BCK-algebra and μ be a fuzzy s-prime filter of X. Then, for $x \in X$, $\mu(x) = \mu(1) = 1$ or $\mu(x) = \mu(0)$.*

Proof. By Lemma 3.14, $x \vee x^* = 1$ and then $\mu(1) = \mu(x \vee x^*) = \mu(x)$ or $\mu(x^*)$ since μ is a fuzzy s-prime filter. By Lemma 3.18, we get $\mu(x) = 1$ or $\mu(x^*) = 1$. If $\mu(x^*) = 1$, then $\mu(0) = \min\{\mu(x), \mu(x^*)\} = \mu(x)$ by Lemma 3.15. If $\mu(x^*) \neq 1$, then $\mu(x) = 1 = \mu(1)$. □

Lemma 3.20. *Let X be a bounded BCK-algebra. Then, a filter F of X is proper if and only if $0 \notin F$.*

Proof. If $0 \notin F$, then clearly F is proper.

Conversely, let F be proper. If $0 \in F$, then for any $x \in X$, $(x^*0^*)^* = (x^*1)^* = 0^* = 1 \in F$ and so $x \in F$. It follows that $F = X$, a contradiction. Therefore, $0 \notin F$. □

Lemma 3.21 (see [13]). *Let X be a bounded commutative BCK-algebra and F be a filter of X. If $x \in X \setminus F$, then there is a prime filter A of X such that $F \subseteq A$ and $x \notin A$.*

Lemma 3.22 (see [13]). *Let X be a bounded implicative BCK-algebra. Then, for any $a \in X$, the filter $\langle a\rangle$, generated by a, is a set of elements x in X satisfying $a \leq x$.*

Lemma 3.23. *Let X be a bounded implicative BCK-algebra and $a \neq 0$. Then, $\langle a\rangle \neq X$.*

Proof. By Lemma 3.22, $0 \notin \langle a\rangle$ and thus $\langle a\rangle \neq X$. □

Lemma 3.24 (see [16]). *For a bounded commutative BCK-algebra X, one gets*

(1) *$x^{**} = x$ for all $x \in X$.*

(2) *$x^* \wedge y^* = (x \vee y)^*$, $x^* \vee y^* = (x \wedge y)^*$ for all $x, y \in X$.*

(3) *$x^*y^* = yx$ for all $x, y \in X$.*

Theorem 3.25. *Let X be a bounded implicative BCK-algebra. Then,*

(i) *if $\beta_1, \beta_2 \in (0,1]$, $\beta = \min\{\beta_1, \beta_2\}$ and $x, y \in X$, then $F(x_{\beta_1}) \cap F(y_{\beta_2}) = F((x \vee y)_\beta)$.*

(ii) *if $\beta \in (\mu(0), 1]$ and $x, y \in X$, then $F(x_\beta) \cup F(y_\beta) = F((x \wedge y)_\beta)$.*

Moreover, $F(x_\beta)$ is both open or closed, where $\mu \in F(X)$.

(iii) *if $F(x_\beta) = F(X)$, where $x \in X$ and $\beta \in (0,1]$, then $x = 0$.*

Proof. (i) If $\mu \in F(x_{\beta_1}) \cap F(y\beta_2)$, then $\mu(x) < \beta_1$ and $\mu(y) < \beta_2$. By Theorem 3.10, μ is a fuzzy prime filter, and then $\mu(x \vee y) \leq \max\{\mu(x), \mu(y)\}$. Since $\mu(x) < \beta_1$ and $\mu(y) < \beta_2$, then $\mu(x) \neq \mu(1)$ and $\mu(y) \neq \mu(1)$. It follows from Theorem 3.19 that $\mu(x) = \mu(0)$ and $\mu(y) = \mu(0)$. Thus, $\mu(x \vee y) \leq \max\{\mu(x), \mu(y)\} = \mu(0) = \min\{\mu(x), \mu(y)\} < \min\{\beta_1, \beta_2\} = \beta$. Therefore, $\mu \in F((x \vee y)_\beta)$.

Conversely, if $\mu \in F(x \vee y)_\beta$, then $\mu(x \vee y) < \beta$ and so $\mu(x \vee y) < \beta_1$, $\mu(x \vee y < \beta_2)$. Note that $x \leq x \vee y$ and $y \leq x \vee y$, we get $\mu(x) \leq \mu(x \vee y) < \beta_1$, and $\mu(y) \leq \mu(x \vee y) < \beta_2$, since μ is order preserving. Thus, $\mu \in F(x_{\beta_1})$ and $\mu \in F(y_{\beta_2})$, or equivalently, $\mu \in F(x_{\beta_1}) \cap F(y_{\beta_2})$.

Therefore, (i) holds.

(ii) Let $\mu \in F(x_\beta) \cup F(y_\beta)$. Then, $\mu(x) < \beta$ or $\mu(y) < \beta$. By Lemma 3.17, we have $\mu(1) = 1 \geq \beta > \mu(x)$, $\mu(1) = 1 \geq \beta > \mu(y)$. Therefore, $x \notin \mu_1$ or $y \notin \mu_1$. On the other hand, by Lemma 3.14, $x \vee x^* = 1 \in \mu_1$, $y \vee y^* = 1 \in \mu_1$. Note that $\mu_1 = \mu_1$ is a prime filter of X by Theorem 3.9. If $x \notin \mu_1$, then $x \vee x^* \in \mu_1$ implies $x^* \in \mu_1$. If $y \notin \mu_1$, then $y \vee y^* \in \mu_1$ implies $y^* \in \mu_1$. Therefore, we get that $x^* \in \mu_1$ or $y^* \in \mu_1$. Note that $x \wedge y \leq x, y$, we get $x^* \leq (x \wedge y)^*$ and $y^* \leq (x \wedge y)^*$. Thus, $\mu(x^*) \leq \mu((x \wedge y)^*)$ and $\mu(y^*) \leq \mu((x \wedge y)^*)$, and so $\max\{\mu(x^*), \mu(y^*)\} \leq \mu((x \wedge y)^*)$. But $\mu(x^*) = \mu(1)$ or $\mu(y^*) = \mu(1)$ implies that $1 = \max\{\mu(x^*), \mu(y^*)\} \leq \mu((x \wedge y)^*)$ or $\mu((x \wedge y)^*) = \mu(1)$. This means, $(x \wedge y)^* \in \mu_1$.

If $x \wedge y \in \mu_1$, then $\mu(0) \geq \min\{\mu(((x \wedge y) * 0)^*), \mu(x \wedge y)\} = \min\{\mu((x \wedge y)^*), \mu(x \wedge y)\} = \mu(1)$. It follows that $\mu(x) = \mu(1)$ for all $x \in X$, a contradiction. Thus, $x \wedge y \notin \mu_1$. By Lemma 3.18, $\mu(x \wedge y) = \mu(0)$. Hence, $\mu(x \wedge y) < \beta$ and so $\mu \in F((x \wedge y)_\beta)$. It follows that $F(x_\beta) \cup F(y_\beta) \subseteq F((x \wedge y)_\beta)$.

Conversely, let $\mu \in F((x \wedge y)_\beta)$. Then, $\mu(x \wedge y) < \beta \leq 1 = \mu(1)$. Thus, $x \wedge y \notin \mu_1$. Since $(x \wedge y) \vee (x \wedge y)^* = 1 \in \mu_1$, then $(x \wedge y)^* \in \mu_1$. By Lemma 3.24, $(x \wedge y)^* = x^* \vee y^*$ and so $x^* \in \mu_1$ or $y^* \in \mu_1$. If $x^* \in \mu_1$ (or $y^* \in \mu_1$), then $\mu(0) \geq \min\{\mu((x*0)^*), \mu(x)\} = \min\{\mu(x^*), \mu(x)\} = \mu(x)$ (or $\mu(0) \geq \mu(y)$). Thus, $x \notin \mu_1$ or $y \notin \mu_1$. It follows that $F((x \wedge y)_\beta) \subseteq F(x_\beta) \cup F(y_\beta)$.

Combining the above arguments, we get $F(x_\beta) \cup F(y_\beta) = F((x \wedge y)_\beta)$.

In order to prove $F(x_\beta)$ is closed, we will show $F(x_\beta) = V((x^*)_\beta)$.

Let $\mu \in F(x_\beta)$. Then, $\mu \in F(\langle x_\beta \rangle)$ by Lemma 3.2. Thus, $\langle x_\beta \rangle \not\subseteq \mu$ and so $\mu(x) < \beta \leq 1 = \mu(1)$. Hence, $x \notin \mu_1$. Note that $x \vee x^* = 1 \in \mu_1$ we get $x^* \in \mu_1$, which implies that $\mu(x^*) = \mu(1) = 1 \geq \beta$ and so $(x^*)_\beta \subseteq \mu$. It follows that $\mu \in V((x^*)_\beta)$ and thus $F(x_\beta) \subseteq V((x^*)_\beta)$.

Conversely, let $\mu \in V((x^*)_\beta)$. Then, $(x^*)_\beta \subseteq \mu$ and so $\mu(x^*) \geq \beta > \mu(0)$. By Theorem 3.19, $\mu(x^*) = \mu(1)$. Note that $\mu(0) \geq \min\{((x * 0)^*), \mu(x)\} = \min\{\mu(x^*), \mu(x)\} = \mu(x)$, we get that $\mu(x) = \mu(0) < \beta$, or equivalently, $\langle x_\beta \rangle \not\subseteq \mu$. Thus, $\mu \in F(x_\beta)$ and so $V((x^*)_\beta) \subseteq F(x_\beta)$.

Combining the above two sides, we get $F(x_\beta) = V((x^*)_\beta)$.

(iii) Let $F(x_\beta) = F(X)$. We claim that $x = 0$. If this is not true, by Lemma 3.23,$\langle x \rangle \neq X$. By Lemma 3.21, there exists a prime filter P of X such that $\langle x \rangle \subseteq P$. On the other hand, by Lemma 3.12, $\chi_P \in F(X) = F(x_\beta)$.

Therefore, $\langle x \rangle \not\subseteq P$, a contradiction. It follows that $x = 0$. □

In general, the converse of Theorem 3.25 (iii) does not hold. Let us see the following counter example.

Example 3.26. Let $X = \{0,1,2,3\}$ and $*$-table and $\vee$-table be given as follows.

$*$	0	1	2	3
0	0	0	0	0
1	1	0	1	0
2	2	2	0	0
3	3	2	1	0

$\vee$	0	1	2	3
0	0	1	2	3
1	1	1	3	3
2	2	3	2	3
3	3	3	3	3

(3.8)

Then $(x; *, 0)$ is a bounded implicative BCK-algebra and 3 is a unit. It is easy to see that $P = \{1,3\}$ is a filter of X. From $\vee$-table, we can see easily that P is prime. So, χ_P is a fuzzy s-prime filter by Lemma 3.12. Let $\beta = 1/2$. Then, $3_\beta \subseteq \mu$ and so $\langle 3_\beta \rangle \subseteq \mu$. Hence, $\mu \notin F(3_\beta)$. Therefore $F(3_\beta) \neq F(X)$.

Theorem 3.27. *Let X be a bounded implicative BCK-algebra and $X_\alpha = \{\mu \in F(X) \mid \mu(0) = \alpha\}$ for $\alpha \in [0,1)$. Then, X_α is a Hausdorff space.*

Proof. Let $\mu, \sigma \in X_\alpha$ and $\mu \neq \sigma$. We claim that $\mu_1 \neq \sigma_1$. Otherwise, if $\mu_1 = \sigma_1$, then for $x \in \mu_1 = \sigma_1$, $\mu(x) = \mu(1) = 1 = \sigma(1) = \sigma(x)$ and for $x \notin \mu_1 = \sigma_1$, $\mu(x) = \mu(0) = \alpha = \sigma(0) = \sigma(x)$ by Theorem 3.19, a contradiction. Thus, $\mu_1 \not\subseteq \sigma_1$ or $\sigma_1 \not\subseteq \mu_1$. Let $\mu_1 \not\subseteq \sigma_1$. Then, $x \in \mu_1 \setminus \sigma_1$ implies $x^* \in \sigma_1$. Moreover, $\mu(0) \geq \min\{\mu(x * 0)^*, \mu(x)\} = \min\{\mu(x^*), \mu(x)\} = \mu(x^*)$ since $\mu(x) = \mu(1) = 1$. Thus $\mu(x^*) = \mu(0) \neq \mu(1)$ and so $x^* \notin \mu_1$. Therefore, $x^* \in \sigma_1 \setminus \mu_1$. Hence,

$$\sigma(x) = \alpha = \mu(x^*), \qquad \mu(x) = 1 = \sigma(x^*). \tag{3.9}$$

Let $t \in (\alpha, 1]$. Then, $(x_t)(x) = t > \sigma(x)$, and so $\sigma \in F(x_t)$. Note that $((x^*)_t)(x^*) = t > \alpha = \mu(x^*)$, we get $\mu \in F((x^*)_t)$. Moreover, we get

$$\begin{aligned} F(x_t) \cap F((x^*)_t) &= F((x \vee x^*)_t) \quad \text{(by Theorem 3.25 (i))} \\ &= F(1_t) \quad \text{(by Lemma 3.14)} \\ &= \emptyset \quad \text{(by Lemma 3.17)}. \end{aligned} \tag{3.10}$$

It follows that X_α is a Hausdorff space. □

Corollary 3.28. *Let X be a bounded implicative BCK-algebra. Then $X_0 = \{\mu \in F(X) \mid \mu(0) = 0\}$ is a Hausdorff space.*

Let X be a bounded commutative BCK-algebra, $L(X)$ be the set of all filters of X, and F-spec(X) stand for all prime filters of X.

For any subset A of X, we define $S(A) = \{P \in F\text{-spec}(X) \mid A \not\subseteq P\}$.

If $A = \{a\}$, we denote $S(\{a\})$ by $S(a)$.

Lemma 3.29. *$S(A) = S(\langle A \rangle)$ and if $A \subseteq B$, then $S(A) \subseteq S(B)$.*

Proof. Since $A \subseteq \langle A \rangle$, then $A \not\subseteq P$ implies $\langle A \rangle \not\subseteq P$. Then, $P \in S(A)$ implies $P \in S(\langle A \rangle)$.

Conversely, if $P \in S(\langle A \rangle)$, then $\langle A \rangle \not\subseteq P$. Hence, $A \not\subseteq P$ since $A \subseteq P$ implies $\langle A \rangle \subseteq P$. Therefore, $P \in S(A)$.

Thus, $S(A) = S(\langle A \rangle)$. Similarly, we can prove that $A \subseteq B$ implies $S(A) \subseteq S(B)$. □

Proposition 3.30. *The family $T(X) = \{S(F) \mid F \in L(X)\}$ forms a topology on F-spec(X).*

Proof. First, we get

$$\begin{aligned} S(1) = S(\langle 1\rangle) &= \{P \in F\text{-spec}(X) \mid \langle 1\rangle \not\subseteq P\} = \emptyset \in T(X), \\ S(X) &= \{P \in F\text{-spec}(X) \mid X \not\subseteq P\} = F\text{-spec}(X) \in T(X). \end{aligned} \tag{3.11}$$

Then, for any family $\{S(F_\alpha)\}_{\alpha\in\Omega}$,

$$\begin{aligned} \bigcup_{\alpha\in\Omega} S(F_\alpha) &= \{P \in F\text{-spec}(X) \mid F_\alpha \not\subseteq P \text{ For some } F_\alpha\} \\ &= \left\{P \in F\text{-spec}(X) \mid \bigcup_{\alpha\in\Omega} F_\alpha \not\subseteq P\right\} \\ &= \left\{P \in F\text{-spec}(X) \mid \left\langle \bigcup_{\alpha\in\Omega} F_\alpha \right\rangle \not\subseteq P\right\} \\ &= S\left(\left\langle \bigcup_{\alpha\in\Omega} F_\alpha \right\rangle\right) \in T(X). \end{aligned} \tag{3.12}$$

Finally,

$$S(F_1) \cap S(F_2) = \{P \in F\text{-spec}(X) \mid F_1 \cap F_2 \not\subseteq P\} = S(F_1 \cap F_2) \in T(X). \tag{3.13}$$

Therefore, $T(X)$ is a topology on F-spec(X). □

Theorem 3.31. *Let X be a bounded implicative BCK-algebra and the map $f : F\text{-spec}(X) \to X_\alpha$ is defined by $f(P) = \chi_P^\alpha$ where χ_P^α is defined in Lemma 3.12. Then, f is a homeomorphism.*

Proof. (a) f is well defined.

By Lemma 3.12, χ_P^α is a fuzzy s-prime filter for any $P \in F$-spec(X). Note that $0 \notin P$, then $\chi_P^\alpha(0) = \alpha$ and so $\chi_P^\alpha \in X_\alpha$. Thus, f is well defined.

(b) Clearly f is injective.

(c) f is surjective.

For any $\mu \in X_\alpha$, by Theorem 3.19, $\mu(x) = \mu(1) = 1$ or $\mu(x) = \mu(0) = \alpha$. Hence, $\mu = \chi_{\mu_1}^\alpha$. By Theorem 3.9, we get μ_1 is a prime filter of X. Thus, $\mu_1 \in F$-spec(X) and so $f(\mu_1) = \chi_{\mu_1}^\alpha = \mu$. It follows that f is surjective.

(d) f is continuous.

Let $F_\alpha(\theta) = F(\theta) \cap X_\alpha$ be an open set of X_α. We will prove that $f^{-1}(F_\alpha(\theta))$ is an open set of F-spec(X). It is sufficient to prove that $f^{-1}(F_\alpha(\theta)) = \bigcup_{\alpha\le t\le 1} S(\theta_t)$, since θ_t is a filter of X by Lemma 3.6.

First, let $P \in \bigcup_{\alpha<t<1} S(\theta_t)$, then there exists some t such that $\alpha < t < 1$ and $P \in S(\theta_t)$. Thus, $\theta_t \not\subseteq P$ and there exists $x \in \theta_t \setminus P$. Hence $\theta(x) \ge t > \alpha = \chi_P^\alpha(x)$. Therefore, $\theta \not\subseteq \chi_P^\alpha$ and so $\chi_P^\alpha \in F_\alpha(\theta)$. This shows that $f(P) \in F_\alpha(\theta)$. It follows that $P \in f^{-1}(F_\alpha(\theta))$.

Conversely, let $P \in f^{-1}(F_\alpha(\theta))$, then $f(P) = \chi_P^\alpha \in F_\alpha(\theta)$. Hence, $\theta \not\subseteq \chi_P^\alpha$, and thus there exists $x \in X$ such that $\chi_P^\alpha < \theta(x)$. Therefore, $x \notin P$ and so $\chi_P^\alpha = \alpha < \theta(x)$. We can take t such that $\alpha < t_1 < \theta(x)$. Then, $x \in \theta_{t_1} \setminus P$. It follows that $\theta_{t1} \not\subseteq P$ and so $P \in S(\theta_{t_1}) \subseteq \bigcup_{\alpha<t<1} S(\theta_t)$.

Combining the above two hands, we get $f^{-1}(F_\alpha(\theta)) = \bigcup_{\alpha<t<1} S(\theta_t)$. So f is continuous.

(e) f^{-1} is continuous. It is sufficient to prove that $f(S(F))$ is an open set of X_α for any $F \in L(X)$.

We will prove that $f(S(F)) = F_\alpha(\chi_F^\alpha)$.

$$\begin{aligned}
\mu \in f(S(F)) &\Longrightarrow \exists P \in S(F) \text{ such that } f(P) = \mu = \chi_P^\alpha \\
&\Longrightarrow F \not\subseteq P, \quad f(P) = \mu = \chi_P^\alpha \\
&\Longrightarrow \exists x \in F \setminus P, \quad \mu = \chi_P^\alpha \\
&\Longrightarrow \chi_P^\alpha < 1 = \chi_F^\alpha(x) \\
&\Longrightarrow \chi_F^\alpha \not\subseteq \chi_P^\alpha = \mu \\
&\Longrightarrow \mu \in F_\alpha(\chi_F^\alpha).
\end{aligned} \tag{3.14}$$

Thus, $f(S(F)) \subseteq F_\alpha(\chi_F^\alpha)$.
Conversely, we get

$$\begin{aligned}
\mu \in F_\alpha(\chi_F^\alpha) &\Longrightarrow \chi_F^\alpha \not\subseteq \mu, \quad \exists P \in F\text{-spec}(X) \text{ such that } f(P) = \mu = \chi_P^\alpha. \\
&\Longrightarrow \exists x \in X \text{ such that } \chi_F^\alpha(x) > \mu(x) = \chi_P^\alpha \\
&\Longrightarrow F \not\subseteq P \\
&\Longrightarrow P \in S(F) \\
&\Longrightarrow \mu = f(P) \in f(S(F)).
\end{aligned} \tag{3.15}$$

So that $F_\alpha(\chi_F^\alpha) \subseteq f(S(F))$.
Therefore, $f(S(F)) = F_\alpha(\chi_F^\alpha)$. By Lemma 3.6, we can easily see that χ_F^α is a fuzzy filter of X and so $F_\alpha(\chi_F^\alpha) = F(\chi_F^\alpha) \cap X_\alpha$ is an open set of X_α. It follows that f^{-1} is continuous. □

By Theorem 3.27 and Theorem 3.31, we get the following corollary.

Corollary 3.32. *Let X be a bounded implicative BCK-algebra. Then, F-spec(X) is a Hausdorff space.*

Acknowledgment

The paper is supported by National Nature Science Foundation of Shaanxi Province (2007A19).

References

[1] K. Iséki, "An algebra related with a propositional calculus," *Proceedings of the Japan Academy*, vol. 42, pp. 26–29, 1966.

[2] K. Iséki and S. Tanaka, "Ideal theory of BCK-algebras," *Mathematica Japonica*, vol. 21, no. 4, pp. 351–366, 1976.

[3] K. Iséki and S. Tanaka, "An introduction to the theory of BCK-algebras," *Mathematica Japonica*, vol. 23, no. 1, pp. 1–26, 1978.

[4] J. Meng and Y. B. Jun, *BCK-Algebras*, Kyung Moon Sa Co., Seoul, Republic of Korea, 1994.

[5] L. A. Zadeh, "Fuzzy sets," *Information and Computation*, vol. 8, pp. 338–353, 1965.

[6] C. S. Hoo, "Fuzzy ideals of BCI and MV-algebras," *Fuzzy Sets and Systems*, vol. 62, no. 1, pp. 111–114, 1994.

[7] A. Rosenfeld, "Fuzzy groups," *Journal of Mathematical Analysis and Applications*, vol. 35, pp. 512–517, 1971.

[8] X. Ougen, "Fuzzy BCK-algebra," *Mathematica Japonica*, vol. 36, no. 5, pp. 935–942, 1991.

[9] Y. B. Jun, S. M. Hong, and J. Meng, "Fuzzy BCK-filters," *Mathematica Japonica*, vol. 47, no. 1, pp. 45–49, 1998.

[10] Y. B. Jun, J. Meng, and X. L. Xin, "On fuzzy BCK-filters," *The Korean Journal of Computational & Applied Mathematics*, vol. 5, no. 1, pp. 91–97, 1998.

[11] J. Meng, Y. B. Jun, and H. S. Kim, "Fuzzy implicative ideals of BCK-algebras," *Fuzzy Sets and Systems*, vol. 89, no. 2, pp. 243–248, 1997.

[12] J. Meng, X. L. Xin, and Y. S. Pu, "Quotient BCK-algebra induced by a fuzzy ideal," *Southeast Asian Bulletin of Mathematics*, vol. 23, no. 2, pp. 243–251, 1999.

[13] J. Meng, "BCK-filters," *Mathematica Japonica*, vol. 44, no. 1, pp. 119–129, 1996.

[14] R. A. Alò and E. Y. Deeba, "Topologies of BCK-algebras," *Mathematica Japonica*, vol. 31, no. 6, pp. 841–853, 1986.

[15] Y. B. Jun, X. L. Xin, and D. S. Lee, "On topological BCI-algebras," *Information Sciences*, vol. 116, no. 2-4, pp. 253–261, 1999.

[16] B.-L. Meng, "Some results of fuzzy BCK-filters," *Information Sciences*, vol. 130, no. 1–4, pp. 185–194, 2000.



























Hindawi
Submit your manuscripts at
http://www.hindawi.com