Research Article

Bernoulli Basis and the Product of Several Bernoulli Polynomials

Dae San Kim¹ and Taekyun Kim²

¹ Department of Mathematics, Sogang University, Seoul 121-742, Republic of Korea

² Division of General Education, Kwangjuon University, Seoul 139-701, Republic of Korea

Correspondence should be addressed to Dae San Kim, dskim@sogang.ac.kr

Received 25 June 2012; Accepted 9 August 2012

Academic Editor: Yilmaz Simsek

Copyright © 2012 D. S. Kim and T. Kim. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We develop methods for computing the product of several Bernoulli and Euler polynomials by using Bernoulli basis for the vector space of polynomials of degree less than or equal to n .

1. Introduction

It is well known that, the n th Bernoulli and Euler numbers are defined by

$$\sum_{l=0}^n \binom{n}{l} B_l - B_n = \delta_{1,n}, \quad \sum_{l=0}^n \binom{n}{l} E_l + E_n = 2\delta_{0,n}, \quad (1.1)$$

where $B_0 = E_0 = 1$ and $\delta_{k,n}$ is the Kronecker symbol (see [1–20]).

The Bernoulli and Euler polynomials are also defined by

$$B_n(x) = \sum_{l=0}^n \binom{n}{l} B_{n-l} x^l, \quad E_n(x) = \sum_{l=0}^n \binom{n}{l} E_{n-l} x^l. \quad (1.2)$$

Note that $\{B_0(x), B_1(x), \dots, B_n(x)\}$ forms a basis for the space $\mathbb{P}_n = \{p(x) \in \mathbb{Q}[x] \mid \deg p(x) \leq n\}$.

So, for a given $p(x) \in \mathbb{P}_n$, we can write

$$p(x) = \sum_{k=0}^n a_k B_k(x), \quad (1.3)$$

(see [8–12]) for uniquely determined $a_k \in \mathbb{Q}$.

Further,

$$\begin{aligned} a_k &= \frac{1}{k!} \left\{ p^{(k-1)}(1) - p^{(k-1)}(0) \right\}, \quad \text{where } p^{(k)}(x) = \frac{d^k p(x)}{dx^k}, \\ a_0 &= \int_0^1 p(t) dt, \quad \text{where } k = 1, 2, \dots, n. \end{aligned} \quad (1.4)$$

Probably, $\{1, x, \dots, x^n\}$ is the most natural basis for the space $\mathbb{P}_n$. But $\{B_0(x), B_1(x), \dots, B_n(x)\}$ is also a good basis for the space $\mathbb{P}_n$, for our purpose of arithmetical and combinatorial applications.

What are common to $B_n(x)$, $E_n(x)$, x^n ? A few proportion common to them are as follows:

- (i) they are all monic polynomials of degree n with rational coefficients;
- (ii) $(B_n(x))' = nB_{n-1}(x)$, $(E_n(x))' = nE_{n-1}(x)$, $(x^n)' = nx^{n-1}$;
- (iii) $\int B_n(x) dx = B_{n+1}(x)/(n+1) + c$, $\int E_n(x) dx = E_{n+1}(x)/(n+1) + c$, $\int x^n dx = x^{n+1}/(n+1) + c$.

In [5, 6], Carlitz introduced the identities of the product of several Bernoulli polynomials:

$$\begin{aligned} B_m(x)B_n(x) &= \sum_{r=0}^{\infty} \left\{ \binom{m}{2r}n + \binom{n}{2r}m \right\} \frac{B_{2r}B_{m+n-2r}(x)}{m+n-2r} + (-1)^{m+1} \\ &\quad \times \frac{m!n!}{(m+n)!} B_{m+n} \quad (m+n \geq 2), \\ \int_0^1 B_m(x)B_n(x)B_p(x)B_q(x) dx &= (-1)^{m+n+p+q} \sum_{r,s=0}^{\infty} \left\{ \binom{m}{2r}n + \binom{n}{2r}m \right\} \left\{ \binom{p}{2s}q + \binom{q}{2s}p \right\} \\ &\quad \times \frac{(m+n-2r-1)!(p+q-2s-1)!}{(m+n+p+q-2r-2s)!} B_r B_s B_{m+n+p+q-2r-2} \\ &\quad + (-1)^{m+p} \frac{m!n!}{(m+n)!} \frac{p!q!}{(p+q)!} B_{m+n} B_{p+q}. \end{aligned} \quad (1.5)$$

In this paper, we will use (1.4) to derive the identities of the product of several Bernoulli and Euler polynomials.

2. The Product of Several Bernoulli and Euler Polynomials

Let us consider the following polynomials of degree n :

$$p(x) = \sum_{i_1+\dots+i_r+j_1+\dots+j_s=n} B_{i_1}(x) \cdots B_{i_r}(x) E_{j_1}(x) \cdots E_{j_s}(x), \quad (2.1)$$

where the sum runs over all nonnegative integers $i_1, \dots, i_r, j_1, \dots, j_s$ satisfying $i_1 + \dots + i_r + j_1 + \dots + j_s = n$, $r + s = 1$, $r, s \geq 0$.

Thus, from (2.1), we have

$$\begin{aligned} p^{(k)}(x) &= (n+r+s-1)(n+r+s-2) \cdots (n+r+s-k) \\ &\times \sum_{i_1+\dots+i_r+j_1+\dots+j_s=n-k}^{\infty} B_{i_1}(x) \cdots B_{i_r}(x) E_{j_1}(x) \cdots E_{j_s}(x). \end{aligned} \quad (2.2)$$

For $k = 1, 2, \dots, n$, by (1.4), we get

$$\begin{aligned} a_k &= \frac{1}{k!} \{p^{(k-1)}(1) - p^{(k-1)}(0)\} \\ &= \frac{\binom{n+r+s}{k}}{n+r+s} \sum_{i_1+\dots+i_r+j_1+\dots+j_s=n-k+1} \{B_{i_1}(1) \cdots B_{i_r}(1) E_{j_1}(1) \cdots E_{j_s}(1) - B_{i_1} \cdots B_{i_r} E_{j_1} \cdots E_{j_s}\} \\ &= \frac{\binom{n+r+s}{k}}{n+r+s} \left\{ \sum_{\substack{0 \leq a \leq r \\ 0 \leq c \leq s \\ k+r-n-1 \leq a \leq r}} \binom{r}{a} \binom{s}{c} (-1)^c 2^{s-c} \right. \\ &\quad \times \sum_{i_1+\dots+i_r+j_1+\dots+j_s=n+a+1-k-r}^{\infty} B_{i_1} \cdots B_{i_a} E_{j_1} \cdots E_{j_c} \\ &\quad \left. - \sum_{i_1+\dots+i_r+j_1+\dots+j_s=n-k+1} B_{i_1} \cdots B_{i_r} E_{j_1} \cdots E_{j_s} \right\}. \end{aligned} \quad (2.3)$$

From (2.3), we note that

$$\begin{aligned} a_n &= \frac{\binom{n+r+s}{n}}{n+r+s} \sum_{i_1+\dots+i_r+j_1+\dots+j_s=1} \{B_{i_1}(1) \cdots B_{i_r}(1) E_{j_1}(1) \cdots E_{j_s}(1) - B_{i_1} \cdots B_{i_r} E_{j_1} \cdots E_{j_s}\} \\ &= \frac{\binom{n+r+s}{n}}{n+r+s} \left\{ \left(-\frac{1}{2} + 1\right)r - \left(-\frac{1}{2}\right)s - \left(-\frac{1}{2}\right)(r+s) \right\} \\ &= \frac{\binom{n+r+s}{n}}{n+r+s} (r+s) = \binom{n+r+s-1}{n}, \end{aligned}$$

$$\begin{aligned}
a_{n-1} &= \frac{1}{n+r+s} \binom{n+r+s}{n-1} \\
&\times \sum_{i_1+\dots+i_r+j_1+\dots+j_s=2} \{B_{i_1}(1) \cdots B_{i_r}(1) E_{j_1}(1) \cdots E_{j_s}(1) - B_{i_1} \cdots B_{i_r} E_{j_1} \cdots E_{j_s}\} \\
&= \frac{1}{n+r+s} \binom{n+r+s}{n-1} \left\{ \frac{1}{6}r + \frac{1}{2} \frac{1}{2} \binom{r+s}{2} - \frac{1}{6}r - \left(-\frac{1}{2}\right) \left(-\frac{1}{2}\right) \binom{r+s}{2} \right\} = 0, \\
a_0 &= \int_0^1 p(t) dt \\
&= \sum_{i_1+\dots+i_r+j_1+\dots+j_s=n}^{\infty} \sum_{l_1=0}^{i_1} \cdots \sum_{l_r=0}^{i_r} \sum_{p_1=0}^{j_1} \cdots \sum_{p_s=0}^{j_s} \binom{i_1}{l_1} \cdots \binom{i_r}{l_r} \binom{j_1}{p_1} \cdots \binom{j_s}{p_s} \\
&\times \frac{B_{i_1-l_1} \cdots B_{i_r-l_r} E_{j_1-p_1} E_{j_s-p_s}}{l_1 + \cdots + l_r + p_1 + \cdots + p_s + 1}.
\end{aligned} \tag{2.4}$$

Therefore, by (1.3), (2.1), (2.3), and (2.4), we obtain the following theorem.

Theorem 2.1. For $n \in \mathbb{N}$ with $n \geq 2$, we have

$$\begin{aligned}
&\sum_{i_1+\dots+i_r+j_1+\dots+j_s=n} B_{i_1}(x) \cdots B_{i_r}(x) E_{j_1}(x) \cdots E_{j_s}(x) \\
&= \frac{1}{n+r+s} \sum_{k=1}^{n-2} \binom{n+r+s}{k} \\
&\times \left\{ \sum_{\substack{0 \leq a \leq r \\ 0 \leq c \leq s \\ k+r-n-1 \leq a \leq r}} \binom{r}{a} \binom{s}{c} (-1)^c 2^{s-c} \sum_{i_1+\dots+i_a+j_1+\dots+j_c=n+a+1-k-r} B_{i_1} \cdots B_{i_a} E_{j_1} \cdots E_{j_c} \right. \\
&\quad \left. - \sum_{i_1+\dots+i_r+j_1+\dots+j_s=n-k+1}^{\infty} B_{i_1} \cdots B_{i_r} E_{j_1} \cdots E_{j_s} \right\} B_k(x) + \binom{n+r+s-1}{n} B_n(x) \\
&+ \sum_{i_1+\dots+i_r+j_1+\dots+j_s=n} \sum_{l_1=0}^{i_1} \cdots \sum_{l_r=0}^{i_r} \sum_{p_1=0}^{j_1} \cdots \sum_{p_s=0}^{j_s} \binom{i_1}{l_1} \cdots \binom{i_r}{l_r} \binom{j_1}{p_1} \cdots \binom{j_s}{p_s} \\
&\times \frac{B_{i_1-l_1} \cdots B_{i_r-l_r} E_{j_1-p_1} E_{j_s-p_s}}{l_1 + \cdots + l_r + p_1 + \cdots + p_s + 1}.
\end{aligned} \tag{2.5}$$

Let us take the polynomial $p(x)$ of degree n as follows:

$$p(x) = \sum_{i_1+\dots+i_r+j_1+\dots+j_s+t=n} B_{i_1}(x) \cdots B_{i_r}(x) E_{j_1}(x) \cdots E_{j_s}(x) x^t, \quad (2.6)$$

Then, from (2.6), we have

$$\begin{aligned} p^{(k)}(x) &= (n+r+s)(n+r+s-1) \cdots (n+r+s-k+1) \\ &\times \sum_{i_1+\dots+i_r+j_1+\dots+j_s+t=n-k} B_{i_1}(x) \cdots B_{i_r}(x) E_{j_1}(x) \cdots E_{j_s}(x) x^t, \end{aligned} \quad (2.7)$$

By (1.4) and (2.7), we get, for $k = 1, 2, \dots, n$,

$$\begin{aligned} a_k &= \frac{1}{k!} \{p^{(k-1)}(1) - p^{(k-1)}(0)\} \\ &= \frac{1}{n+r+s+1} \binom{n+r+s+1}{k} \\ &\times \sum_{i_1+\dots+i_r+j_1+\dots+j_s+t=n-k+1} \{B_{i_1}(1) \cdots B_{i_r}(1) E_{j_1}(1) \cdots E_{j_s}(1) - B_{i_1} \cdots B_{i_r} E_{j_1} \cdots E_{j_s} 0^t\} \\ &= \frac{\binom{n+r+s+1}{k}}{n+r+s+1} \left\{ \sum_{\substack{0 \leq a \leq r \\ 0 \leq c \leq s \\ k+r-n-1 \leq a \leq r}} \binom{r}{a} \binom{s}{c} (-1)^c 2^{s-c} \right. \\ &\quad \times \sum_{t=0}^{n+a+1-k-r} \sum_{i_1+\dots+i_a+j_1+\dots+j_c=t} B_{i_1} \cdots B_{i_a} E_{j_1} \cdots E_{j_c} \\ &\quad \left. - \sum_{i_1+\dots+i_r+j_1+\dots+j_s=n-k+1} B_{i_1} \cdots B_{i_r} E_{j_1} \cdots E_{j_s} \right\}, \end{aligned} \quad (2.8)$$

Now, we look at a_n and a_{n-1} .

$$\begin{aligned} a_n &= \frac{\binom{n+r+s+1}{n}}{n+r+s+1} \sum_{i_1+\dots+i_r+j_1+\dots+j_s+t=1} \{B_{i_1}(1) \cdots B_{i_r}(1) E_{j_1}(1) \cdots E_{j_s}(1) - B_{i_1} \cdots B_{i_r} E_{j_1} \cdots E_{j_s} 0^t\} \\ &= \frac{\binom{n+r+s+1}{n}}{n+r+s+1} \left\{ \frac{1}{2}(r+s) + 1 - \left(-\frac{1}{2}\right)(r+s) \right\} \\ &= \frac{r+s+1}{n+r+s+1} \binom{n+r+s+1}{n} = \binom{n+r+s}{n}, \end{aligned}$$

$$\begin{aligned}
a_{n-1} &= \frac{\binom{n+r+s+1}{n-1}}{n+r+s+1} \sum_{i_1+\dots+i_r+j_1+\dots+j_s+t=2} \{B_{i_1}(1) \cdots B_{i_r}(1)E_{j_1}(1) \cdots E_{j_s}(1) - B_{i_1} \cdots B_{i_r}E_{j_1} \cdots E_{j_s}0^t\} \\
&= \frac{\binom{n+r+s+1}{n-1}}{n+r+s+1} \left\{ \frac{1}{6}r+1 + \frac{1}{2} \frac{1}{2} \binom{r+s}{2} + \frac{1}{2}(r+s) - \frac{1}{6}r - \left(-\frac{1}{2}\right) \left(-\frac{1}{2}\right) \binom{r+s}{2} \right\} \\
&= \frac{1}{n+r+s+1} \binom{n+r+s+1}{n-1} \frac{r+s+2}{2} = \frac{1}{2} \binom{n+r+s}{n-1},
\end{aligned} \tag{2.9}$$

From (2.6), we note that

$$\begin{aligned}
a_0 &= \int_0^1 p(t)dt \\
&= \sum_{i_1+\dots+i_r+j_1+\dots+j_s+t=n} \sum_{l_1=0}^{i_1} \cdots \sum_{l_r=0}^{i_r} \sum_{p_1=0}^{j_1} \cdots \sum_{p_s=0}^{j_s} \binom{i_1}{l_1} \cdots \binom{i_r}{l_r} \binom{j_1}{p_1} \cdots \binom{j_s}{p_s} \\
&\quad \times B_{i_1-l_1} \cdots B_{i_r-l_r} E_{j_1-p_1} E_{j_s-p_s} \frac{1}{l_1 + \dots + l_r + p_1 + \dots + p_s + t + 1}.
\end{aligned} \tag{2.10}$$

Therefore, by (1.3), (2.6), (2.8), (2.9), and (2.10), we obtain the following theorem.

Theorem 2.2. For $n \in \mathbb{N}$ with $n \geq 2$, one has

$$\begin{aligned}
&\sum_{i_1+\dots+i_r+j_1+\dots+j_s+t=n} B_{i_1}(x) \cdots B_{i_r}(x)E_{j_1}(x) \cdots E_{j_s}(x)x^t \\
&= \frac{1}{n+r+s+1} \sum_{k=1}^{n-2} \binom{n+r+s+1}{k} \\
&\quad \times \left\{ \sum_{\substack{0 \leq a \leq r \\ 0 \leq c \leq s \\ k+r-n-1 \leq a \leq r}} \binom{r}{a} \binom{s}{c} (-1)^c 2^{s-c} \sum_{t=0}^{n+a+1-k-r} \sum_{i_1+\dots+i_a+j_1+\dots+j_c=t}^{\infty} B_{i_1} \cdots B_{i_a} E_{j_1} \cdots E_{j_c} \right. \\
&\quad \left. - \sum_{i_1+\dots+i_r+j_1+\dots+j_s=n-k+1} B_{i_1} \cdots B_{i_r} E_{j_1} \cdots E_{j_s} \right\} B_k(x) \\
&+ \frac{1}{2} \binom{n+r+s}{n-1} B_{n-1}(x) + \binom{n+r+s}{n} B_n(x) \\
&+ \sum_{i_1+\dots+i_r+j_1+\dots+j_s+t=n} \sum_{l_1=0}^{i_1} \cdots \sum_{l_r=0}^{i_r} \sum_{p_1=0}^{j_1} \cdots \sum_{p_s=0}^{j_s} \left\{ \binom{i_1}{l_1} \cdots \binom{i_r}{l_r} \binom{j_1}{p_1} \cdots \binom{j_s}{p_s} \right. \\
&\quad \times B_{i_1-l_1} \cdots B_{i_r-l_r} E_{j_1-p_1} \cdots E_{j_s-p_s} \\
&\quad \left. \times \frac{1}{l_1 + \dots + l_r + p_1 + \dots + p_s + t + 1} \right\}.
\end{aligned} \tag{2.11}$$

Consider the following polynomial of degree n :

$$p(x) = \sum_{i_1+\dots+i_r+j_1+\dots+j_s=n}^{\infty} \frac{1}{i_1!i_2!\dots i_r!j_1!\dots j_s!} B_{i_1}(x) \cdots B_{i_r}(x) E_{j_1}(x) \cdots E_{j_s}(x). \quad (2.12)$$

Then, from (2.12), one has

$$p^{(k)}(x) = (r+s)^k \sum_{i_1+\dots+i_r+j_1+\dots+j_s=n-k} \frac{B_{i_1}(x) \cdots B_{i_r}(x) E_{j_1}(x) \cdots E_{j_s}(x)}{i_1!i_2!\dots i_r!j_1!\dots j_s!}. \quad (2.13)$$

By (1.4) and (2.13), one gets, for $k = 1, 2, \dots, n$,

$$\begin{aligned} a_k &= \frac{1}{k!} \left\{ p^{(k-1)}(1) - p^{(k-1)}(0) \right\} \\ &= \frac{(r+s)^{k-1}}{k!} \sum_{i_1+\dots+i_r+j_1+\dots+j_s+t=n-k+1} \left\{ \frac{B_{i_1}(1) \cdots B_{i_r}(1) E_{j_1}(1) \cdots E_{j_s}(1) - B_{i_1} \cdots B_{i_r} E_{j_1} \cdots E_{j_s}}{i_1!i_2!\dots i_r!j_1!\dots j_s!} \right\} \\ &= \frac{(r+s)^{k-1}}{k!} \left\{ \sum_{\substack{0 \leq a \leq r \\ 0 \leq c \leq s \\ k+r-n-1 \leq a \leq r}} \binom{r}{a} \binom{s}{c} (-1)^c 2^{s-c} \sum_{i_1+\dots+i_a+j_1+\dots+j_c=n+a+1-k-r} \frac{B_{i_1} \cdots B_{i_a} E_{j_1} \cdots E_{j_c}}{i_1!i_2!\dots i_a!j_1!\dots j_c!} \right. \\ &\quad \left. - \sum_{i_1+\dots+i_r+j_1+\dots+j_s=n-k+1} \frac{1}{i_1!i_2!\dots i_r!j_1!\dots j_s!} B_{i_1} \cdots B_{i_r} E_{j_1} \cdots E_{j_s} \right\}. \end{aligned} \quad (2.14)$$

Now look at a_n and a_{n-1} :

$$\begin{aligned} a_n &= \frac{(r+s)^{n-1}}{n!} \sum_{i_1+\dots+i_r+j_1+\dots+j_s=1} \left\{ \frac{B_{i_1}(1) \cdots B_{i_r}(1) E_{j_1}(1) \cdots E_{j_s}(1) - B_{i_1} \cdots B_{i_r} E_{j_1} \cdots E_{j_s}}{i_1!i_2!\dots i_r!j_1!\dots j_s!} \right\} \\ &= \frac{(r+s)^{n-1}}{n!} \left\{ \frac{1}{2}(r+s) - \left(-\frac{1}{2}\right)(r+s) \right\} = \frac{(r+s)^n}{n!}, \\ a_{n-1} &= \frac{(r+s)^{n-2}}{(n-1)!} \sum_{i_1+\dots+i_r+j_1+\dots+j_s=2} \left\{ \frac{B_{i_1}(1) \cdots B_{i_r}(1) E_{j_1}(1) \cdots E_{j_s}(1) - B_{i_1} \cdots B_{i_r} E_{j_1} \cdots E_{j_s}}{i_1!i_2!\dots i_r!j_1!\dots j_s!} \right\} \\ &= \frac{(r+s)^{n-2}}{(n-1)!} \left\{ \frac{1}{2} \frac{1}{6} r + \frac{1}{2} \frac{1}{2} \binom{r+s}{2} - \frac{1}{2} \frac{1}{6} r - \left(\frac{1}{2}\right) \left(-\frac{1}{2}\right) \binom{r+s}{2} \right\} = 0. \end{aligned} \quad (2.15)$$

It is easy to show that

$$a_0 = \int_0^1 p(t) dt = \sum_{i_1+\dots+i_r+j_1+\dots+j_s=n} \frac{1}{i_1! \dots i_r! j_1! \dots j_s!} \quad (2.16)$$

$$\times \sum_{l_1=0}^{i_1} \dots \sum_{l_r=0}^{i_r} \sum_{p_1=0}^{j_1} \dots \sum_{p_s=0}^{j_s} \left\{ \frac{B_{i_1-l_1} \dots B_{i_r-l_r} E_{j_1-p_1} E_{j_s-p_s}}{l_1 + \dots + l_r + p_1 + \dots + p_s + 1} \binom{i_1}{l_1} \dots \binom{i_r}{l_r} \binom{j_1}{p_1} \dots \binom{j_s}{p_s} \right\}.$$

Therefore, by (1.3), (2.14), and (2.15), one obtains the following theorem.

Theorem 2.3. For $n \in \mathbb{N}$ with $n \geq 2$, one has

$$\begin{aligned} & \sum_{i_1+\dots+i_r+j_1+\dots+j_s=n} \frac{1}{i_1! i_2! \dots i_r! j_1! \dots j_s!} B_{i_1}(x) \dots B_{i_r}(x) E_{j_1}(x) \dots E_{j_s}(x) \\ &= \sum_{k=1}^{n-2} \frac{(r+s)^{k-1}}{k!} \left\{ \sum_{\substack{0 \leq a \leq r \\ 0 \leq c \leq s \\ k+r-n-1 \leq a \leq r}} \binom{r}{a} \binom{s}{c} (-1)^c 2^{s-c} \right. \\ & \quad \times \sum_{i_1+\dots+i_a+j_1+\dots+j_c=n+a+1-k-r} \frac{B_{i_1} \dots B_{i_a} E_{j_1} \dots E_{j_c}}{i_1! i_2! \dots i_a! j_1! \dots j_c!} \\ & \quad - \sum_{i_1+\dots+i_r+j_1+\dots+j_s=n-k+1} \frac{1}{i_1! i_2! \dots i_r! j_1! \dots j_s!} \\ & \quad \left. \times B_{i_1} \dots B_{i_r} E_{j_1} \dots E_{j_s} \right\} B_k(x) + \frac{(r+s)^n}{n!} B_n(x) \\ & + \sum_{i_1+\dots+i_r+j_1+\dots+j_s=n} \sum_{l_1=0}^{i_1} \dots \sum_{l_r=0}^{i_r} \sum_{p_1=0}^{j_1} \dots \sum_{p_s=0}^{j_s} \binom{i_1}{l_1} \dots \binom{i_r}{l_r} \binom{j_1}{p_1} \dots \binom{j_s}{p_s} \\ & \times \frac{B_{i_1-l_1} \dots B_{i_r-l_r} E_{j_1-p_1} E_{j_s-p_s}}{i_1! i_2! \dots i_r! j_1! \dots j_s! (l_1 + \dots + l_r + p_1 + \dots + p_s + 1)}. \end{aligned} \quad (2.17)$$

Take the polynomial $p(x)$ of degree n as follows:

$$p(x) = \sum_{i_1+\dots+i_r+j_1+\dots+j_s+t=n} \frac{1}{i_1! i_2! \dots i_r! j_1! \dots j_s! t!} B_{i_1}(x) \dots B_{i_r}(x) E_{j_1}(x) \dots E_{j_s}(x) x^t. \quad (2.18)$$

Then, from (2.18), one gets

$$p^{(k)}(x) = (r+s+1)^k \times \sum_{i_1+\dots+i_r+j_1+\dots+j_s+t=n-k} \frac{1}{i_1!i_2!\dots i_r!j_1!\dots j_s!t!} B_{i_1}(x) \dots B_{i_r}(x) E_{j_1}(x) \dots E_{j_s}(x) x^t. \quad (2.19)$$

By (1.4) and (2.19), one gets, for $k = 1, \dots, n$,

$$\begin{aligned} a_k &= \frac{1}{k!} \left\{ p^{(k-1)}(1) - p^{(k-1)}(0) \right\} \\ &= \frac{(r+s+1)^{k-1}}{k!} \sum_{i_1+\dots+i_r+j_1+\dots+j_s+t=n-k+1} \frac{1}{i_1!i_2!\dots i_r!j_1!\dots j_s!t!} \\ &\quad \times \{ B_{i_1}(1) \dots B_{i_r}(1) E_{j_1}(1) \dots E_{j_s}(1) - B_{i_1} \dots B_{i_r} E_{j_1} \dots E_{j_s} 0^t \} \\ &= \frac{(r+s+1)^{k-1}}{k!} \left\{ \sum_{\substack{0 \leq a \leq r \\ 0 \leq c \leq s \\ k+r-n-1 \leq a \leq r}} \binom{r}{a} \binom{s}{c} (-1)^c 2^{s-c} \sum_{t=0}^{n+a+1-k-r} \frac{1}{(n+a+1-k-r-t)!} \right. \\ &\quad \times \sum_{i_1+\dots+i_a+j_1+\dots+j_c=t} \frac{1}{i_1!i_2!\dots i_a!j_1!\dots j_c!} B_{i_1} \dots B_{i_a} E_{j_1} \dots E_{j_c} \\ &\quad \left. - \sum_{i_1+\dots+i_r+j_1+\dots+j_s=n-k+1} B_{i_1} \dots B_{i_r} E_{j_1} \dots E_{j_s} \right\}. \quad (2.20) \end{aligned}$$

Now look at a_n and a_{n-1} :

$$\begin{aligned} a_n &= \frac{(r+s+1)^{n-1}}{n!} \sum_{i_1+\dots+i_r+j_1+\dots+j_s+t=1} \frac{1}{i_1!i_2!\dots i_r!j_1!\dots j_s!t!} \\ &\quad \times \{ B_{i_1}(1) \dots B_{i_r}(1) E_{j_1}(1) \dots E_{j_s}(1) - B_{i_1} \dots B_{i_r} E_{j_1} \dots E_{j_s} 0^t \} \\ &= \frac{(r+s+1)^{n-1}}{n!} \left\{ \frac{1}{2}(r+s) + 1 - \left(-\frac{1}{2} \right) (r+s) \right\} \\ &= \frac{(r+s+1)^{n-1}}{n!} (r+s+1) = \frac{(r+s+1)^n}{n!}, \end{aligned}$$

$$\begin{aligned}
a_{n-1} &= \frac{(r+s+1)^{n-2}}{(n-1)!} \sum_{i_1+\dots+i_r+j_1+\dots+j_s+t=2} \frac{1}{i_1!i_2!\dots i_r!j_1!\dots j_s!t!} \\
&\quad \times \{B_{i_1}(1)\dots B_{i_r}(1)E_{j_1}(1)\dots E_{j_s}(1) - B_{i_1}\dots B_{i_r}E_{j_1}\dots E_{j_s}0^t\} \\
&= \frac{(r+s+1)^{n-2}}{(n-1)!} \left\{ \frac{1}{2}\frac{1}{6}r + \frac{1}{2} + \frac{1}{2}\frac{1}{2}\binom{r+s}{2} + \frac{1}{2}(r+s) - \frac{1}{2}\frac{1}{6}r - \left(-\frac{1}{2}\right)\left(-\frac{1}{2}\right)\binom{r+s}{2} \right\} \\
&= \frac{(r+s+1)^{n-2}}{(n-1)!} \frac{r+s+1}{2} \\
&= \frac{(r+s+1)^{n-1}}{2(n-1)!}.
\end{aligned} \tag{2.21}$$

From (2.18), one can derive the following identity:

$$\begin{aligned}
a_0 &= \int_0^1 p(t) dt \\
&= \sum_{i_1+\dots+i_r+j_1+\dots+j_s+t=n} \frac{1}{i_1!\dots i_r!j_1!\dots j_s!t!} \int_0^1 B_{i_1}(x)\dots B_{i_r}(x)E_{j_1}(x)\dots E_{j_s}(x)x^t dt \\
&= \sum_{i_1+\dots+i_r+j_1+\dots+j_s+t=n} \frac{1}{i_1!\dots i_r!j_1!\dots j_s!t!} \sum_{l_1=0}^{i_1} \dots \sum_{l_r=0}^{i_r} \sum_{p_1=0}^{j_1} \dots \\
&\quad \times \sum_{p_s=0}^{j_s} \binom{i_1}{l_1} \dots \binom{i_r}{l_r} \binom{j_1}{p_1} \dots \binom{j_s}{p_s} B_{i_1-l_1} \dots B_{i_r-l_r} E_{j_1-p_1} E_{j_s-p_s} \frac{1}{l_1+\dots+l_r+p_1+\dots+p_s+t+1}.
\end{aligned} \tag{2.22}$$

Therefore, by (1.3), (2.20), (2.21), and (2.22), one obtains the following theorem.

Theorem 2.4. For $n \in \mathbb{N}$ with $n \geq 2$, one has

$$\begin{aligned}
&\sum_{i_1+\dots+i_r+j_1+\dots+j_s+t=n} \frac{1}{i_1!i_2!\dots i_r!j_1!\dots j_s!t!} B_{i_1}(x)\dots B_{i_r}(x)E_{j_1}(x)\dots E_{j_s}(x)x^t \\
&= \sum_{k=1}^{n-2} \frac{(r+s+1)^{k-1}}{k!} \left\{ \sum_{\substack{0 \leq a \leq r \\ 0 \leq c \leq s \\ k+r-n-1 \leq a \leq r}} \binom{r}{a} \binom{s}{c} (-1)^c 2^{s-c} \sum_{t=0}^{n+a+1-k-r} \frac{1}{(n+a+1-k-r-t)!} \right. \\
&\quad \times \sum_{i_1+\dots+i_a+j_1+\dots+j_c=t} \frac{1}{i_1!i_2!\dots i_a!j_1!\dots j_c!} B_{i_1}\dots B_{i_a}E_{j_1}\dots E_{j_c} \\
&\quad \left. - \sum_{i_1+\dots+i_r+j_1+\dots+j_s=n-k+1} \frac{B_{i_1}\dots B_{i_r}E_{j_1}\dots E_{j_s}}{i_1!i_2!\dots i_r!j_1!\dots j_s!} \right\} B_k(x)
\end{aligned}$$

$$\begin{aligned}
& + \frac{(r+s+1)^{n-1}}{2(n-1)!} B_{n-1}(x) + \frac{(r+s+1)^n}{n!} \sum_{i_1+\dots+i_r+j_1+\dots+j_s+t=n} \sum_{l_1=0}^{i_1} \dots \sum_{l_r=0}^{i_r} \sum_{p_1=0}^{j_1} \dots \sum_{p_s=0}^{j_s} \\
& \times \frac{\binom{i_1}{l_1} \dots \binom{i_r}{l_r} \binom{j_1}{p_1} \dots \binom{j_s}{p_s}}{i_1! \dots i_r! j_1! \dots j_s! t!} B_{i_1-l_1} \dots B_{i_r-l_r} E_{j_1-p_1} E_{j_s-p_s} \frac{1}{l_1 + \dots + l_r + p_1 + \dots + p_s + t + 1}.
\end{aligned} \tag{2.23}$$

Acknowledgments

This research was supported by Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education, Science and Technology 2012R1A1A2003786.

References

- [1] S. Araci, D. Erdal, and J. J. Seo, "A study on the fermionic p -adic q -integral representation on $\mathbb{Z}_p$ associated with weighted q -Bernstein and q -Genocchi polynomials," *Abstract and Applied Analysis*, vol. 2011, Article ID 649248, 10 pages, 2011.
- [2] A. Bayad, "Modular properties of elliptic Bernoulli and Euler functions," *Advanced Studies in Contemporary Mathematics*, vol. 20, no. 3, pp. 389–401, 2010.
- [3] A. Bayad and T. Kim, "Identities involving values of Bernstein, q -Bernoulli, and q -Euler polynomials," *Russian Journal of Mathematical Physics*, vol. 18, no. 2, pp. 133–143, 2011.
- [4] L. Carlitz, "Note on the integral of the product of several Bernoulli polynomials.," *Journal of the London Mathematical Society. Second Series*, vol. 34, pp. 361–363, 1959.
- [5] L. Carlitz, "Multiplication formulas for products of Bernoulli and Euler polynomials," *Pacific Journal of Mathematics*, vol. 9, pp. 661–666, 1959.
- [6] L. Carlitz, "Arithmetic properties of generalized Bernoulli numbers," *Journal für die Reine und Angewandte Mathematik*, vol. 202, pp. 174–182, 1959.
- [7] N. S. Jung, H. Y. Lee, and C. S. Ryoo, "Some relations between twisted (h, q) -Euler numbers with weight α and q -Bernstein polynomials with weight α ," *Discrete Dynamics in Nature and Society*, vol. 2011, Article ID 176296, 11 pages, 2011.
- [8] D. S. Kim, "Identities of symmetry for q -Euler polynomials," *Open Journal of Discrete Mathematics*, vol. 1, no. 1, pp. 22–31, 2011.
- [9] D. S. Kim, "Identities of symmetry for generalized Euler polynomials," *International Journal of Combinatorics*, vol. 2011, Article ID 432738, 12 pages, 2011.
- [10] T. Kim, "On the weighted q -Bernoulli numbers and polynomials," *Advanced Studies in Contemporary Mathematics*, vol. 21, no. 2, pp. 207–215, 2011.
- [11] T. Kim, "Symmetry of power sum polynomials and multivariate fermionic p -adic invariant integral on $\mathbb{Z}_p$," *Russian Journal of Mathematical Physics*, vol. 16, no. 1, pp. 93–96, 2009.
- [12] T. Kim, "Some identities on the q -Euler polynomials of higher order and q -Stirling numbers by the fermionic p -adic integral on $\mathbb{Z}_p$," *Russian Journal of Mathematical Physics*, vol. 16, no. 4, pp. 484–491, 2009.
- [13] B. Kurt and Y. Simsek, "Notes on generalization of the Bernoulli type polynomials," *Applied Mathematics and Computation*, vol. 218, no. 3, pp. 906–911, 2011.
- [14] H. Y. Lee, N. S. Jung, and C. S. Ryoo, "A note on the q -Euler numbers and polynomials with weak weight α ," *Journal of Applied Mathematics*, vol. 2011, Article ID 497409, 14 pages, 2011.
- [15] H. Ozden, " p -adic distribution of the unification of the Bernoulli, Euler and Genocchi polynomials," *Applied Mathematics and Computation*, vol. 218, no. 3, pp. 970–973, 2011.
- [16] H. Ozden, I. N. Cangul, and Y. Simsek, "On the behavior of two variable twisted p -adic Euler q - l -functions," *Nonlinear Analysis. Theory, Methods & Applications*, vol. 71, no. 12, pp. e942–e951, 2009.

- [17] S.-H. Rim, A. Bayad, E.-J. Moon, J.-H. Jin, and S.-J. Lee, "A new construction on the q -Bernoulli polynomials," *Advances in Difference Equations*, vol. 2011, article 34, 2011.
- [18] C. S. Ryoo, "Some relations between twisted q -Euler numbers and Bernstein polynomials," *Advanced Studies in Contemporary Mathematics*, vol. 21, no. 2, pp. 217–223, 2011.
- [19] Y. Simsek, "Complete sum of products of (h, q) -extension of Euler polynomials and numbers," *Journal of Difference Equations and Applications*, vol. 16, no. 11, pp. 1331–1348, 2010.
- [20] Y. Simsek, "Generating functions of the twisted Bernoulli numbers and polynomials associated with their interpolation functions," *Advanced Studies in Contemporary Mathematics*, vol. 16, no. 2, pp. 251–278, 2008.