Research Article

Fredholm Weighted Composition Operators on Dirichlet Space

Liankuo Zhao

School of Mathematics and Computer Science, Shanxi Normal University, Linfen 041004, China

Correspondence should be addressed to Liankuo Zhao, liankuozhao@sina.com

Received 3 June 2012; Accepted 3 August 2012

Academic Editor: Henryk Hudzik

Copyright © 2012 Liankuo Zhao. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

This paper characterizes bounded Fredholm, bounded invertible, and unitary weighted composition operators on Dirichlet space.

1. Introduction

Let H be a Hilbert space of analytic functions on the unit disk D . For an analytic function ψ on D , we can define the multiplication operator $M_\psi : f \rightarrow \psi f$, $f \in H$. For an analytic self-mapping φ of D , the composition operator C_φ defined on H as $C_\varphi f = f \circ \varphi$, $f \in H$. These operators are two classes of important operators in the study of operator theory in function spaces [1–3]. Furthermore, for ψ and φ , we define the weighted composition operator $C_{\psi,\varphi}$ on H as

$$C_{\psi,\varphi} : f \longrightarrow \psi(f \circ \varphi), \quad f \in H. \quad (1.1)$$

Recently, the boundedness, compactness, norm, and essential norm of weighted composition operators on various spaces of analytic functions have been studied intensively, see [4–9] and so on. In this paper, we characterize bounded Fredholm weighted composition operators on Dirichlet space of the unit disk.

Recall the Dirichlet space $\mathfrak{D}$ that consists of analytic function f on D with finite Dirichlet integral:

$$D(f) = \int_D |f'|^2 dA < \infty, \quad (1.2)$$

where dA is the normalized Lebesgue area measure on D . It is well known that $\mathfrak{D}$ is the only möbius invariant Hilbert space up to an isomorphism [10]. Endow $\mathfrak{D}$ with norm

$$\|f\| = \left(|f(0)|^2 + D(f) \right)^{1/2}, \quad f \in \mathfrak{D}. \quad (1.3)$$

$\mathfrak{D}$ is a Hilbert space with inner product

$$\langle f, g \rangle = f(0)\overline{g(0)} + \int_D f'(z)\overline{g'(z)}dA(z), \quad f, g \in \mathfrak{D}. \quad (1.4)$$

Furthermore $\mathfrak{D}$ is a reproducing function space with reproducing kernel

$$K_\lambda(z) = 1 + \log \frac{1}{1 - \lambda z}, \quad \lambda, z \in D. \quad (1.5)$$

Denote $\mathcal{M} = \{\psi : \psi \text{ is analytic on } D, \psi f \in \mathfrak{D} \text{ for } f \in \mathfrak{D}\}$. $\mathcal{M}$ is called the multiplier space of $\mathfrak{D}$. By the closed graph theorem, the multiplication operator M_ψ defined by $\psi \in \mathcal{M}$ is bounded on $\mathfrak{D}$. For the characterization of the element in $\mathcal{M}$, see [11].

For analytic function ψ on D and analytic self-mapping φ of D , the weighted composition operator $C_{\psi, \varphi}$ on $\mathfrak{D}$ is not necessarily bounded. Even the composition operator C_φ is not necessarily bounded on $\mathfrak{D}$, which is different from the cases in Hardy space and Bergman space. See [12] for more information about the properties of composition operators acting on the Dirichlet space.

The main result of the paper reads as the following.

Theorem 1.1. *Let ψ and φ be analytic functions on D with $\varphi(D) \subset D$. Then $C_{\psi, \varphi}$ is a bounded Fredholm operator on $\mathfrak{D}$ if and only if $\psi \in \mathcal{M}$, bounded away from zero near the unit circle, and φ is an automorphism of D .*

If $\psi(z) = 1$, then the result above gives the characterization of bounded Fredholm composition operator C_φ on $\mathfrak{D}$, which was obtained in [12].

As corollaries, in the end of this paper one gives the characterization of bounded invertible and unitary weighted composition operator on $\mathfrak{D}$, respectively. Some idea of this paper is derived from [4, 13], which characterize normal and bounded invertible weighted composition operator on the Hardy space, respectively.

2. Proof of the Main Result

In the following, ψ and φ denote analytic functions on D with $\varphi(D) \subset D$. It is easy to verify that $\psi \in \mathfrak{D}$ if $C_{\psi, \varphi}$ is defined on $\mathfrak{D}$.

Proposition 2.1. *Let $C_{\psi,\varphi}$ be a bounded Fredholm operator on $\mathfrak{D}$. Then ψ has at most finite zeroes in D and φ is an inner function.*

Proof. If $C_{\psi,\varphi}$ is a bounded Fredholm operator, then there exist a bounded operator T and a compact operator S on $\mathfrak{D}$ such that

$$T(C_{\psi,\varphi})^* = I + S, \quad (2.1)$$

where I is the identity operator.

Since

$$\begin{aligned} (C_{\psi,\varphi})^* K_w(z) &= \langle C_{\psi,\varphi}^* K_w, K_z \rangle = \langle K_w, C_{\psi,\varphi} K_z \rangle \\ &= \langle K_w, \psi K_z \circ \varphi \rangle = \overline{\psi(w) K_z(\varphi(w))} \\ &= \overline{\psi(w)} K_{\varphi(w)}(z), \end{aligned} \quad (2.2)$$

we have

$$\begin{aligned} \|T\| |\psi(w)| \frac{\|K_{\varphi(w)}\|}{\|K_w\|} &\geq \|T(C_{\psi,\varphi})^* k_w\| \\ &\geq \|k_w\| - \|Sk_w\| \\ &= 1 - \|Sk_w\|, \end{aligned} \quad (2.3)$$

where $k_w = K_w / \|K_w\|$ is the normalization of reproducing kernel function K_w .

Since S is compact and k_w weakly converges to 0 as $|w| \rightarrow 1$, $\|Sk_w\| \rightarrow 0$ as $|w| \rightarrow 1$. It follows that there exists constant r , $0 < r < 1$, such that $\|Sk_w\| < 1/2$ for all w with $r < |w| < 1$. Inequality (2.3) shows that

$$\frac{|\psi(w)|}{\|K_w\|} \geq \frac{1}{2\|T\|\|K_{\varphi(w)}\|}, \quad r < |w| < 1, \quad (2.4)$$

which implies that ψ has no zeroes in $\{w \in D, r < |w| < 1\}$, and, hence, ψ has at most finite zeroes in $\{w \in D, |w| \leq r\}$.

Since k_w weakly converges to 0 as $|w| \rightarrow 1$, $\langle \psi, k_w \rangle \rightarrow 0$ as $|w| \rightarrow 1$, that is,

$$\frac{\psi(w)}{\|K_w\|} \rightarrow 0, \quad |w| \rightarrow 1. \quad (2.5)$$

It follows from (2.4) that $\|K_{\varphi(w)}\| = (1 + \log(1/(1 - |\varphi(w)|^2)))^{1/2} \rightarrow \infty$ and hence $|\varphi(w)| \rightarrow 1$ as $|w| \rightarrow 1$, that is, φ is an inner function. $\square$

For the proof of the following lemma, we cite Carleson's formula for the Dirichlet integral [14].

Let $f \in \mathfrak{D}$, $f = BSF$ be the canonical factorization of f as a function in the Hardy space, where $B = \prod_{j=1}^{\infty} (\bar{a}_j / |a_j|) ((a_j - z) / (1 - \bar{a}_j z))$, is a Blaschke product, S is the singular part of f and F is the outer part of f . Then

$$D(f) = \int_{\mathbb{T}} \sum_{n=1}^{\infty} P_{\alpha_n}(\xi) |f(\xi)|^2 \frac{|d\xi|}{2\pi} + \iint_{\mathbb{T}} \frac{2}{|\zeta - \xi|^2} |f(\xi)|^2 d\mu(\xi) \frac{|d\xi|}{2\pi} \\ + \iint_{\mathbb{T}} \frac{(e^{2u(\zeta)} - e^{2u(\xi)})(u(\zeta) - u(\xi))}{|\zeta - \xi|^2} \frac{|d\zeta|}{2\pi} \frac{|d\xi|}{2\pi}, \quad (2.6)$$

where $\mathbb{T}$ is the unit circle, $u(\xi) = \log |f(\xi)|$, $P_{\alpha}(\xi)$ is the Poisson kernel, and μ is the singular measure corresponding to S .

Lemma 2.2. Let $C_{\psi, \varphi}$ be a bounded operator on $\mathfrak{D}$, $\psi = BF$ with B a finite Blaschke product. Then $C_{F, \varphi}$ is bounded.

Proof. Let M_B be the multiplication operator on $\mathfrak{D}$. Then $C_{\psi, \varphi} = M_B C_{F, \varphi}$. Since B is a finite Blaschke product, by the Carleson's formula, we have

$$D(\psi(f \circ \varphi)) = D(BF(f \circ \varphi)) \geq D(F(f \circ \varphi)), \quad f \in \mathfrak{D}. \quad (2.7)$$

Since $\|f\|^2 = |f(0)|^2 + D(f)$, $f \in \mathfrak{D}$, by the inequality above it is easy to verify that $C_{F, \varphi}$ is bounded on $\mathfrak{D}$ if $C_{\psi, \varphi}$ is bounded. $\square$

Lemma 2.3. Let F be an analytic function on D with zero-free. If $C_{F, \varphi}$ is a bounded Fredholm operator on $\mathfrak{D}$, then φ is univalent.

Proof. If $\varphi(a) = \varphi(b)$ for $a, b \in D$ with $a \neq b$, by a similar reasoning as [1, Lemma 3.26], there exist infinite sets $\{a_n\}$ and $\{b_n\}$ in D which is disjoint such that $\varphi(a_n) = \varphi(b_n)$. Hence,

$$(C_{F, \varphi})^* \left(\frac{K_{a_n}}{F(a_n)} - \frac{K_{b_n}}{F(b_n)} \right) = 0, \quad (2.8)$$

which contradicts to that kernel of $(C_{F, \varphi})^*$ is finite dimensional. $\square$

Corollary 2.4. If $C_{\psi, \varphi}$ is a bounded Fredholm operator on $\mathfrak{D}$, then φ is an automorphism of D and $\psi \in \mathcal{M}$.

Proof. By Proposition 2.1, ψ has the factorization of BF with B a finite Blaschke product and F zero free in D . By Lemma 2.2, $C_{F, \varphi}$ is a bounded operator on $\mathfrak{D}$. Since $C_{\psi, \varphi} = M_B C_{F, \varphi}$ and M_B is a Fredholm operator, $C_{F, \varphi}$ is a Fredholm operator also. By Proposition 2.1 and Lemma 2.3, φ is an univalent inner function, it follows from [1, Corollary 3.28] that φ is an automorphism of D .

Since $C_{\psi, \varphi} C_{\varphi^{-1}} = M_{\psi}$, M_{ψ} is a bounded multiplication operator on $\mathfrak{D}$, which implies that $\psi \in \mathcal{M}$. $\square$

The following lemmas is well-known. It is easy to verify by the fact $M_{\psi}^* K_w = \overline{\psi(w)} K_w$ also.

Lemma 2.5. Let $\psi \in \mathcal{M}$. Then M_ψ is an invertible operator on $\mathfrak{D}$ if and only if ψ is invertible in $\mathcal{M}$.

Lemma 2.6. Let $\psi \in \mathcal{M}$. Then M_ψ is a Fredholm operator on $\mathfrak{D}$ if and only if ψ is bounded away from the unit circle.

Now we give the proof of Theorem 1.1.

Proof of Theorem 1.1. If $C_{\psi,\varphi}$ is a bounded Fredholm operator on $\mathfrak{D}$, by Corollary 2.4, $\psi \in \mathcal{M}$ and φ is an automorphism of D . Since C_φ is invertible, M_ψ is a Fredholm operator. So ψ is bounded away from the unit circle follows from Lemma 2.6.

On the other hand, if $\psi \in \mathcal{M}$ and bounded away from the unit circle, then M_ψ is a bounded Fredholm operator on $\mathfrak{D}$. If φ is an automorphism of D , then C_φ is invertible. Hence $C_{\psi,\varphi} = M_\psi C_\varphi$ is a bounded Fredholm operator on $\mathfrak{D}$. $\square$

As corollaries, in the following, we characterize bounded invertible and unitary weighted composition operators on $\mathfrak{D}$.

Corollary 2.7. Let ψ and φ be analytic functions on D with $\varphi(D) \subset D$. Then $C_{\psi,\varphi}$ is a bounded invertible operator on $\mathfrak{D}$ if and only if $\psi \in \mathcal{M}$, invertible in $\mathcal{M}$, and φ is an automorphism of D .

Proof. Since a bounded invertible operator is a bounded Fredholm operator, the proof is similar to the proof of Theorem 1.1. $\square$

Corollary 2.8. Let ψ and φ be analytic functions on D with $\varphi(D) \subset D$. $C_{\psi,\varphi}$ is a bounded operator on $\mathfrak{D}$. Then $C_{\psi,\varphi}$ is a unitary operator if and only if ψ is a constant with $|\psi| = 1$ and φ is a rotation of D .

Proof. If $C_{\psi,\varphi}$ is a unitary operator, then it must be an invertible operator. By Corollary 2.7, φ is an automorphism of D and ψ is invertible in $\mathcal{M}$.

Let n be nonnegative integer, $e_n(z) = z^n$, $z \in D$. A unitary is also an isometry, so we have

$$\|\psi\| = \|C_{\psi,\varphi}e_0\| = \|e_0\| = 1, \quad (2.9)$$

$$\|\psi\varphi^n\| = \|C_{\psi,\varphi}e_n\| = \|e_n\| = \sqrt{n}, \quad n \geq 1. \quad (2.10)$$

Let $\alpha \in D$ such that $\varphi(\alpha) = 0$. Since φ is an automorphism of D , φ^n is a finite Blaschke product with zero α of order n . By Carleson's formula for Dirichlet integral, we have

$$D(\psi\varphi^n) = n \int_{\mathbb{T}} P_\alpha(\xi) |\psi(\xi)|^2 \frac{|d\xi|}{2\pi} + D(\psi). \quad (2.11)$$

Hence,

$$\begin{aligned} n &= \|\psi\varphi^n\|^2 = |\psi(0)\varphi(0)^n|^2 + D(\psi\varphi^n) \\ &= |\psi(0)\varphi(0)^n|^2 + n \int_{\mathbb{T}} P_\alpha(\xi) |\psi(\xi)|^2 \frac{|d\xi|}{2\pi} + D(\psi), \quad n \geq 1. \end{aligned} \quad (2.12)$$

That is,

$$1 = \frac{|\varphi(0)\varphi(0)^n|^2}{n} + \int_{\mathbb{T}} P_{\alpha}(\xi) |\varphi(\xi)|^2 \frac{|d\xi|}{2\pi} + \frac{D(\varphi)}{n}, \quad n \geq 1. \quad (2.13)$$

Let $n \rightarrow \infty$, then $1 = \int_{\mathbb{T}} P_{\alpha}(\xi) |\varphi(\xi)|^2 (|d\xi|/2\pi)$.

By (2.12), we have $D(\varphi) = 0$ and $|\varphi(0)\varphi(0)| = 0$. By (2.9), we obtain φ is a constant with $|\varphi| = 1$, which implies that $\varphi(0) = 0$, that is, φ is a rotation of D .

The sufficiency is easy to verify. $\square$

Remark 2.9. The key step in the proof of the main result is to analyze zeros of the symbol φ and univalence of φ . The following result pointed out by the referee gives a simple characterization of the symbols φ and φ for the bounded Fredholm operator $C_{\varphi,\varphi}$ on $\mathfrak{D}$.

Proposition 2.10. *Let φ and φ be analytic functions on D with $\varphi(D) \subset D$. $C_{\varphi,\varphi}$ is a bounded Fredholm operator on $\mathfrak{D}$. Then φ has only finitely many zeros in D and φ is univalent.*

Proof. If $\varphi(a) = 0$ for $a \in D$, then $C_{\varphi,\varphi}^* K_a = \overline{\varphi(a)} K_{\varphi(a)} = 0$, which implies that K_a is in the kernel of $C_{\varphi,\varphi}^*$. Thus if φ had infinitely many zeros, the kernel of $C_{\varphi,\varphi}^*$ would be infinite dimensional and hence this operator would not be Fredholm.

If $\varphi(a) = \varphi(b)$ for $a, b \in D$ with $a \neq b$, by a similar reasoning as [1, Lemma 3.26], there exist infinite sets $\{a_n\}$ and $\{b_n\}$ in D which is disjoint such that $\varphi(a_n) = \varphi(b_n)$. Since φ has only finitely many zeros in D , we can choose infinitely many a_n and b_n such that $\varphi(a_n) \neq 0$, $\varphi(b_n) \neq 0$. Hence,

$$(C_{\varphi,\varphi})^* \left(\frac{K_{a_n}}{\varphi(a_n)} - \frac{K_{b_n}}{\varphi(b_n)} \right) = 0. \quad (2.14)$$

Since $C_{\varphi,\varphi}$ is a Fredholm operator, φ must be univalent. $\square$

Acknowledgments

Thanks are for referee for many helpful suggestions which promote the author to think the related issues deeply. This work is supported by YSF of Shanxi (2010021002-2) and NSFC (11201274).

References

- [1] C. C. Cowen and B. D. MacCluer, *Composition Operators on Spaces of Analytic Functions*, Studies in Advanced Mathematics, CRC Press, Boca Raton, Fla, USA, 1995.
- [2] R. G. Douglas, *Banach Algebra Techniques in Operator Theory*, vol. 179 of *Graduate Texts in Mathematics*, Springer, New York, NY, USA, 2nd edition, 1998.
- [3] K. H. Zhu, *Operator Theory in Function Spaces*, vol. 139 of *Monographs and Textbooks in Pure and Applied Mathematics*, Marcel Dekker, New York, NY, USA, 1990.
- [4] P. S. Bourdon and S. K. Narayan, "Normal weighted composition operators on the Hardy space $H^2(U)$," *Journal of Mathematical Analysis and Applications*, vol. 367, no. 1, pp. 278–286, 2010.
- [5] M. D. Contreras and A. G. Hernández-Díaz, "Weighted composition operators on Hardy spaces," *Journal of Mathematical Analysis and Applications*, vol. 263, no. 1, pp. 224–233, 2001.

- [6] M. D. Contreras and A. G. Hernández-Díaz, "Weighted composition operators between different Hardy spaces," *Integral Equations and Operator Theory*, vol. 46, no. 2, pp. 165–188, 2003.
- [7] Ž. Čučković and R. Zhao, "Weighted composition operators on the Bergman space," *Journal of the London Mathematical Society. Second Series*, vol. 70, no. 2, pp. 499–511, 2004.
- [8] Ž. Čučković and R. Zhao, "Weighted composition operators between different weighted Bergman spaces and different Hardy spaces," *Illinois Journal of Mathematics*, vol. 51, no. 2, pp. 479–498, 2007.
- [9] S. Ohno and K. Stroethoff, "Weighted composition operators from reproducing Hilbert spaces to Bloch spaces," *Houston Journal of Mathematics*, vol. 37, no. 2, pp. 537–558, 2011.
- [10] J. Arazy and S. D. Fisher, "The uniqueness of the Dirichlet space among Möbius-invariant Hilbert spaces," *Illinois Journal of Mathematics*, vol. 29, no. 3, pp. 449–462, 1985.
- [11] D. A. Stegenga, "Multipliers of the Dirichlet space," *Illinois Journal of Mathematics*, vol. 24, no. 1, pp. 113–139, 1980.
- [12] M. Jovović and B. MacCluer, "Composition operators on Dirichlet spaces," *Acta Scientiarum Mathematicarum*, vol. 63, no. 1-2, pp. 229–247, 1997.
- [13] G. Gunatillake, "Invertible weighted composition operators," *Journal of Functional Analysis*, vol. 261, no. 3, pp. 831–860, 2011.
- [14] L. Carleson, "A representation formula for the Dirichlet integral," *Mathematische Zeitschrift*, vol. 73, pp. 190–196, 1960.