Hindawi Publishing Corporation
International Journal of Mathematics and Mathematical Sciences
Volume 2012, Article ID 981059, 14 pages
doi:10.1155/2012/981059

Research Article

On Subspaces of an Almost φ-Lagrange Space

P. N. Pandey and Suresh K. Shukla

Department of Mathematics, University of Allahabad, Allahabad 211002, India

Correspondence should be addressed to Suresh K. Shukla, shuklasureshk@gmail.com

Received 29 March 2012; Accepted 4 June 2012

Academic Editor: Zhongmin Shen

We discuss the subspaces of an almost φ-Lagrange space (APL space in short). We obtain the induced nonlinear connection, coefficients of coupling, coefficients of induced tangent and induced normal connections, the Gauss-Weingarten formulae, and the Gauss-Codazzi equations for a subspace of an APL-space. Some consequences of the Gauss-Weingarten formulae have also been discussed.

1. Introduction

The credit for introducing the geometry of Lagrange spaces and their subspaces goes to the famous Romanian geometer Miron [1]. He developed the theory of subspaces of a Lagrange space together with Bejancu [2]. Miron and Anastasiei [3] and Sakaguchi [4] studied the subspaces of generalized Lagrange spaces (GL spaces in short). Antonelli and Hrimiuc [5, 6] introduced the concept of φ-Lagrangians and studied φ-Lagrange manifolds. Generalizing the notion of a φ-Lagrange manifold, the present authors recently studied the geometry of an almost φ-Lagrange space (APL space briefly) and obtained the fundamental entities related to such space [7]. This paper is devoted to the subspaces of an APL space.

Let $F^n = (M, F(x, y))$ be an n-dimensional Finsler space and $\varphi : \mathbb{R}^+ \to \mathbb{R}$ a smooth function. If the function φ has the following properties:

(a) $\varphi'(t) \neq 0$,

(b) $\varphi'(t) + \varphi''(t) \neq 0$, for every $t \in \mathrm{Im}(F^2)$,

then the Lagrangian given by

$$L(x, y) = \varphi\left(F^2\right) + A_i(x)y^i + U(x), \tag{1.1}$$

where $A_i(x)$ is a covector and $U(x)$ is a smooth function, is a regular Lagrangian [7]. The space $L^n = (M, L(x,y))$ is a Lagrange space. The present authors [7] called such space as an almost φ-Lagrange space (shortly APL space) associated to the Finsler space F^n. An APL space reduces to a φ-Lagrange space if and only if $A_i(x) = 0$ and $U(x) = 0$. We take

$$g_{ij} = \frac{1}{2}\dot{\partial}_i\dot{\partial}_j F^2, \qquad a_{ij} = \frac{1}{2}\dot{\partial}_i\dot{\partial}_j L, \quad \text{where } \dot{\partial}_i \equiv \frac{\partial}{\partial y^i}. \tag{1.2}$$

We indicate all the geometrical objects related to F^n by putting a small circle "$\circ$" over them. Equations (1.2), in view of (1.1), provide the following expressions for a_{ij} and its inverse (cf. [7]):

$$a_{ij} = \varphi' \cdot \left(g_{ij} + \frac{2\varphi''}{\varphi'}\overset{\circ}{y}_i\overset{\circ}{y}_j\right), \qquad a^{ij} = \frac{1}{\varphi'}\left(g^{ij} - \frac{2\varphi''}{\varphi' + 2F^2\varphi''}y^i\ y^j\right), \tag{1.3}$$

where $g_{ij}y^j = \overset{\circ}{y}_i$.

Let $\check{M}$ be a smooth manifold of dimension m, $1 < m < n$, immersed in M by immersion $i : \check{M} \to M$. The immersion i induces an immersion $T_i : T\check{M} \to TM$ making the following diagram commutative:

$$\begin{array}{ccc} T\check{M} & \xrightarrow{T_i} & TM \\ \check{\pi}\downarrow & & \downarrow\pi \\ \check{M} & \xrightarrow[i]{} & M. \end{array} \tag{1.4}$$

Let (u^α, v^α) (throughout the paper, the Greek indices $\alpha, \beta, \gamma, \ldots$ run from 1 to m) be local coordinates on $T\check{M}$. The restriction of the Lagrangian L on $T\check{M}$ is $L(u,v) = L(x(u), y(u,v))$. Let $a_{\alpha\beta} = (1/2)(\partial^2\check{L}/\partial u^\alpha\partial u^\beta)$. Then, we have (cf. [8]) $a_{\alpha\beta} = B^i_\alpha B^j_\beta a_{ij}$ where $B^i_\alpha(u) = \partial x^i/\partial u^\alpha$ are the projection factors. The pair $\check{L}^m = (\check{M}, \check{L}(u,v))$ is also a Lagrange space, called the subspace of L^n. For the natural bases $(\partial/\partial x^i, \partial/\partial y^i)$ on TM and $(\partial/\partial u^\alpha, \partial/\partial v^\alpha)$ on $T\check{M}$, we have [8]

$$\frac{\partial}{\partial u^\alpha} = B^i_\alpha\frac{\partial}{\partial x^i} + B^i_{0\alpha}\frac{\partial}{\partial y^i}, \qquad \frac{\partial}{\partial v^\alpha} = B^i_\alpha\frac{\partial}{\partial y^i}, \tag{1.5}$$

where $B^i_{0\alpha} = B^i_{\beta\alpha}v^\beta$, $B^i_{\beta\alpha} = \partial^2 x^i/\partial u^\alpha\partial u^\beta$.

For the bases (dx^i, dy^i) and (du^α, dv^α), we have

$$dx^i = B^i_\alpha du^\alpha, \qquad dy^i = B^i_\alpha dv^\alpha + B^i_{0\alpha}du^\alpha. \tag{1.6}$$

Since (B^i_α) are m linearly independent vector fields tangent to $\check{M}$, a vector field $\xi^i(x,y)$ is normal to $\check{M}$ along $T\check{M}$ if on $T\check{M}$, we have

$$a_{ij}(x(u), y(u,v))B^i_\alpha\xi^j = 0, \quad \forall\alpha = 1, 2, \ldots, m. \tag{1.7}$$

There are, at least locally, $(n-m)$ unit vector fields $B_a^i(u,v)$ $(a = m+1, m+2, \ldots, n)$ normal to $\check{M}$ and mutually orthonormal, that is,

$$a_{ij}B_\alpha^i B_b^j = 0, \quad a_{ij}B_a^i B_b^j = \delta_{ab}, \quad (a,b = m+1, m+2, \ldots, n). \tag{1.8}$$

Thus, at every point $(u,v) \in T\check{M}$, we have a moving frame $\Re = ((u,v), B_\alpha^i(u,v), B_a^i(u,v))$. Using (1.3) in the first expression of (1.8) and keeping $\overset{\circ}{y}_i B_a^i = 0$ (this fact is clear from $g_{ij}y^i B_a^j = 0$) in view, we observe that B_a^i's are normal to $\check{M}$ with respect to L^n if and only if they are so with respect to F^n. The dual frame of $\Re$ is $\Re^* = ((u,v), B_i^\alpha(u,v), B_i^a(u,v))$ with the following duality conditions:

$$B_\alpha^i B_i^\beta = \delta_\alpha^\beta, \qquad B_a^i B_i^\beta = 0, \qquad B_\alpha^i B_i^b = 0, \qquad B_a^i B_i^b = \delta_a^b, \qquad B_a^i B_j^a + B_\alpha^i B_j^\alpha = \delta_j^i. \tag{1.9}$$

We will make use of the following results due to the present authors [7], during further discussion.

Theorem 1.1 (cf. [7]). *The canonical nonlinear connection of an APL space L^n has the local coefficients given by*

$$N_j^i = \overset{\circ}{N}{}_j^i - V_j^i, \tag{1.10}$$

where $V_j^i = (1/2)F_j^i - S_j^{ir}(2F_{rk}y^k + \partial_r U)$,

$$S_j^{ir} = \frac{1}{2\varphi'}\overset{\circ}{C}{}_{qj}^i g^{qr} + \frac{1}{2}\frac{\varphi''}{\varphi'^2}g^{ir}\overset{\circ}{y}_j + \frac{\varphi''\left(\delta_j^r y^i + \delta_j^i y^r\right)}{2\varphi'\left(\varphi' + 2F^2\varphi''\right)} + \frac{\varphi'^2\varphi''' - 2\varphi''^3F^2 - 4\varphi'\varphi''^2}{2\varphi'^2\left(\varphi' + 2F^2\varphi''\right)^2}y^i\overset{\circ}{y}_j y^r,$$
$$F_{rk}(x) = \frac{1}{2}(\partial_r A_k - \partial_k A_r), \qquad F_j^i = a^{ik}F_{kj}. \tag{1.11}$$

Theorem 1.2 (cf. [7]). *The coefficients of the canonical metrical d-connection $C\Gamma(N)$ of an APL space L^n are given by*

$$C_{jk}^i = \overset{\circ}{C}{}_{jk}^i + \frac{\varphi''}{\varphi'}\left(\delta_j^i\overset{\circ}{y}_k + \delta_k^i\overset{\circ}{y}_j\right) + \frac{\varphi''}{\varphi' + 2F^2\varphi''}g_{jk}y^i + \frac{2\left(\varphi'''\varphi' - 2\varphi''^2\right)}{\varphi'\left(\varphi' + 2F^2\varphi''\right)}y^i\overset{\circ}{y}_j\overset{\circ}{y}_k, \tag{1.12}$$

$$L_{jk}^i = \overset{\circ}{L}{}_{jk}^i + V_k^r C_{jr}^i + V_j^r C_{kr}^i + V_p^r a^{ip}C_{rkj}. \tag{1.13}$$

For basic notations related to a Finsler space, a Lagrange space, and their subspaces, we refer to the books [8, 9].

2. Induced Nonlinear Connection

Let $\check{N} = (\check{N}^{\alpha}_{\beta}(u,v))$ be a nonlinear connection for $\check{L}^m = (\check{M}, \check{L}(u,v))$. The adapted basis of $T_{(u,v)}T\check{M}$ induced by $\check{N}$ is $(\delta/\delta u^{\alpha} = \delta_{\alpha}, \partial/\partial v^{\alpha} = \dot{\partial}_{\alpha})$, where

$$\delta_{\alpha} = \partial_{\alpha} - \check{N}^{\beta}_{\alpha}\dot{\partial}_{\beta}. \tag{2.1}$$

The dual basis (cobasis) of the adapted basis $(\delta_{\alpha}, \dot{\partial}_{\alpha})$ is $(du^{\alpha}, \delta v^{\alpha} = dv^{\alpha} + \check{N}^{\alpha}_{\beta}du^{\beta})$.

Definition 2.1 (cf. [8]). A nonlinear connection $\check{N} = (\check{N}^{\alpha}_{\beta}(u,v))$ of $\check{L}^m$ is said to be induced by the canonical nonlinear connection N if the following equation holds good:

$$\delta v^{\alpha} = B^{\alpha}_{i}\delta y^{i}. \tag{2.2}$$

The local coefficients of the induced nonlinear connection $\check{N} = (\check{N}^{\alpha}_{\beta}(u,v))$ for the subspace $\check{L}^m = (\check{M}, \check{L}(u,v))$ of a Lagrange space $L^n = (M, L(x,y))$ are given by (cf. [8])

$$\check{N}^{\alpha}_{\beta} = B^{\alpha}_{i}\left(N^{i}_{j}B^{j}_{\beta} + B^{i}_{0\beta}\right), \tag{2.3}$$

N^{i}_{j} being the local coefficients of canonical nonlinear connection N of the Lagrange space $L^n = (M, L(x,y))$. Now using (1.10) in (2.3), we get

$$\check{N}^{\alpha}_{\beta} = B^{\alpha}_{i}\left(\overset{\circ}{N}{}^{i}_{j}B^{j}_{\beta} + B^{i}_{0\beta}\right) - B^{\alpha}_{i}V^{i}_{j}B^{j}_{\beta}. \tag{2.4}$$

If we take $\overset{\circ}{\check{N}}{}^{\alpha}_{\beta} = B^{\alpha}_{i}(\overset{\circ}{N}{}^{i}_{j}B^{j}_{\beta} + B^{i}_{0\beta})$, it follows from (2.4) that

$$\check{N}^{\alpha}_{\beta} = \overset{\circ}{\check{N}}{}^{\alpha}_{\beta} - B^{\alpha}_{i}V^{i}_{j}B^{j}_{\beta}. \tag{2.5}$$

Thus, we have the following.

Theorem 2.2. *The local coefficients of the induced nonlinear connection $\check{N}$ of the subspace $\check{L}^m$ of an APL space L^n are given by* (2.5).

In view of (2.5), (2.1) takes the following form, for the subspace $\check{L}^m$ of an APL space L^n:

$$\delta_{\beta} = \overset{\circ}{\delta}_{\beta} + B^{\alpha}_{p}V^{p}_{j}B^{j}_{\beta}\dot{\partial}_{\alpha}, \tag{2.6}$$

where $\overset{\circ}{\delta}_{\beta} = \partial_{\beta} - \overset{\circ}{\check{N}}{}^{\alpha}_{\beta}\dot{\partial}_{\alpha}$.

We can put $(dx^i, \delta y^i)$ as (cf. [8])

$$dx^i = B^i_\alpha du^\alpha, \qquad \delta y^i = B^i_\alpha \delta y^\alpha + B^i_a H^a_\alpha du^\alpha, \tag{2.7}$$

where

$$H^a_\alpha = B^a_i \left(N^i_j B^j_\alpha + B^i_{0\alpha} \right). \tag{2.8}$$

Using (1.10) in (2.8) and simplifying, we get

$$H^a_\alpha = B^a_i \left(\overset{\circ}{N}{}^i_j B^j_\alpha + B^i_{0\alpha} \right) - B^a_i V^i_j B^j_\alpha. \tag{2.9}$$

Taking $\overset{\circ}{H}{}^a_\alpha = B^a_i (\overset{\circ}{N}{}^i_j B^j_\alpha + B^i_{0\alpha})$, in (2.9), it follows that

$$H^a_\alpha = \overset{\circ}{H}{}^a_\alpha - B^a_i V^i_j B^j_\alpha. \tag{2.10}$$

Now, $dx^i = B^i_\alpha du^\alpha$, $\delta y^i = B^i_\alpha \delta y^\alpha$ if and only if $H^a_\alpha = 0$, that is, if and only if $\overset{\circ}{H}{}^a_\alpha = B^a_i V^i_j B^j_\alpha$. Thus, we have the following.

Theorem 2.3. *The adapted cobasis $(dx^i, \delta y^i)$ of the basis $(\partial/\partial x^i, \partial/\partial y^i)$ induced by the nonlinear connection N of an APL space L^n is of the form $dx^i = B^i_\alpha du^\alpha, \delta y^i = B^i_\alpha \delta y^\alpha$ if and only if $\overset{\circ}{H}{}^a_\alpha = B^a_i V^i_j B^j_\alpha$.*

Definition 2.4 (cf. [8]). Let $D = D\Gamma(N)$ be the canonical metrical d-connection of L^n. An operator $\check{D}$ is said to be a coupling of D with $\check{N}$ if

$$\check{D} X^i = X^i_{|\alpha} du^\alpha + X^i|_\alpha \delta v^\alpha, \tag{2.11}$$

where $X^i_{|\alpha} = \delta_\alpha X^i + X^j \check{L}^i_{j\alpha}$, $X^i|_\alpha = \dot{\partial}_\alpha X^i + X^j \check{C}^i_{j\alpha}$.

The coefficients $(\check{L}^i_{j\alpha}, \check{C}^i_{j\alpha})$ of coupling $\check{D}$ of D with $\check{N}$ are given by

$$\check{L}^i_{j\alpha} = L^i_{jk} B^k_\alpha + C^i_{jk} B^k_a H^a_\alpha, \tag{2.12}$$

$$\check{C}^i_{j\alpha} = C^i_{jk} B^k_\alpha. \tag{2.13}$$

Using (1.12) and (1.13) in (2.12), we have

$$\begin{aligned}\check{L}^i_{j\beta} = {} & \left(\overset{\circ}{L}{}^i_{jk} + V^r_k C^i_{jr} + V^r_j C^i_{kr} + V^r_p a^{ip} C_{rkj}\right) B^k_\beta \\ & + \left[\overset{\circ}{C}{}^i_{jk} + \frac{\varphi''}{\varphi'}\left(\delta^i_j \overset{\circ}{y}_k + \delta^i_k \overset{\circ}{y}_j\right) + \frac{\varphi''}{\varphi' + 2F^2\varphi''} g_{jk} y^i \right. \\ & \left. + \frac{2\left(\varphi'''\varphi' - 2\varphi''^2\right)}{\varphi'\left(\varphi' + 2F^2\varphi''\right)} y^i \overset{\circ}{y}_j \overset{\circ}{y}_k \right] B^k_a H^a_\beta.\end{aligned} \tag{2.14}$$

In view of (2.10) and $\overset{\circ}{y}_i B^i_a = 0$, (2.14) becomes

$$\begin{aligned}\check{L}^i_{j\beta} = {} & \left(\overset{\circ}{L}{}^i_{jk} B^k_\beta + \overset{\circ}{C}{}^i_{jk} B^k_a \overset{\circ}{H}{}^a_\beta\right) + \left(V^r_k C^i_{jr} + V^r_j C^i_{kr} + V^r_p a^{ip} C_{rkj} - \overset{\circ}{C}{}^i_{jr} B^r_b B^b_p V^p_k\right) B^k_\beta \\ & + \left(\frac{\varphi''}{\varphi'} \overset{\circ}{y}_j \delta^i_k + \frac{\varphi''}{\varphi' + 2F^2\varphi''} g_{jk} y^i\right) B^k_a H^a_\beta,\end{aligned} \tag{2.15}$$

that is,

$$\begin{aligned}\check{L}^i_{j\beta} = {} & \overset{\circ}{\check{L}}{}^i_{j\beta} + \left(V^r_k C^i_{jr} + V^r_j C^i_{kr} + V^r_p a^{ip} C_{rkj} - \overset{\circ}{C}{}^i_{jr} B^r_b B^b_p V^p_k\right) B^k_\beta \\ & + \left(\frac{\varphi''}{\varphi'} \overset{\circ}{y}_j \delta^i_k + \frac{\varphi''}{\varphi' + 2F^2\varphi''} g_{jk} y^i\right) B^k_a H^a_\beta,\end{aligned} \tag{2.16}$$

where $\overset{\circ}{\check{L}}{}^i_{j\beta} = \overset{\circ}{L}{}^i_{jk} B^k_\beta + \overset{\circ}{C}{}^i_{jk} B^k_a \overset{\circ}{H}{}^a_\beta$.

Using (1.12) in (2.13), we find that

$$\begin{aligned}\check{C}^i_{j\beta} = {} & \overset{\circ}{C}{}^i_{jk} B^k_\beta + \left(\frac{\varphi''}{\varphi'}\left(\delta^i_j \overset{\circ}{y}_k + \delta^i_k \overset{\circ}{y}_j\right) + \frac{\varphi''}{\varphi' + 2F^2\varphi''} g_{jk} y^i \right. \\ & \left. + \frac{2\left(\varphi'''\varphi' - 2\varphi''^2\right)}{\varphi'\left(\varphi' + 2F^2\varphi''\right)} y^i \overset{\circ}{y}_j \overset{\circ}{y}_k \right) B^k_\beta,\end{aligned} \tag{2.17}$$

that is,

$$\begin{aligned}\check{C}^i_{j\beta} = {} & \overset{\circ}{\check{C}}{}^i_{j\beta} + \left(\frac{\varphi''}{\varphi'}\left(\delta^i_j \overset{\circ}{y}_k + \delta^i_k \overset{\circ}{y}_j\right) + \frac{\varphi''}{\varphi' + 2F^2\varphi''} g_{jk} y^i \right. \\ & \left. + \frac{2\left(\varphi'''\varphi' - 2\varphi''^2\right)}{\varphi'\left(\varphi' + 2F^2\varphi''\right)} y^i \overset{\circ}{y}_j \overset{\circ}{y}_k \right) B^k_\beta,\end{aligned} \tag{2.18}$$

where $\overset{\circ}{\check{C}}{}^{i}_{j\beta} = \overset{\circ}{C}{}^{i}_{jk} B^k_\beta$. Thus, we have the following.

Theorem 2.5. *The coefficients of coupling for the subspace $\check{L}^m$ of an APL space L^n are given by* (2.16) *and* (2.18).

Definition 2.6 (cf. [8]). An operator D^T given by

$$D^T X^\alpha = X^\alpha_{|\beta} du^\beta + X^\alpha|_\beta \delta v^\beta, \tag{2.19}$$

where $X^\alpha_{|\beta} = \delta_\beta X^\alpha + X^\gamma L^\alpha_{\gamma\beta}$, $X^\alpha|_\beta = \dot{\partial}_\beta X^\alpha + X^\gamma C^\alpha_{\gamma\beta}$, is called the induced tangent connection by D. This defines an N-linear connection for $\check{L}^m$.

The coefficients $(L^\alpha_{\gamma\beta}, C^\alpha_{\gamma\beta})$ of D^T are given by

$$L^\alpha_{\beta\gamma} = B^\alpha_i \left(B^i_{\beta\gamma} + B^j_\beta \check{L}^i_{j\gamma} \right), \tag{2.20}$$

$$C^\alpha_{\beta\gamma} = B^\alpha_i B^j_\beta \check{C}^i_{j\gamma}. \tag{2.21}$$

Using (2.16) in (2.20), we get

$$\begin{aligned} L^\alpha_{\beta\gamma} = B^\alpha_i B^i_{\beta\gamma} + B^j_\beta B^\alpha_i \Bigg[& \overset{\circ}{\check{L}}{}^{i}_{j\gamma} + \left(V^r_k C^i_{jr} + V^r_j C^i_{kr} + V^r_p a^{ip} C_{rkj} - \overset{\circ}{C}{}^{i}_{jr} B^r_b B^b_p V^p_k \right) B^k_\gamma \\ & + \left(\frac{\varphi''}{\varphi'} \overset{\circ}{y}_j \delta^i_k + \frac{\varphi''}{\varphi' + 2F^2\varphi''} g_{jk} y^i \right) B^k_a H^a_\gamma \Bigg], \end{aligned} \tag{2.22}$$

that is,

$$\begin{aligned} L^\alpha_{\beta\gamma} = B^\alpha_i \left(B^i_{\beta\gamma} + \overset{\circ}{\check{L}}{}^{i}_{j\gamma} B^j_\beta \right) + B^\alpha_i B^j_\beta \Bigg[& \left(V^r_k C^i_{jr} + V^r_j C^i_{kr} + V^r_p a^{ip} C_{rkj} - \overset{\circ}{C}{}^{i}_{jr} B^r_b B^b_p V^p_k \right) B^k_\gamma \\ & + \left(\frac{\varphi''}{\varphi'} \overset{\circ}{y}_j \delta^i_k + \frac{\varphi''}{\varphi' + 2F^2\varphi''} g_{jk} y^i \right) B^k_a H^a_\gamma \Bigg]. \end{aligned} \tag{2.23}$$

If we take $\overset{\circ}{L}{}^{\alpha}_{\beta\gamma} = B^\alpha_i (B^i_{\beta\gamma} + \overset{\circ}{\check{L}}{}^{i}_{j\gamma} B^j_\beta)$, the last expression gives

$$\begin{aligned} L^\alpha_{\beta\gamma} = \overset{\circ}{L}{}^{\alpha}_{\beta\gamma} + B^\alpha_i B^j_\beta \Bigg[& \left(V^r_k C^i_{jr} + V^r_j C^i_{kr} + V^r_p a^{ip} C_{rkj} - \overset{\circ}{C}{}^{i}_{jr} B^r_b B^b_p V^p_k \right) B^k_\gamma \\ & + \left(\frac{\varphi''}{\varphi'} \overset{\circ}{y}_j \delta^i_k + \frac{\varphi''}{\varphi' + 2F^2\varphi''} g_{jk} y^i \right) B^k_a H^a_\gamma \Bigg]. \end{aligned} \tag{2.24}$$

Next, using (2.18) in (2.21), we obtain

$$C^{\alpha}_{\beta\gamma} = B^{\alpha}_{i} B^{j}_{\beta} \overset{\circ}{\check{C}}{}^{i}_{j\gamma} + \left(\frac{\varphi''}{\varphi'} \left(\delta^{i}_{j} \overset{\circ}{y}_{k} + \delta^{i}_{k} \overset{\circ}{y}_{j} \right) + \frac{\varphi''}{\varphi' + 2F^{2}\varphi''} g_{jk} y^{i} + \frac{2\left(\varphi'''\varphi' - 2\varphi''^{2}\right)}{\varphi'\left(\varphi' + 2F^{2}\varphi''\right)} y^{i} \overset{\circ}{y}_{j} \overset{\circ}{y}_{k} \right) B^{k}_{\gamma} B^{\alpha}_{i} B^{j}_{\beta}. \tag{2.25}$$

If we take $\overset{\circ}{C}{}^{\alpha}_{\beta\gamma} = B^{\alpha}_{i} B^{j}_{\beta} \overset{\circ}{\check{C}}{}^{i}_{j\gamma}$, (2.25) becomes

$$C^{\alpha}_{\beta\gamma} = \overset{\circ}{C}{}^{\alpha}_{\beta\gamma} + \left(\frac{\varphi''}{\varphi'} \left(\delta^{i}_{j} \overset{\circ}{y}_{k} + \delta^{i}_{k} \overset{\circ}{y}_{j} \right) + \frac{\varphi''}{\varphi' + 2F^{2}\varphi''} g_{jk} y^{i} + \frac{2\left(\varphi'''\varphi' - 2\varphi''^{2}\right)}{\varphi'\left(\varphi' + 2F^{2}\varphi''\right)} y^{i} \overset{\circ}{y}_{j} \overset{\circ}{y}_{k} \right) B^{k}_{\gamma} B^{\alpha}_{i} B^{j}_{\beta}. \tag{2.26}$$

Thus, we have the following.

Theorem 2.7. *The coefficients of the induced tangent connection D^{T} for the subspace $\check{L}^{m}$ of an APL space are given by* (2.24) *and* (2.26).

Remarks. The torsion $T^{\alpha}_{\beta\gamma} = L^{\alpha}_{\beta\gamma} - L^{\alpha}_{\gamma\beta}$ does not vanish, in general, while $S^{\alpha}_{\beta\gamma} = C^{\alpha}_{\beta\gamma} - C^{\alpha}_{\gamma\beta} = 0$. These facts may be observed from (2.24) and (2.26).

Definition 2.8 (cf. [8]). An operator $D^{\perp}$ given by

$$D^{\perp} X^{a} = X^{a}_{|\alpha} du^{\alpha} + X^{a}|_{\alpha} \delta v^{\alpha}, \tag{2.27}$$

where $X^{a}_{|\alpha} = \delta_{\alpha} X^{a} + X^{b} L^{a}_{b\alpha}$, $X^{a}|_{\alpha} = \dot{\partial}_{\alpha} X^{a} + X^{b} C^{a}_{b\alpha}$, is called the induced normal connection by D.

The coefficients $(L^{a}_{b\gamma}, C^{a}_{b\gamma})$ of $D^{\perp}$ are given by

$$L^{a}_{b\gamma} = B^{a}_{i} \left(\delta_{\gamma} B^{i}_{b} + B^{j}_{b} \check{L}^{i}_{j\gamma} \right), \tag{2.28}$$

$$C^{a}_{b\gamma} = B^{a}_{i} \left(\dot{\partial}_{\gamma} B^{i}_{b} + B^{j}_{b} \check{C}^{i}_{j\gamma} \right). \tag{2.29}$$

Using (2.6) and (2.16) in (2.28), we find

$$\begin{aligned} L^{a}_{b\gamma} = {} & B^{a}_{i} \overset{\circ}{\delta}_{\gamma} B^{i}_{b} + B^{a}_{i} B^{\alpha}_{p} V^{p}_{j} B^{j}_{\gamma} \dot{\partial}_{\alpha} B^{i}_{b} \\ & + B^{j}_{b} B^{a}_{i} \left[\overset{\circ}{\check{L}}{}^{i}_{j\gamma} + \left(V^{r}_{k} C^{i}_{jr} + V^{r}_{j} C^{i}_{kr} + V^{r}_{p} a^{ip} C_{rkj} - \overset{\circ}{C}{}^{i}_{jr} B^{r}_{c} B^{c}_{p} V^{p}_{k} \right) B^{k}_{\gamma} \right. \\ & \left. + \left(\frac{\varphi''}{\varphi'} \overset{\circ}{y}_{j} \delta^{i}_{k} + \frac{\varphi''}{\varphi' + 2F^{2}\varphi''} g_{jk} y^{i} \right) B^{k}_{c} H^{c}_{\gamma} \right]. \end{aligned} \tag{2.30}$$

Taking $\overset{\circ}{L}{}^{a}_{b\gamma} = B^a_i(\overset{\circ}{\delta}_\gamma B^i_b + B^j_b \overset{\circ}{\check{L}}{}^{i}_{j\gamma})$ and using $\overset{\circ}{y}_j B^j_b = 0$, (2.30) reduces to

$$L^a_{b\gamma} = \overset{\circ}{L}{}^{a}_{b\gamma} + B^a_i B^\alpha_p V^p_j B^j_\gamma \dot{\partial}_\alpha B^i_b + \left(V^r_k C^i_{jr} + V^r_j C^i_{kr} + V^r_p a^{ip} C_{rkj} - \overset{\circ}{C}{}^{i}_{jr} B^r_b B^b_p V^p_k \right) B^a_i B^j_b B^k_\gamma + \frac{\varphi''}{\varphi' + 2F^2\varphi''} g_{jk} y^i B^k_c H^c_\gamma B^a_i B^j_b. \tag{2.31}$$

Next, using (2.18) in (2.29), we have

$$C^a_{b\gamma} = B^a_i \left(\dot{\partial}_\gamma B^i_b + B^j_b \overset{\circ}{\check{C}}{}^{i}_{j\gamma} \right) + \left[\frac{\varphi''}{\varphi'} \left(\delta^i_j \overset{\circ}{y}_k + \delta^i_k \overset{\circ}{y}_j \right) + \frac{\varphi''}{\varphi' + 2F^2\varphi''} g_{jk} y^i + \frac{2(\varphi'''\varphi' - 2\varphi''^2)}{\varphi'(\varphi' + 2F^2\varphi'')} y^i \overset{\circ}{y}_j \overset{\circ}{y}_k \right] B^k_\gamma B^a_i B^j_b. \tag{2.32}$$

Taking $\overset{\circ}{C}{}^{a}_{b\gamma} = B^a_i(\dot{\partial}_\gamma B^i_b + B^j_b \overset{\circ}{\check{C}}{}^{i}_{j\gamma})$ and using (1.9) and $\overset{\circ}{y}_j B^j_b = 0$, the last equation yields

$$C^a_{b\gamma} = \overset{\circ}{C}{}^{a}_{b\gamma} + \frac{\varphi''}{\varphi'} \delta^a_b \overset{\circ}{y}_k B^k_\gamma + \frac{\varphi''}{\varphi' + 2F^2\varphi''} g_{jk} y^i B^k_\gamma B^a_i B^j_b. \tag{2.33}$$

Thus, we have the following.

Theorem 2.9. *The coefficients of induced normal connection $D^\perp$ for the subspace $\check{L}^m$ of an APL space L^n are given by* (2.31) *and* (2.33).

Definition 2.10 (cf. [8]). The (mixed) derivative of a mixed d-tensor field $T^{i\cdots\alpha\cdots a}_{j\cdots\beta\cdots b}$ is given by

$$\begin{aligned}
\nabla T^{i\cdots\alpha\cdots a}_{j\cdots\beta\cdots b} = {} & \Big(\delta_\eta T^{i\cdots\alpha\cdots a}_{j\cdots\beta\cdots b} + T^{k\cdots\alpha\cdots a}_{j\cdots\beta\cdots b} \check{L}^i_{k\eta} + \cdots + T^{i\cdots\gamma\cdots a}_{j\cdots\beta\cdots b} L^\alpha_{\gamma\eta} + \cdots + T^{i\cdots\alpha\cdots c}_{j\cdots\beta\cdots b} L^a_{c\eta} \\
& - T^{i\cdots\alpha\cdots a}_{k\cdots\beta\cdots b} \check{L}^k_{j\eta} - \cdots - T^{i\cdots\alpha\cdots a}_{j\cdots\gamma\cdots b} L^\gamma_{\beta\eta} - \cdots - T^{i\cdots\alpha\cdots a}_{j\cdots\beta\cdots c} \check{L}^c_{b\eta} \Big) du^\eta \\
& + \Big(\dot{\partial}_\eta T^{i\cdots\alpha\cdots a}_{j\cdots\beta\cdots b} + T^{k\cdots\alpha\cdots a}_{j\cdots\beta\cdots b} \check{C}^i_{k\eta} + \cdots + T^{i\cdots\gamma\cdots a}_{j\cdots\beta\cdots b} C^\alpha_{\gamma\eta} + \cdots + T^{i\cdots\alpha\cdots c}_{j\cdots\beta\cdots b} C^a_{c\eta} \\
& - T^{i\cdots\alpha\cdots a}_{k\cdots\beta\cdots b} \check{C}^k_{j\eta} - \cdots - T^{i\cdots\alpha\cdots a}_{j\cdots\gamma\cdots b} C^\gamma_{\beta\eta} - \cdots - T^{i\cdots\alpha\cdots a}_{j\cdots\beta\cdots c} \check{C}^c_{b\eta} \Big) \delta v^\eta.
\end{aligned} \tag{2.34}$$

The connection 1-forms,

$$\check{\omega}^i_j =: \check{L}^i_{j\alpha} du^\alpha + \check{C}^i_{j\alpha} \delta v^\alpha, \tag{2.35}$$

$$\omega^\alpha_\beta =: L^\alpha_{\beta\gamma} du^\gamma + C^\alpha_{\beta\gamma} \delta v^\gamma, \tag{2.36}$$

$$\omega^a_b =: L^a_{b\gamma} du^\gamma + C^a_{b\gamma} \delta v^\gamma, \tag{2.37}$$

are called the connection 1-forms of ∇. We have the following structure equations of ∇.

Theorem 2.11 (cf. [8]). *The structure equations of ∇ are as follows:*

$$\begin{aligned}
&d(du^\alpha) - du^\beta \wedge \omega^\alpha_\beta = -\Omega^\alpha,\\
&d(\delta u^\alpha) - \delta u^\beta \wedge \omega^\alpha_\beta = -\dot{\Omega}^\alpha,\\
&d\check{\omega}^i_j - \check{\omega}^h_j \wedge \check{\omega}^i_h = -\check{\Omega}^i_j,\\
&d\omega^\alpha_\beta - \omega^\gamma_\beta \wedge \omega^\alpha_\gamma = -\Omega^\alpha_\beta,\\
&d\omega^a_b - \omega^c_b \wedge \omega^a_c = -\Omega^a_b,
\end{aligned} \tag{2.38}$$

where the 2-forms of torsions $\Omega^\alpha, \dot{\Omega}^\alpha$ are given by

$$\begin{aligned}
\Omega^\alpha &= \frac{1}{2}T^\alpha_{\beta\gamma}du^\beta \wedge du^\gamma + C^\alpha_{\beta\gamma}du^\beta \wedge \delta v^\gamma,\\
\dot{\Omega}^\alpha &= \frac{1}{2}R^\alpha_{\beta\gamma}du^\beta \wedge du^\gamma + P^\alpha_{\beta\gamma}du^\beta \wedge \delta v^\gamma,
\end{aligned} \tag{2.39}$$

with $P^\alpha_{\beta\gamma} = \dot{\partial}_\gamma \check{N}^\alpha_\beta - L^\alpha_{\beta\gamma}$, and the 2-forms of curvature $\check{\Omega}^i_j, \Omega^\alpha_\beta$ and Ω^a_b, are given by

$$\begin{aligned}
\check{\Omega}^i_j &= \frac{1}{2}\check{R}^i_{j\alpha\beta}du^\alpha \wedge du^\beta + \check{P}^i_{j\alpha\beta}du^\alpha \wedge \delta v^\beta + \frac{1}{2}\check{S}^i_{j\alpha\beta}\delta v^\alpha \wedge \delta v^\beta,\\
\Omega^\alpha_\beta &= \frac{1}{2}R^\alpha_{\beta\gamma\delta}du^\gamma \wedge du^\delta + P^\alpha_{\beta\gamma\delta}du^\gamma \wedge \delta v^\delta + \frac{1}{2}S^\alpha_{\beta\gamma\delta}\delta v^\gamma \wedge \delta v^\delta,\\
\Omega^a_b &= \frac{1}{2}R^a_{b\alpha\beta}du^\alpha \wedge du^\beta + P^a_{b\alpha\beta}du^\alpha \wedge \delta v^\beta + \frac{1}{2}S^a_{b\alpha\beta}\delta v^\alpha \wedge \delta v^\beta.
\end{aligned} \tag{2.40}$$

We will use the following notations in Section 4:

$$\text{(a) } \check{\Omega}_{ij} = \check{\Omega}^h_i a_{hj}, \qquad \text{(b) } \Omega_{\alpha\beta} = \Omega^\gamma_\alpha a_{\gamma\beta}, \qquad \text{(c) } \Omega_{ab} = \Omega^c_b \delta_{ac}. \tag{2.41}$$

3. The Gauss-Weingarten Formulae

The Gauss-Weingarten formulae for the subspace $\check{L}^m = (\check{M}, \check{L}(u,v))$ of a Lagrange space L^n are given by (cf. [8])

$$\nabla B^i_\alpha = B^i_a \Pi^a_\alpha, \qquad \nabla B^i_a = -B^i_\beta \Pi^\beta_a, \tag{3.1}$$

where

$$\Pi^a_\alpha = H^a_{\alpha\beta} du^\beta + K^a_{\alpha\beta} \delta v^\beta,$$
$$\Pi^\beta_a = g^{\beta\gamma} \delta_{ab} \Pi^b_\gamma, \tag{3.2}$$

$$\text{(a)}\ H^a_{\alpha\beta} = B^a_i \left(\delta_\beta B^i_\alpha + B^j_\alpha \check{L}^i_{j\beta} \right), \qquad \text{(b)}\ K^a_{\alpha\beta} = B^a_i B^j_\alpha \check{C}^i_{j\beta}. \tag{3.3}$$

Using (2.6) and (2.16) in (3.3)(a), we have

$$\begin{aligned} H^a_{\alpha\beta} = {} & B^a_i \left(\overset{\circ}{\delta}_\beta B^i_\alpha + B^j_\alpha \overset{\circ}{\check{L}}{}^i_{j\beta} \right) + B^a_i B^\gamma_p V^p_j B^j_\beta B^i_{\alpha\gamma} \\ & + \left(V^r_k C^i_{jr} + V^r_j C^i_{kr} + V^r_p a^{ip} C_{rkj} - \overset{\circ}{C}{}^i_{jr} B^r_b B^b_p V^p_k \right) B^a_i B^j_\alpha B^k_\beta \\ & + \left(\frac{\varphi''}{\varphi'} \overset{\circ}{y}_j \delta^i_k + \frac{\varphi''}{\varphi' + 2F^2\varphi''} g_{jk} y^i \right) B^k_b H^b_\beta B^a_i B^j_\alpha. \end{aligned} \tag{3.4}$$

If we take $\overset{\circ}{H}{}^a_{\alpha\beta} = B^a_i (\overset{\circ}{\delta}_\beta B^i_\alpha + B^j_\alpha \overset{\circ}{\check{L}}{}^i_{j\beta})$, the last expression provides

$$\begin{aligned} H^a_{\alpha\beta} = {} & \overset{\circ}{H}{}^a_{\alpha\beta} + B^a_i B^\gamma_p V^p_j B^j_\beta B^i_{\alpha\gamma} + \left(V^r_k C^i_{jr} + V^r_j C^i_{kr} + V^r_p a^{ip} C_{rkj} - \overset{\circ}{C}{}^i_{jr} B^r_b B^b_p V^p_k \right) B^a_i B^j_\alpha B^k_\beta \\ & + \left(\frac{\varphi''}{\varphi'} \overset{\circ}{y}_j \delta^i_k + \frac{\varphi''}{\varphi' + 2F^2\varphi''} g_{jk} y^i \right) B^k_b H^b_\beta B^a_i B^j_\alpha. \end{aligned} \tag{3.5}$$

Next, using (2.18) in (3.3)(b) and keeping (1.9) in view, we find

$$K^a_{\alpha\beta} = \overset{\circ}{K}{}^a_{\alpha\beta} + \left(\frac{\varphi''}{\varphi' + 2F^2\varphi''} g_{jk} y^i + \frac{2(\varphi'''\varphi' - \varphi''^2)}{\varphi'(\varphi' + 2F^2\varphi'')} y^i \overset{\circ}{y}_j \overset{\circ}{y}_k \right) B^a_i B^j_\alpha B^k_\beta, \tag{3.6}$$

where $\overset{\circ}{K}{}^a_{\alpha\beta} = B^a_i B^j_\alpha \overset{\circ}{\check{C}}{}^i_{j\beta}$. Thus, we have the following.

Theorem 3.1. *The following Gauss-Weingarten formulae for the subspace $\check{L}^m$ of an APL space hold:*

$$\nabla B^i_\alpha = B^i_a \Pi^a_\alpha, \qquad \nabla B^i_a = -B^i_\beta \Pi^\beta_a, \tag{3.7}$$

where

$$\Pi^a_\alpha = H^a_{\alpha\beta} du^\beta + K^a_{\alpha\beta}\delta v^\beta, \qquad \Pi^\beta_a = g^{\beta\gamma}\delta_{ab}\Pi^b_\gamma,$$

$$\begin{aligned} H^a_{\alpha\beta} &= \overset{\circ}{H}{}^a_{\alpha\beta} + B^a_i B^\gamma_p V^p_j B^j_\beta B^i_{\alpha\gamma} + \left(V^r_k C^i_{jr} + V^r_j C^i_{kr} + V^r_p a^{ip} C_{rkj} - \overset{\circ}{C}{}^i_{jr} B^r_b B^b_p V^p_k\right) B^a_i B^j_\alpha B^k_\beta \\ &\quad + \left(\frac{\varphi''}{\varphi'}\overset{\circ}{y}_j \delta^i_k + \frac{\varphi''}{\varphi' + 2F^2\varphi''} g_{jk} y^i\right) B^k_b H^b_\beta B^a_i B^j_\alpha, \\ K^a_{\alpha\beta} &= \overset{\circ}{K}{}^a_{\alpha\beta} + \left(\frac{\varphi''}{\varphi' + 2F^2\varphi''} g_{jk} y^i + \frac{2(\varphi'''\varphi' - \varphi''^2)}{\varphi'(\varphi' + 2F^2\varphi'')} y^i \overset{\circ}{y}_j \overset{\circ}{y}_k\right) B^a_i B^j_\alpha B^k_\beta. \end{aligned} \tag{3.8}$$

Remark 3.2. $H^a_{\alpha\beta}$ and $K^a_{\alpha\beta}$ given, respectively, by (3.5) and (3.6) are called the second fundamental d-tensor fields of immersion i.

The following consequences of Theorem 3.1 are straightforward.

Corollary 3.3. *In a subspace $\check{L}^m$ of an APL space, we have the following:*

$$\begin{aligned} &(a)\ \nabla a_{\alpha\beta} = 0, \\ &(b)\ \nabla B^i_\alpha = 0, \end{aligned} \tag{3.9}$$

if and only if

$$\begin{aligned} \overset{\circ}{H}{}^a_{\alpha\beta} &= -\left[B^a_i B^\gamma_p V^p_j B^j_\beta B^i_{\alpha\gamma} + \left(V^r_k C^i_{jr} + V^r_j C^i_{kr} + V^r_p a^{ip} C_{rkj} - \overset{\circ}{C}{}^i_{jr} B^r_b B^b_p V^p_k\right) B^a_i B^j_\alpha B^k_\beta \right. \\ &\quad \left. + \left(\frac{\varphi''}{\varphi'}\overset{\circ}{y}_j \delta^i_k + \frac{\varphi''}{\varphi' + 2F^2\varphi''} g_{jk} y^i\right) B^k_b H^b_\beta B^a_i B^j_\alpha\right], \\ \overset{\circ}{K}{}^a_{\alpha\beta} &= -\left(\frac{\varphi''}{\varphi' + 2F^2\varphi''} g_{jk} y^i + \frac{2\left(\varphi'''\varphi' - \varphi''^2\right)}{\varphi'(\varphi' + 2F^2\varphi'')} y^i \overset{\circ}{y}_j \overset{\circ}{y}_k\right) B^a_i B^j_\alpha B^k_\beta. \end{aligned} \tag{3.10}$$

4. The Gauss-Codazzi Equations

The Gauss-Codazzi Equations for the subspace $\check{L}^m = (\check{M}, \check{L}(u,v))$ of a Lagrange space L^n are given by (cf. [8])

$$B^i_\alpha B^j_\beta \check{\Omega}_{ij} - \Omega_{\alpha\beta} = \Pi_{\beta a} \wedge \Pi^a_\alpha, \tag{4.1}$$

$$B^i_a B^j_b \check{\Omega}_{ij} - \Omega_{ab} = \Pi_{\gamma b} \wedge \Pi^\gamma_a, \tag{4.2}$$

$$-B^i_\alpha B^j_a \check{\Omega}_{ij} = \delta_{ab}\left(d\Pi^b_\alpha + \Pi^b_\beta \wedge \omega^\beta_\alpha - \Pi^c_\alpha \wedge \omega^b_c\right), \tag{4.3}$$

where

$$\text{(a) } \Pi_{\alpha a} = g_{\alpha\beta}\Pi^{\beta}_{a}, \qquad \text{(b) } \Pi_{\gamma b} = \delta_{bc}\Pi^{c}_{\gamma}. \tag{4.4}$$

Using (1.3) in (2.41)(a), we find that

$$\check{\Omega}_{ij} = \varphi'\check{\Omega}^{h}_{i} g_{hj} + 2\varphi''\check{\Omega}^{h}_{i}\overset{\circ}{y}_{h}\overset{\circ}{y}_{j}. \tag{4.5}$$

Applying $a_{\gamma\beta} = B^{i}_{\gamma}B^{j}_{\beta}a_{ij}$ in (2.41)(b), we have $\Omega_{\alpha\beta} = B^{i}_{\gamma}B^{j}_{\beta}\Omega^{\gamma}_{\alpha}a_{ij}$, which in view of (1.3) becomes

$$\Omega_{\alpha\beta} = \varphi' g_{ij}B^{i}_{\gamma}B^{j}_{\beta}\Omega^{\gamma}_{\alpha} + 2\varphi''\overset{\circ}{y}_{i}\overset{\circ}{y}_{j}B^{i}_{\gamma}B^{j}_{\beta}\Omega^{\gamma}_{\alpha}, \tag{4.6}$$

that is,

$$\Omega_{\alpha\beta} = \varphi' g_{\gamma\beta}\Omega^{\gamma}_{\alpha} + 2\varphi''\overset{\circ}{y}_{i}\overset{\circ}{y}_{j}B^{i}_{\gamma}B^{j}_{\beta}\Omega^{\gamma}_{\alpha}. \tag{4.7}$$

For the subspace $\check{L}^{m}$ of an APL space, (4.4)(a) is of the form $\Pi_{\alpha a} = a_{\alpha\beta}\Pi^{\beta}_{a}$, which in view of $a_{\alpha\beta} = B^{i}_{\alpha}B^{j}_{\beta}a_{ij}$ and (1.3) becomes $\Pi_{\alpha a} = \varphi' B^{i}_{\alpha}B^{j}_{\beta}a_{ij}\Pi^{\beta}_{a} + 2\varphi''\overset{\circ}{y}_{i}\overset{\circ}{y}_{j}B^{i}_{\alpha}B^{j}_{\beta}\Pi^{\beta}_{a}$, that is,

$$\Pi_{\alpha a} = \varphi' g_{\alpha\beta}\Pi^{\beta}_{a} + 2\varphi''\overset{\circ}{y}_{i}\overset{\circ}{y}_{j}B^{i}_{\alpha}B^{j}_{\beta}\Pi^{\beta}_{a}. \tag{4.8}$$

Thus, we have the following.

Theorem 4.1. *The Gauss-Codazzi equations for a Lagrange subspace $\check{L}^{m}$ of an APL space are given by* (4.1)–(4.3) *with* $\Pi_{\alpha a}$, $\Pi_{\gamma b}$, $\check{\Omega}_{ij}$, $\Omega_{\alpha\beta}$, *and* ω^{b}_{c}, *respectively, given by* (4.8), (4.4)*(b)*, (4.5), (4.7), *and* (2.37).

Acknowledgments

Authors are thankful to the reviewers for their valuable comments and suggestions. S. K. Shukla gratefully acknowledges the financial support provided by the Council of Scientific and Industrial Research (CSIR), India.

References

[1] R. Miron, "Lagrange geometry," *Mathematical and Computer Modelling*, vol. 20, no. 4-5, pp. 25–40, 1994.
[2] R. Miron and A. Bejancu, "A nonstandard theory of Finsler subspaces," in *Topics in Differential Geometry*, vol. 46 of *Colloquia Mathematica Societatis János Bolyai*, pp. 815–851, 1984.
[3] R. Miron and M. Anastasiei, *Vector Bundles and Lagrange Spaces with Applications to Relativity*, vol. 1, Geometry Balkan Press, Bucharest, Romania, 1997.
[4] T. Sakaguchi, *Subspaces in Lagrange spaces [Ph.D. thesis]*, University "Al. I. Cuza", Iasi, Romania, 1986.
[5] P. L. Antonelli and D. Hrimiuc, "A new class of spray-generating Lagrangians," in *Lagrange and Finsler Geometry*, vol. 76 of *Applications to Physics and Biology*, pp. 81–92, Kluwer Academic Publishers, Dordrecht, The Netherlands, 1996.

[6] P. L. Antonelli and D. Hrimiuc, "On the theory of φ-Lagrange manifolds with applications in biology and physics," *Nonlinear World*, vol. 3, no. 3, pp. 299–333, 1996.
[7] P. N. Pandey and S. K. Shukla, "On almost φ-Lagrange spaces," *ISRN Geometry*, vol. 2011, Article ID 505161, 16 pages, 2011.
[8] P. L. Antonelli, Ed., *Handbook of Finsler Geometry*, Kluwer Academic Publishers, Dordrecht, The Netherlands, 2001.
[9] R. Miron and M. Anastasiei, *The Geometry of Lagrange Spaces: Theory and Applications*, Kluwer Academic Publishers, 1994.