Research Article

On the Structures of Abelian π -Regular Rings

Jianwei Lu and Ligu He

Department of Mathematics, Shenyang University of Technology, Shenyang 110870, China

Correspondence should be addressed to Jianwei Lu; jianwei191@sohu.com

Received 14 November 2013; Revised 24 February 2014; Accepted 5 March 2014; Published 1 April 2014

Academic Editor: A. Cigdem Ozcan

Copyright © 2014 J. Lu and L. He. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Assume that R is an Abelian ring. In this paper, we characterize the structure of R whenever R is π -regular. It is also proved that an Abelian π -regular ring is isomorphic to the subdirect sum of some metadivision rings.

1. Introduction

Throughout this paper, we only consider the associative rings with identity 1. By $E(R)$, we denote the full set of idempotent elements of the ring R . Write $C(R)$ to denote the center of the ring R , and write $\text{Nil}(R)$, the full set of nilpotent elements of R . The ring R is called *Abelian* if $E(R) \subseteq C(R)$. The ring R is said to be *regular* if, for any $a \in R$, there exists an $x \in R$ such that $a = axa$. And R is called *π -regular* if, for any $a \in R$, there exist $x \in R$ and a positive integer n such that $a^n = a^n x a^n$. We also say a ring R is *strongly π -regular* if, for any $a \in R$, there are $x \in R$ and positive integer n such that $a^n = x a^{n+1}$. It follows that strongly π -regular rings are π -regular and Abelian π -regular rings are strongly π -regular. The ring R is said to be a *metadivision* if any element in R is either a unit or a nilpotent element.

In ring theory, there has been much interest in finding connections between the π -regularity and the condition that every prime ideal is maximal, which has been studied, for example, in [1–6]. A pretty result due to Storrer in [6] is if R is a commutative ring with identity then R is π -regular if and only if every prime ideal of R is maximal. A commutative ring is obviously an Abelian ring; thus, the following result is a generalization of Storrer's result.

Theorem A. *Let R be an Abelian ring. Then R is π -regular if and only if the following statements hold:*

- (1) $\text{Nil}(R)$ is an ideal of R ;
- (2) every one-sided ideal containing $\text{Nil}(R)$ is an ideal in R ;
- (3) every completely prime ideal is a maximal ideal in R .

A classical result due to McCoy in [7] states that a commutative π -regular ring is isomorphic to the subdirect sum of some metadivision rings. We generalize this result as follows.

Theorem B. *Assume that R is an Abelian π -regular ring and $\mathcal{M}$ denotes the collection of all completely prime ideals of R . Then R is isomorphic to the subdirect sum of the $R/E(P_\alpha)P_\alpha$'s for all $P_\alpha \in \mathcal{M}$.*

2. Abelian π -Regular Rings

Recall that an ideal Q of R is *primary* if $ab \in Q$ for any $a, b \in R$; we always obtain $a^n \in Q$ or $b^n \in Q$ for some positive integer n .

Definition 1. The ring R is said to be a metadivision ring if any element in R is either a unit or a nilpotent element.

In fact, there is an intimate connection between the primary ideals and the metadivision rings as follows.

Lemma 2. *Let R be an Abelian π -regular ring. Then Q is a primary ideal of R if and only if R/Q is a metadivision ring.*

Proof. The “only if” part: for any element $a \in R$, there exists the equality $a^n = a^n r a^n$ for a positive integer n and $r \in R$. Set $e = r a^n$; then e is an idempotent. If $\bar{a} \in R/Q$ is not nilpotent, then $\bar{a}^n \neq \bar{0}$ and so $\bar{e} \neq \bar{0}$; thus, $e \notin Q$. Since $e(e-1) = 0$ lies in the primary ideal Q , it follows that $(e-1)^m \in Q$ for some $m > 0$. Also $1-e$ is an idempotent; we have $1-e = (1-e)^m \in Q$, and so $\bar{e} = \bar{1}$; that is, $\bar{r} \bar{a}^n = \bar{1}$.

In the preceding proof, if we set $e = a^n r$, in a similar manner, then we can obtain another element $\bar{s}$ such that $\bar{a}\bar{s} = \bar{1}$. Hence we obtain that $\bar{a}$ is unit and thus R/Q is a metadivision ring.

The “if” part: assume that $a, b \in R$ with $ab \in Q$. If neither a^n nor b^n lies in Q for any positive integer n , then both of $\bar{a}$ and $\bar{b}$ are unit in R/Q (which is a metadivision ring). Thus there exist $r_1, r_2 \in R$ such that $\bar{r}_1 \bar{a} = \bar{1} = \bar{b} \bar{r}_2$, and so $\bar{r}_1 \bar{a} \bar{b} \bar{r}_2 = \bar{1}$. Also $ab \in Q$; it follows that $\bar{0} = \bar{r}_1 \bar{0} \bar{r}_2 = \bar{r}_1 \bar{a} \bar{b} \bar{r}_2 = \bar{1}$, a contradiction. Therefore we conclude that Q is a primary ideal of R . $\square$

Proposition 3. Suppose that R is an Abelian π -regular ring and $a^n = a^n x a^n$ for $a, x \in R$. Then, for any $m > n$, $a^m = a^m y a^m$ for some $y \in R$.

Proof. Let $e = a^n x$; then e is an idempotent and so lies in $C(R)$. Thus it is immediate that $a^n = e a^n = a^n e$. Because

$$\begin{aligned} a^{n+1} &= a^n a = a^n e a = a^{2n} x^2 a^{n+1} \\ &= a^{n+1} (a^{n-1} x^2) a^{n+1} = a^{n+1} y a^{n+1}, \end{aligned} \quad (1)$$

where $y = a^{n-1} x^2$, the application of the inductive argument on m yields the desired result. $\square$

Lemma 4. Let R be an Abelian π -regular ring and P a prime ideal of R . Then $E(P)P$ is a minimal primary ideal of R .

Proof. It is no loss to assume that P is a proper ideal of R . Because $E(P)P = PE(P)$, $E(P)P$ is an ideal. For $a, b \in R$ and $ab \in E(P)P$, the application of the above proposition yields that there exists a positive integer n such that $a^n = a^n r_1 a^n$ and $b^n = b^n r_2 b^n$. Given $e_1 = r_1 a^n$ and $e_2 = r_2 b^n$, then e_i are idempotent for $i = 1, 2$, and thus $e_1 e_2 = e_1 r_2 b^n = r_2 b^{n-1} e_1 b = r_2 b^{n-1} r_1 a^{n-1} ab \in E(P)P \subseteq P$. Considering P is prime, either e_1 or e_2 lies in P and so in $E(P)P$. We further get that $a^n = a^n e_1 \in E(P)P$ or $b^n = b^n e_2 \in E(P)P$, and hence $E(P)P$ is primary in R . Suppose now that $Q \subsetneq E(P)P$ is a primary ideal of R . Then there exists $e \in E(P) - Q$; we therefore have that $e(1 - e) = 0 \in Q$ and $1 - e = (1 - e)^m \in Q$ for some $m > 0$. It follows that $1 - e \in E(P)P$ and thus $1 = 1 - e + e \in E(P)P$, a contradiction. Therefore $E(P)P$ is a minimal primary ideal of R . $\square$

Lemma 5. Let R be an Abelian ring. Then R is Abelian π -regular if and only if $\text{Nil}(R)$ is an ideal and $R/\text{Nil}(R)$ is strongly regular.

Proof. This is Theorem 3 of [1]. $\square$

Lemma 6. Assume that R is an Abelian π -regular ring, and I is a one-sided ideal of R . If $I \supseteq \text{Nil}(R)$, then I is an ideal of R and $Ra + I = aR + I$ for any $a \in R$. In particular, a maximal one-sided ideal of R is a maximal two-sided ideal of R .

Proof. First we assert that if R is Abelian and regular, then a left ideal A of R is an ideal. For $r \in A$, $r = ryr$ with some $y \in R$. Let $e = yr$; then e is idempotent, and thus $rx = rex = rxe \in A$ for any $x \in R$. We get that A is an ideal.

Now assume that I is a left ideal of R and $I \supseteq \text{Nil}(R)$. Then $I/\text{Nil}(R)$ is a left ideal of $R/\text{Nil}(R)$. By Lemma 5, $R/\text{Nil}(R)$ is strongly regular, so $I/\text{Nil}(R)$ is an ideal of $R/\text{Nil}(R)$. For any $r \in I$, $x \in R$, $\bar{r}\bar{x} \in \bar{I} = I/\text{Nil}(R)$, and so $rx \in I$; hence, I is a right ideal. By the same argument, we may also deduce that if $I \supseteq \text{Nil}(R)$ is a right ideal, then I is a left ideal. Therefore, the one-sided ideal I is an ideal in R when I contains $\text{Nil}(R)$.

For $a \in R$, both $Ra + I$ and $aR + I$ are ideals of R . Since $ax \in Ra + I$, for any $x \in R$, $aR + I \subseteq Ra + I$. Likewise, $Ra + I \subseteq aR + I$, and thus we get that $Ra + I = aR + I$, as desired.

If P is a maximal one-sided ideal of R , then $P \supseteq J(R) = \text{Nil}(R)$, and thus P is an ideal by the next to last paragraph; it is automatically a maximal ideal. $\square$

Lemma 7. Assume that R is an Abelian π -regular ring; then the following statements are equivalent.

- (1) P is a completely prime ideal.
- (2) P is a prime ideal and $P \supseteq \text{Nil}(R)$.
- (3) R/P is a division ring.
- (4) P is a maximal ideal of R .

Proof. (2) $\Rightarrow$ (3). First we claim that if A is a prime ideal of an Abelian regular R , then R/A is a division ring. Let $0 \neq \bar{a} \in R/A$ and $a = ara = ae$ for $r \in R$ and $e = ra$. Then, $\bar{e} \neq \bar{0}$ and $e \notin A$. Also since $e(e - 1) = 0 \in A$, we have $e - 1 \in A$ and so $\bar{e} = \bar{1}$; that is, $\bar{r}\bar{a} = \bar{1}$. Thus R/A is a division ring. Now assume that P is a prime ideal of the Abelian π -regular ring R and $P \supseteq \text{Nil}(R)$. Applying the above lemma, $\bar{P} = P/\text{Nil}(R)$ is a prime ideal of the Abelian regular ring $\bar{R} = R/\text{Nil}(R)$, and then $R/P \cong \bar{R}/\bar{P}$ is a division ring, as wanted. The proofs of the others are obvious. $\square$

A regular ring is obviously a π -regular ring. The following lemma is its converse under some conditions.

Lemma 8. Let R be an Abelian ring and $\text{Nil}(R) = 0$. If R is π -regular, then R is regular.

Proof. Let $a \in R$. Then $a^n = a^n x a^n = a^n e$ where $e = ra^n$ for some $r \in R$ and a positive integer n ; then $(a(1 - e))^n = a^n(1 - e)^n = a^n(1 - e) = a^n - a^n e = 0$. Since $\text{Nil}(R) = 0$, it follows that $a(1 - e) = 0$, and so $a = ae = ara^n = a(ra^{n-1})a$; thus, R is a regular ring. $\square$

The next result is Theorem B.

Theorem 9. Let R be an Abelian ring. Then R is π -regular if and only if the following statements hold.

- (1) $\text{Nil}(R)$ is an ideal of R .
- (2) Every one-sided ideal containing $\text{Nil}(R)$ is an ideal in R .
- (3) Every completely prime ideal is a maximal ideal in R .

Proof. The “if part” immediately follows from Lemmas 5–7.

Now we prove the “only if” part. Suppose to the contrary that R is not a π -regular ring. Lemmas 5 and 8 imply that

$\bar{R} = R/\text{Nil}(R)$ is not π -regular, and so not a strong π -regular ring; then there exists $\bar{a} \in \bar{R}$ such that $\bar{R}\bar{a} \not\supseteq \bar{R}\bar{a}^2 \not\supseteq \cdots \not\supseteq \bar{R}\bar{a}^n \not\supseteq \cdots$, which is obviously an infinite descending chain of left ideals of R . Let $\mathcal{S}$ be a collection of ideals $\bar{I}$ of $\bar{R}$ such that the left ideal chains $\bar{R}\bar{a}^i + \bar{I}, i = 1, 2, \dots$, are infinitely descending. And $\mathcal{S}$ is not empty in view of $\bar{0} \in \mathcal{S}$. For any ascending chain in R , $\bar{I}_1 \subset \bar{I}_2 \subset \cdots$, it is clear $\cup \bar{I}_i \in \mathcal{S}$; thus, $\mathcal{S}$ has a maximal element $\bar{H}$. We claim that $\bar{H}$ is not a completely prime ideal. Write $\sigma^{-1}(\bar{H})$ to denote the preimage of the natural homomorphism σ from R onto $\bar{R}$ and let $a', b' \in R$ with $a'b' \in \sigma^{-1}(\bar{H})$; then, $\overline{a'b'} \in \bar{H}$. If $\bar{H}$ is a prime ideal of $\bar{R}$, then $\bar{a}' \in \bar{H}$ or $\bar{b}' \in \bar{H}$, and so $a' \in \sigma^{-1}(\bar{H})$ or $b' \in \sigma^{-1}(\bar{H})$. Hence $\sigma^{-1}(\bar{H})$ is a completely prime ideal of R . Part (3) implies that $\sigma^{-1}(\bar{H})$ is a maximal ideal of R , and thus $\bar{H}$ is a maximal ideal of $\bar{R}$. Part (2) implies that $\bar{H} + \bar{R}\bar{a}^{k+1} = \bar{H}$ or $\bar{H} + \bar{R}\bar{a}^{k+1} = \bar{R}$ for a positive integer k , and thus $\bar{H} + \bar{R}\bar{a}^{k+1} = \bar{H} + \bar{R}\bar{a}^{k+2}$ or $\bar{H} + \bar{R}\bar{a}^{k+1} = \bar{H} + \bar{R}\bar{a}^k$. This is a contradiction, however, because $\bar{H} \in \mathcal{S}$. Hence $\bar{H}$ is not a completely prime ideal, and so there exist $\bar{a}_1, \bar{b}_1 \in \bar{R}$ such that $\bar{a}_1\bar{b}_1 \in \bar{H}$, whereas neither $\bar{a}_1$ nor $\bar{b}_1$ lies in $\bar{H}$. Set $\bar{H}_1 = \bar{R}\bar{a}_1 + \bar{H}$ and $\bar{H}_2 = \bar{R}\bar{b}_1 + \bar{H}$; then, both $\bar{H}_1$ and $\bar{H}_2$ are ideals of $\bar{R}$, which properly contain $\bar{H}$. Thus neither $\bar{H}_1$ nor $\bar{H}_2$ lies in $\mathcal{S}$; consequently, there exists some positive integer n such that $\bar{R}\bar{a}^n + \bar{H}_1 = \bar{R}\bar{a}^{n+1} + \bar{H}_1$ and $\bar{R}\bar{a}^n + \bar{H}_2 = \bar{R}\bar{a}^{n+1} + \bar{H}_2$. And so there are $\bar{x}, \bar{y} \in \bar{R}$ such that $\bar{a}^n - \bar{x}\bar{a}^{n+1} \in \bar{H}_1$ and $\bar{a}^n - \bar{y}\bar{a}^{n+1} \in \bar{H}_2$; hence, $(\bar{a}^n - \bar{x}\bar{a}^{n+1})(\bar{a}^n - \bar{y}\bar{a}^{n+1}) \in \bar{H}_1\bar{H}_2$. Since $\bar{r}\bar{R} = \bar{R}\bar{r}$ for any $\bar{r} \in \bar{R}$ (by Part (2)), it follows that there exist $\bar{y}_1, \bar{y}_2 \in \bar{R}$ such that $\bar{a}^n\bar{y} = \bar{y}_1\bar{a}^n$ and $\bar{a}^{n+1}\bar{y} = \bar{y}_2\bar{a}^{n+1}$. Thus we get that

$$\begin{aligned} & (\bar{a}^n - \bar{x}\bar{a}^{n+1})(\bar{a}^n - \bar{y}\bar{a}^{n+1}) \\ &= \bar{a}^{2n} - \bar{x}\bar{a}^{2n+1} - \bar{a}^n\bar{y}\bar{a}^{n+1} + \bar{x}\bar{a}^{n+1}\bar{y}\bar{a}^{n+1} \\ &= \bar{a}^{2n} - \bar{x}\bar{a}^{2n+1} - \bar{y}_1\bar{a}^{2n+1} + \bar{x}\bar{y}_2\bar{a}^{2n+2} \\ &= \bar{a}^{2n} - \bar{x}_0\bar{a}^{2n+1} \in \bar{H}_1\bar{H}_2, \quad \text{here } \bar{x}_0 = \bar{x} + \bar{y}_1 - \bar{x}\bar{y}_2\bar{a}. \end{aligned} \quad (2)$$

Also

$$\begin{aligned} \bar{H}_1\bar{H}_2 &= (\bar{R}\bar{a}_1 + \bar{H})(\bar{R}\bar{b}_1 + \bar{H}) \\ &= \bar{R}\bar{a}_1\bar{b}_1\bar{R} + \bar{H}\bar{b}_1\bar{R} + \bar{R}\bar{a}_1\bar{H} \\ &\quad + \bar{H}^2 \subseteq \bar{R}\bar{a}_1\bar{b}_1\bar{R} + \bar{H} \subseteq \bar{H}, \end{aligned} \quad (3)$$

and we have that $\bar{a}^{2n} - \bar{x}_0\bar{a}^{2n+1} \in \bar{H}$ and so $\bar{R}\bar{a}^{2n} + \bar{H} = \bar{R}\bar{a}^{2n+1} + \bar{H}$. This is a contradiction, however, because $\bar{H}$ is a maximal element of $\mathcal{S}$. Therefore R is π -regular, and the proof is finished. $\square$

Corollary 10. Assume that R is an Abelian ring such that $\text{Nil}(R)$ is an ideal of R and the one-sided ideals containing $\text{Nil}(R)$ are all ideals. Then the following statements are equivalent.

- (1) R is a π -regular ring.
- (2) $R/E(P)P$ is a metadivision ring for any completely prime ideal P .
- (3) For any completely prime ideals P_1, P_2 , if $E(P_1) = E(P_2)$, then $P_1 = P_2$.
- (4) For any completely prime ideal P , R/P is π -regular.

Proof. (1) $\Rightarrow$ (2). $E(P)P$ is primary ideal of R by Lemma 4 thus $E(P)P$ is a metadivision ring by Lemma 2.

(2) $\Rightarrow$ (1). If P is a completely prime ideal, then, under the natural homomorphism $R \rightarrow R/E(P)P = \bar{R}$, $\bar{P}$ is a completely prime ideal of $\bar{R}$. Because $\bar{R}$ is a metadivision ring, it follows that $\text{Nil}(\bar{R})$ is a unique maximal ideal, and so $\bar{P} \supseteq \text{Nil}(\bar{R})$; thus, $\bar{P} = \text{Nil}(\bar{R})$. We get that $\bar{R}/\bar{P} = \bar{R}/\text{Nil}(\bar{R})$ is a division ring. Hence $R/P \cong \bar{R}/\bar{P}$ is also a division ring and P is a maximal ideal of R . Theorem 9 shows that R is an Abelian π -regular ring.

(1) $\Rightarrow$ (3). Let P_1, P_2 be completely prime ideals of R and $E(P_1) = E(P_2)$. If $P_1 \neq P_2$, then it is no loss to assume that $a \in P_2$ but $a \notin P_1$. Set $a^n = a^n r a^n = a^n e$ where $e = r a^n$ for some $r \in R$ and positive integer n . Since P_1 is a completely prime ideal, it follows that $a^n \notin P_1$ and so $e \notin P_1$. Also $e = r a^n \in P_2$ and $e^2 = e$; we have $e \in E(P_2)$ and so $E(P_1) \neq E(P_2)$. This contradiction shows that $P_1 = P_2$.

(3) $\Rightarrow$ (1). Let P be a completely prime ideal of R ; then there exists a maximal ideal $H \supseteq P$. Note that $H \supseteq \text{Nil}(R)$ since $P \supseteq \text{Nil}(R)$. If H were not completely prime, we may pick $a, b \in R$ with $ab \in H$ such that $a \notin H$ and $b \notin H$. If $a \notin H$ and $b \notin H$, then $Ra + H = R$ and $bR + H = R$. Thus, we get

$$\begin{aligned} R^2 &= (Ra + H)(bR + H) \\ &= RabR + RaH + HbR + H^2 \subseteq RabR + H. \end{aligned} \quad (4)$$

Since $R^2 = R$, $ab \in H$, we have $R \subseteq H$. This contradiction implies that H is a completely prime ideal. Fix $e \in E(H)$, because $1 - e \notin H$, $1 - e \notin P$. Also $(1 - e)e = 0 \in P$; it follows that $e \in P$, and then $E(H) \subseteq E(P)$; thus, $E(H) = E(P)$; hence $H = P$; that is, P is a maximal ideal. Theorem 9 shows that R is an Abelian π -regular ring.

(1) $\Rightarrow$ (4). Every division ring is π -regular and R/P is a division ring by Lemma 7, thus R/P is π -regular.

(4) $\Rightarrow$ (1). By Theorem 9, it suffices to prove that every completely prime ideal is maximal. If P was not maximal, then there exists a maximal ideal $Q \not\supseteq P$. Thus, there exists an $a \in Q$ but $a \notin P$, and so $a^n \notin P$ for all n . Let $\bar{a}^n = \bar{a}^n \bar{r} \bar{a}^n = \bar{a}^n \bar{e}$ for $\bar{r} \in \bar{R} = R/P$. Since $\bar{a}^n \neq \bar{0}$, we get $\bar{e} \neq \bar{0}$ and so $e \notin P$. Since $(1 - e)e = (\bar{1} - \bar{e})\bar{e} = \bar{0}$, we have $(1 - e)e \in P$ and so $1 - e \in P \not\subseteq Q$. Also $e - r a^n \in Q$ and $a \in Q$; it follows that $e \in Q$ and so $1 = 1 - e + e \in Q$. This is a contradiction, which shows P is a maximal ideal, as wanted. $\square$

The following result is Theorem A.

Theorem 11. Assume that R is an Abelian π -regular ring and $\mathcal{M}$ denotes the collection of all completely prime ideals of R . Then R is isomorphic to the subdirect sum of the $R/E(P_\alpha)P_\alpha$'s for all $P_\alpha \in \mathcal{M}$.

Proof. Let $I = \bigcap_{P_\alpha \in \mathcal{M}} E(P_\alpha)P_\alpha$. It suffices to prove that $I = \{0\}$ in order to prove the desired result. Suppose to the contrary that $I \neq \{0\}$. Then there exists $0 \neq x \in I$. We first claim that there exist $e_\alpha \in E(P_\alpha)$ with $e_\alpha x = x$ for any P_α . Since $x \in I$, $x \in E(P_\alpha)P_\alpha$ for all $P_\alpha \in \mathcal{M}$. Let $x = e_1 x_1 + e_2 x_2 + \cdots + e_r x_r$ where $e_i \in E(P_\alpha)$, $x_i \in P_\alpha$, $i = 1, 2, \dots, r$, and then $(1 - e_1) \cdots (1 - e_r) x = 0$. Set $e = 1 - (1 - e_1) \cdots (1 - e_r)$; then $e^2 = e$ and $ex = x$. If $e \notin E(P_\alpha)$, then $e \notin P_\alpha$; thus, $1 - e \in P_\alpha$ (since $(1 - e)e = 0 \in P_\alpha$); hence $(1 - e_1)(1 - e_2) \cdots (1 - e_r) \in P_\alpha$. We get that $1 - e_k \in P_\alpha$ for some $1 \leq k \leq r$, and so $1 = 1 - e_k + e_k \in P_\alpha$, a contradiction. Therefore we have $e \in E(P_\alpha)$.

Set $\mathcal{S} = \{I \mid I \text{ is an ideal of } R, I \supseteq \text{Nil}(R), \text{ and } ex \neq x \text{ for any } e \in E(I)\}$. Since $x \neq 0$, it follows that $\text{Nil}(R) \in \mathcal{S}$ and so $\mathcal{S}$ is not empty. For any ascending series of $\mathcal{S}$, $I_1 \subset I_2 \subset I_3 \subset \cdots$, it is clear that $\bigcup_i I_i \in \mathcal{S}$, and so $\mathcal{S}$ has a maximal element H . We assert that H is a completely prime ideal. If not, then there exist $a, b \in R$ such that $ab \in H$ but neither a nor b lies in H . Applying Lemma 6, both $Ra + H$ and $bR + H$ are ideals of R . Because either a or b does not lie in H , it follows that none of $Ra + H$ or $bR + H$ lies in $\mathcal{S}$, and so there exist idempotent elements $e' \in Ra + H$ and $e'' \in bR + H$ such that $e'x = x$ and $e''x = x$; thus, $e'e''x = e'x = x$. Also $ab \in H$; it follows that

$$\begin{aligned} e'e'' &\in (Ra + H)(bR + H) = RabR \\ &\quad + RaH + HbR + H^2 \subset H. \end{aligned} \quad (5)$$

Moreover $e'e'' \in E(H)$. This is a contradiction since H is a maximal element of $\mathcal{S}$. Hence the assertion is true. By the last paragraph, there is an element $e \in E(H)$ such that $ex = x$, again a contradiction since $H \in \mathcal{S}$. Therefore $I = \{0\}$ and the desired result follows. The proof is complete. $\square$

Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

Acknowledgments

This project is supported by NSF of China (no. 61074005) and by NSF of Liaoning Education Department (no. 2008516).

References

- [1] A. Badawi, "On abelian π -regular rings," *Communications in Algebra*, vol. 25, no. 4, pp. 1009–1021, 1997.
- [2] A. Badawi, "On semicommutative π -regular rings," *Communications in Algebra*, vol. 22, no. 1, pp. 151–157, 1994.
- [3] G. F. Birkenmeier, J. Y. Kim, and J. K. Park, "Regularity conditions and the simplicity of prime factor rings," *Journal of Pure and Applied Algebra*, vol. 115, no. 3, pp. 213–230, 1997.
- [4] G. F. Birkenmeier, J. Y. Kim, and J. K. Park, "A connection between weak regularity and the simplicity of prime factor rings," *Proceedings of the American Mathematical Society*, vol. 122, no. 1, pp. 53–58, 1994.
- [5] N. K. Kim and Y. Lee, "On rings whose prime ideals are completely prime," *Journal of Pure and Applied Algebra*, vol. 170, no. 2-3, pp. 255–265, 2002.
- [6] H. H. Storrer, "Epimorphismen von kommutativen Ringen," *Commentarii Mathematici Helvetici*, vol. 43, no. 1, pp. 378–401, 1968.
- [7] N. H. McCoy, "Generalized regular rings," *Bulletin of the American Mathematical Society*, vol. 45, no. 2, pp. 175–178, 1939.