Research Article

Some Applications of New Modified q -Integral Type Operators

R. P. Pathak and Shiv Kumar Sahoo

Department of Mathematics, National Institute of Technology, G E Road, Raipur 492010, India

Correspondence should be addressed to Shiv Kumar Sahoo; maths10sks@gmail.com

Received 23 March 2015; Revised 29 June 2015; Accepted 12 July 2015

Academic Editor: Ram N. Mohapatra

Copyright © 2015 R. P. Pathak and S. K. Sahoo. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We introduce a new sequence of q -integral operators. We show that it is a weighted approximation process in the polynomial space of continuous functions defined on unit interval. Weighted statistical approximation theorem, Korovkin type theorems for fuzzy continuous functions, and an estimate for the rate of convergence for these operators.

1. Introduction

The study of q -Calculus is a generalization of any subjects, such as hyper geometric series, complex analysis, and particle physics. Currently it continues being an important subject of study. It has been shown that positive linear operators constructed by q -numbers are quite effective as far as the rate of convergence is concerned and we can have some unexpected results, which are not observed for classical case. In the last decade, some new generalizations of well-known positive linear operators, based on q -integers, were introduced and studied by several authors. For example, q -Meyer König and Zeller operators were studied by Trif [1], Dogru and Duman [2], Aral and Gupta [3], and so forth. In 2001, Aral and Gupta [4, 5] introduced a q -generalization of the classical Baskakov operators. In 2012, Sharma [6, 7] introduced the q -Durrmeyer type operators. Orkcu and Dogru [8] introduced Kantorovich type generalization of q -Szász-Mirakjan operators and discussed their A -statistical approximation properties. In this paper motivated by Sharma we introduced a q -analogue of the q -Durrmeyer operators and we study better rate of convergence and statistical approximation properties.

We mention some important definitions of q -Calculus.

Definition 1. For any fixed real numbers $q > 0$ and $k \in \mathbb{N}$, the q -integers are defined by

$$[k]_q = \begin{cases} k, & \text{if } q = 1, \\ 1 + q + q^1 + q^2 + \cdots + q^{k-1}, & \text{if } q \neq 1. \end{cases} \quad (1)$$

In this way for real number n one may write

$$[n]_q = \frac{1 - q^n}{1 - q}; \quad q \neq 1. \quad (2)$$

Definition 2. The q -factorial is defined by

$$[k]_q! = \begin{cases} 1, & \text{if } k = 0, \\ [1]_q \cdot [2]_q \cdots [k]_q, & \text{if } k = 1, 2, \dots \end{cases} \quad (3)$$

Definition 3. For any number $k \in (0, n)$, the q -binomial coefficient is defined by

$$\binom{n}{k}_q = \frac{[n]_q!}{[k]_q! [n-k]_q!}. \quad (4)$$

Aral and Gupta [4] introduced a q -generalization of the classical Baskakov operators. For all $f \in C[0, \infty)$, $q \in (0, 1)$, and each positive integer n , the operators are defined as

$$(B_{n,q}f)(x) = \sum_{k=0}^{\infty} \binom{n+k-1}{k}_q \frac{q^{k(k-1)/2} (kx)^k}{(1+qx)^{n+k}} f\left(\frac{[k]_q}{q^{q-1}[n]_q}\right). \quad (5)$$

For $q = 1$, the above operators become classical Baskakov operators [9].

Deo et al. [10] introduced new version of Bernstein-Durrmeyer type operators defined as follows: for $f \in CI_n$, where $I_n = [0, n/(n+1)]$,

$$(B_{n,q}f)(x) = n \left(1 + \frac{1}{n}\right)^2 \sum_{k=0}^n p_{n,k}(x) \int_0^{n/(n+1)} p_{n,k}(t) f(t) dt, \quad (6)$$

where

$$p_{n,k}(x) = \left(1 + \frac{1}{n}\right)^n \binom{n}{k} x^k \left(\frac{n}{n+1} - x\right)^{n-k}, \quad (7)$$

and they established some approximation results on it.

Sharma [6] introduced the following q -Durrmeyer type operators defined as follows: for $f \in CI_{n,q}$, where $I_{n,q} = [0, [n]_q/[n+1]_q]$,

$$(M_{n,q}^*f)(x) = \frac{[n+1]_q^2}{[n]_q} \sum_{k=0}^n q^{-k} p_{n,k}^*(q; x) \times \dots \times \int_0^{[n]_q/[n+1]_q} p_{n,k}^*(q; qt) f(t) d_q t, \quad (8)$$

where

$$p_{n,k}^*(q; x) = \binom{n}{k}_q \left(\frac{[n+1]_q}{[n]_q} x\right)^k \left(1 - \frac{[n+1]_q}{[n]_q} x\right)_q^{n-k}, \quad (9)$$

and they established some approximation results on it.

In this paper motivated by Sharma [6, 7, 10–12] we introduce a q -analogue of the q -Baskakov-Durrmeyer type operators defined as follows: for $f \in CI_{n,q}$,

$$(M_{n,q}f)(x) = \frac{[n+1]_q^2}{[n]_q} \sum_{k=0}^{\infty} b_{n,k}(q; x) \int_0^{[n]_q/[n+1]_q} p_{n,k}^*(q; qt) f(t) d_q t, \quad (10)$$

where we set $b_{n,k}(q; x) = \binom{n+k-1}{k}_q (q^{(k^2-k-2)/2} x^k / (1+x)_q^{n+k})$.

Kasana et al. [13] obtained a sequence of modified Szász type operators for integrable function on $[0, \infty)$ defined as

$$(M_{n,x}f)(x) \equiv M_{n,x}(f(y); t) = n \sum_{k=0}^{\infty} b_{n,k}(t) \int_0^{\infty} b_{n,k}(y) f(x+y) dy, \quad (11)$$

where x and t belong to $[0, \infty)$ and x is fixed.

In this paper, motivated by Kasana and Sharma, we introduce a q -analogue of the q -Baskakov-Durrmeyer type operators defined as follows: for $f \in CI_{n,q}$,

$$(M_{n,q,x}^*f)(t) = \frac{[n+1]_q^2}{[n]_q} \sum_{k=0}^{\infty} b_{n,k}(q; t) \times \dots \times \int_0^{[n]_q/[n+1]_q} p_{n,k}^*(q; qy) f(x+y) d_q y, \quad (12)$$

where x and t belong to $I_{n,q}$ and x is fixed.

The aim of this paper is to study some approximation properties of a new generalization of operators based on q -integers. We estimate moments for these operators. Also, we study statistical convergence and Korovkin type theorems for fuzzy continuous functions. Finally, we give better error estimations for operators (10) and (12).

2. Estimation of Moments

We use Lemma 5 [6] for $s = 1, 2, \dots$ and, by the definition of q -Beta function, we get $\int_0^{[n]_q/[n+1]_q} p_{n,k}^*(q; qt) t^s d_q t = ([n]_q^{s+1} / [n+1]_q^{s+1}) q^k ([n]_q! [k+s]_q! / [k]_q! [s+n+1]_q!)$.

Theorem 4. Let the sequence of positive linear operators $(M_{n,q}(f))(x)$ be defined by (10). For all $n \in \mathbb{N}$; $q \in (0, 1)$; $f \in CI_{n,q}$; $x \in I_{n,q}$, one gets

$$\begin{aligned} (M_{n,q}1)(x) &= 1, \\ (M_{n,q}t)(x) &= \frac{[n]_q([n]_q x + 1)}{[n+2]_q[n+1]_q}, \\ (M_{n,q}t^2)(x) &= \frac{(1+q)[n]_q^2 + q(1+q)^2 x [n]_q^3}{[n+3]_q[n+2]_q[n+1]_q^2} \\ &\quad + \frac{q^3 x^2 [n]_q^3 [n+1]_q}{[n+3]_q[n+2]_q[n+1]_q^2}. \end{aligned} \quad (13)$$

Proof. By using Lemma 5 and letting $f(t) = 1$ in the operators $M_{n,q}$ be defined by (10), we get

$$\begin{aligned} (M_{n,q}1)(x) &= \frac{[n+1]_q^2}{[n]_q} \sum_{k=0}^{\infty} b_{n,k}(q; x) \int_0^{[n]_q/[n+1]_q} p_{n,k}(q; qt) 1 d_q t \\ &= \frac{[n+1]_q^2}{[n]_q} \sum_{k=0}^{\infty} b_{n,k}(q; x) \frac{[n]_q}{[n+1]_q} q^k \frac{[n]_q!}{[n+1]_q!} \\ &= \sum_{k=0}^{\infty} q^{k(k-2)/2} \binom{n+k-1}{k}_q (x)^k (1+x)_q^{-(n+k)} = 1. \end{aligned} \quad (14)$$

Again, we set $f(t) = t$ in the operators $M_{n,q}$; we get

$$\begin{aligned} (M_{n,q}t)(x) &= \frac{[n+1]_q^2}{[n]_q} \sum_{k=0}^{\infty} b_{n,k}(q; x) \frac{[n]_q^2}{[n+1]_q^2} q^k \frac{[n]_q! [k+1]_q!}{[k]_q! [n+2]_q!} \\ &= \frac{[n]_q}{[n+2]_q [n+1]_q} \sum_{k=0}^{\infty} b_{n,k}(q; x) [k+1]_q \\ &= \frac{[n]_q ([n]_q x + 1)}{[n+2]_q [n+1]_q}. \end{aligned} \quad (15)$$

Similarly, we set $f(t) = t^2$ in the operators $M_{n,q}$; we get

$$\begin{aligned} (M_{n,q}t^2)(x) &= \frac{[n+1]_q^2}{[n]_q} \sum_{k=0}^{\infty} b_{n,k}(q; x) \frac{[n]_q^3}{[n+1]_q^3} q^k \frac{[n]_q! [k+2]_q!}{[k]_q! [n+3]_q!} \\ &= \frac{[n]_q^2 \sum_{k=0}^{\infty} b_{n,k}(q; x) [k+1]_q [k+2]_q}{[n+3]_q [n+2]_q [n+1]_q^2} \\ &= \frac{[n]_q^2 [1+q+q(1+q)^2 [n]_q x]}{[n+3]_q [n+2]_q [n+1]_q^2} \\ &\quad + \frac{[n]_q^2 [q^4 (([n+1]_q [n]_q x^2/q + [n]_q x) - [n]_q x)]}{[n+3]_q [n+2]_q [n+1]_q^2} \\ &= \frac{(1+q) [n]_q^2 + q(1+q)^2 x [n]_q^3 + q^3 x^2 [n]_q^3 [n+1]_q}{[n+3]_q [n+2]_q [n+1]_q^2}. \end{aligned} \quad (16)$$

This completes the proof. $\square$

Lemma 5. For the special case $q = 1$, one gets [14]

$$\begin{aligned} (M_{n,1}1)(x) &= 1; \\ (M_{n,1}t)(x) &= \frac{n^2 x + n}{(n+2)(n+1)}; \\ (M_{n,1}t^2)(x) &= \frac{n^2 [n^2 x^2 + (4nx + x)x + 2]}{(n+3)(n+2)(n+1)^2}. \end{aligned} \quad (17)$$

Lemma 6. For the sequence of positive linear operators $M_{n,q}$, one gets the following central moments: let $\phi^i = (t-x)^i$, $i = 1, 2, \dots$

$$\begin{aligned} (M_{n,q}\phi^1)(x) &= (M_{n,q}t)(x) - x(M_{n,q}1)(x) \\ &= \frac{[n]_q ([n]_q x + 1)}{[n+2]_q [n+1]_q} - x \cdot 1 = \frac{(1-3x) [n]_q - 2x}{[n+2]_q [n+1]_q}; \end{aligned}$$

$$\begin{aligned} (M_{n,q}\phi^2)(x) &= (M_{n,q}t^2)(x) - 2x(M_{n,q}t)(x) + x^2(M_{n,q}1)(x) \\ &= \frac{(1+q) [n]_q^2 + q(1+q)^2 x [n]_q^3 + q^3 x^2 [n]_q^3 [n+1]_q}{[n+3]_q [n+2]_q [n+1]_q^2} \\ &\quad - 2x \frac{[n]_q ([n]_q x + 1)}{[n+2]_q [n+1]_q} + x^2 \cdot 1 \\ &= x^2 \left(1 - \frac{2 [n]_q^2 [n+3]_q + q^3 [n]_q^3}{[n+3]_q [n+2]_q [n+1]_q} \right) \\ &\quad + x \left(\frac{q(1+q)^2 [n]_q^3 - 2 [n]_q [n+3]_q}{[n+3]_q [n+2]_q [n+1]_q^2} \right) \\ &\quad + \frac{(1+q) [n]_q^2}{[n+3]_q [n+2]_q [n+1]_q^2}. \end{aligned} \quad (18)$$

Lemma 7. For the special case $q = 1$, one has the following central moment [14]:

$$\begin{aligned} (M_{n,q}\phi^1)(x) &= \frac{n(1-3x) - 2x}{(n+2)(n+1)} \\ (M_{n,q}\phi^2)(x) &= \frac{2xn^3 + n^2 [11x^2 - 8x + 2]}{(n+3)(n+2)(n+1)^2} \\ &\quad + \frac{n [17x^2 - 6x + 2] + 6x^2}{(n+3)(n+2)(n+1)^2}. \end{aligned} \quad (19)$$

3. Weighted Statistical Approximation Theorem

The aim of this section is to use statistical convergence to study Korovkin type approximation of function f by means of sequence of positive linear operators from a weighted space into a weighted subspace. Let $C_{\rho_0} I_{n,q}$ be the space of the continuous and bounded functions defined on $I_{n,q}$ such that $|f| \leq C(1+x^2)$, where C is a constant depending on f . Our operators acting from similar methods of [3]; we obtain the following results.

Theorem 8. Let sequence $(q_n)_n$, $q_n \in (0, 1)$, such that $st - \lim_{n \rightarrow \infty} q_n = 1$, and the sequence of positive linear operators M_{n,q_n} , $n \in N$, be defined by (10). Then, for all $x \in I_{n,q}$ and $f \in C_{\rho_0} I_{n,q}$, one gets

$$st - \lim_{n \rightarrow \infty} |(M_{n,q_n} f)(x) - f(x)|_{\rho_\alpha} = 0; \quad \alpha > 0. \quad (20)$$

Proof. The weight functions $\rho_0(x)$ and weighted subspace $C_{\rho_0} I_{n,q}$ are defined by $x \in I_{n,q}$; $\alpha > 0$; $\rho_0(x) = 1 + x^2$; $\rho_\alpha(x) = 1 + x^{2+\alpha}$; and $f \in C_{\rho_0} I_{n,q}$ such that f is continuous

on $I_{n,q}$ with norm; here $B_\rho(I_{n,q})$ and $C_\rho(I_{n,q})$ are Banach space. By using Theorem 4, we get

$$\text{st} - \lim_{n \rightarrow \infty} \left| (M_{n,q_n} 1)(x) - 1 \right|_{\rho_0} = 0. \quad (21)$$

Since

$$\begin{aligned} & \left| \frac{(M_{n,q_n} t)(x) - x}{1 + x^2} \right| \\ &= \left| \frac{[n]_q ([n]_q x + 1) / [n+2]_q [n+1]_q - x}{1 + x^2} \right| \\ &\leq \frac{1}{[n]_{q_n}} \end{aligned} \quad (22)$$

and $\text{st} - \lim_{n \rightarrow \infty} q_n = 1$, this implies $\text{st} - \lim_{n \rightarrow \infty} 1/[n]_{q_n} = 0$, and we get

$$\text{st} - \lim_{n \rightarrow \infty} \left| (M_{n,q_n} t)(x) - x \right|_{\rho_0} = 0. \quad (23)$$

Again, since

$$\begin{aligned} \left| \frac{(M_{n,q_n} t^2)(x) - x^2}{1 + x^2} \right| &\leq \frac{1 + q_n}{[n]_{q_n}^2} + \frac{q_n (1 + q_n)^2}{[n]_{q_n}} \\ &\quad + \frac{q^3}{[n]_{q_n}}, \end{aligned} \quad (24)$$

we get

$$\text{st} - \lim_{n \rightarrow \infty} \left| (M_{n,q_n} t^2)(x) - x^2 \right|_{\rho_0} = 0. \quad (25)$$

By using A-statistical convergence theorem given by Duman and Orhan [15], here we let $A = C_1$ ((21), (23), and (25)), and we get $\text{st} - \lim_{n \rightarrow \infty} |(M_{n,q_n} t^k)(x) - x^k|_{\rho_0} = 0$ for $k = 0, 1, 2$ if and only if

$$\text{st} - \lim_{n \rightarrow \infty} \left| (M_{n,q_n} f)(x) - f(x) \right|_{\rho_\alpha} = 0. \quad (26)$$

This completes the proof. $\square$

Theorem 9. Let sequence $(q_n)_n$, $q_n \in (0, 1)$, such that $\text{st} - \lim_{n \rightarrow \infty} q_n = 1$, and the sequence of positive linear operators M_{n,x,q_n}^* , $n \in N$, be defined by (12). Then, for all functions $f \in C_{\rho_0} I_{n,q}$, one gets

$$\text{st} - \lim_{n \rightarrow \infty} \left| (M_{n,x,q_n}^* f)(t) - f(t) \right|_{\rho_\alpha} = 0; \quad \alpha > 0, \quad (27)$$

where x and t belong to I_n and x is fixed.

Proof. The proof is analogous as Theorem 8. $\square$

Theorem 10. Let sequence $(q_n)_n$, $q_n \in (0, 1)$, such that $\text{st} - \lim_{n \rightarrow \infty} q_n = 1$, and the sequence of positive linear operators M_{n,q_n}^* , $n \in N$, be defined by (10). Then, for all $x \in I_n$ and functions $f \in C_{x^2} I_{n,q}$, we get

$$\lim_{n \rightarrow \infty} \left| (M_{n,q_n} f)(x) - f(x) \right|_{x^2} = 0. \quad (28)$$

Proof. To prove the theorem, we use modulus of continuity of f on closed interval I_n given by

$$\omega(f, \delta) = \sup_{|t-x| \leq \delta} \sup_{t \in I_{n,q}} |f(t) - f(x)|. \quad (29)$$

We see that $f \in C_{x^2} I_n$; the modulus of continuity $\omega(f, \delta)$ tends to zero. Consider

$$\begin{aligned} & \left| (M_{n,q_n} t)(x) - x \right|_{x^2} \\ &\leq \frac{[n]_{q_n} - q[n-1]_{q_n}}{[n+2]_{q_n} [n+1]_{q_n}} \sup_{x \in I_{n,q}} \frac{x}{1 + x^2} \\ &\quad + \frac{[n]_{q_n}}{[n+2]_{q_n} [n+1]_{q_n}} \sup_{x \in I_{n,q}} \frac{1}{1 + x^2}, \end{aligned} \quad (30)$$

and we get

$$\lim_{n \rightarrow \infty} \left| (M_{n,q_n} t)(x) - x \right|_{x^2} = 0. \quad (31)$$

Again

$$\begin{aligned} & \left| (M_{n,q_n} t^2)(x) - x^2 \right|_{x^2} \\ &\leq \left(\frac{[n]_{q_n}^3 + q^3 [n]_{q_n}^4}{[n+3]_{q_n} [n+2]_{q_n} [n+1]_{q_n}^2} - 1 \right) \sup_{x \in I_{n,q}} \frac{x^2}{1 + x^2} \\ &\quad + \frac{q(1+q)^2 x [n]_{q_n}^3}{[n+3]_{q_n} [n+2]_{q_n} [n+1]_{q_n}^2} \sup_{x \in I_{n,q}} \frac{x}{1 + x^2} \\ &\quad + \frac{(1+q) [n]_{q_n}^2}{[n+3]_{q_n} [n+2]_{q_n} [n+1]_{q_n}^2} \sup_{x \in I_{n,q}} \frac{1}{1 + x^2}, \end{aligned} \quad (32)$$

and we get

$$\lim_{n \rightarrow \infty} \left| (M_{n,q_n} t^2)(x) - x^2 \right|_{x^2} = 0. \quad (33)$$

By (31) and (33), we get $\lim_{n \rightarrow \infty} |(M_{n,q_n} t^k)(x) - x^k|_{x^2} = 0$ for $k = 0, 1, 2$ if and only if

$$\lim_{n \rightarrow \infty} \left| (M_{n,q_n} f)(x) - f(x) \right|_{x^2} = 0. \quad (34)$$

This completes the proof. $\square$

Theorem 11. Let sequence $(q_n)_n$, $q_n \in (0, 1)$, such that $\text{st} - \lim_{n \rightarrow \infty} q_n = 1$, and the sequence of positive linear operators M_{n,x,q_n}^* , $n \in N$, be defined by (12). Then, for all functions $f \in C_{t^2} I_{n,q}$, one gets

$$\lim_{n \rightarrow \infty} \left| (M_{n,x,q_n}^* f)(t) - f(t) \right|_{t^2} = 0, \quad (35)$$

where x and t belong to $I_{n,q}$ and x is fixed.

Proof. The proof is analogous as Theorem 10. $\square$

4. Korovkin Type Theorems for Fuzzy Continuous Functions

In this section we mention some important definitions given by Burgin and Duman [16].

Definition 12. A number a is called an r -limit of a sequence S , if for any $\epsilon \in \mathbb{R}$, the inequality $|a - a_i| < r + \epsilon$ is valid for almost all a_i , that is, there is such n that for any $i > n$, we have $|a - a_i| < r + \epsilon$. It is denoted by $a = r - \lim S$.

Definition 13. A sequence S that has an r -limit is called r -convergent and it is said that S , r -converges to its r -limit a . It is denoted by $S \rightarrow ra$.

Definition 14. A function $f : R \rightarrow R$ is called r -continuous in $X \subset R$ if $\gamma(f, X) \leq r$ and is called fuzzy continuous in X if $\gamma(f, X) \leq \infty$, where $\gamma(f, X)$ defined as:

$$\gamma(f, X) \geq \inf \{ \sup \{ |f(x) - g(x)| : x \in X \} : g(x) \in C(X) \}. \quad (36)$$

For example the functions $f(x) = x^n$ when $x \in [n, n+1)$, $n \in \mathbb{Z}$ and $g(x) = [x]^n$ are fuzzy continuous in each finite interval of the real line $\mathbb{R}$, but they are not continuous in any interval with the length larger than 1. To define the Riemann integral for a continuous function $f(x)$, step functions are utilized. If the integral of $f(x)$ exists, then any such step function is fuzzy continuous.

Theorem 15. Let a sequence $(q_n)_n$; $q_n \in (0, 1)$ such that $r - \lim_{n \rightarrow \infty} q_n = 1$ and let the sequence of positive linear operators M_{n,q_n} ; $n \in \mathbb{N}$ be defined by (10). If $r_i - \lim_{n \rightarrow \infty} |(M_{n,q_n} e_i)(x) - e_i| = 0$ for $i = 0, 1, 2$. Then for all functions $f \in CI_{n,q}$, we get

$$r - \lim_{n \rightarrow \infty} |(M_{n,q_n} f)(x) - f(x)| = 0, \quad (37)$$

where r is any real number such that $r \geq K_3(r_0 + r_1 + r_2)$ for some $K_3 > 0$.

Proof. Let the functions e_i defined as: $e_i(x) = t^i$ for all $x \in I_{n,q}$. Now, for each $\epsilon > 0$, there corresponds $\delta > 0$ such that $|\lambda(t-x)| \leq \epsilon$ whenever $|t-x| \leq \delta$. Again for $|t-x| > \delta$, then there exist a positive number M such that $|\lambda(t-x)| \leq M \leq M(t-x)^2/\delta^2$. Thus for all t and $x \in I_{n,q}$, we get

$$|\lambda(t-x)| \leq \epsilon + M \frac{(t-x)^2}{\delta^2}. \quad (38)$$

Applying M_{n,q_n} on (38), we get

$$\begin{aligned} & |(M_{n,q_n} f)(x) - f(x)| \\ & \leq \epsilon (M_{n,q_n} e_0)(x) + \frac{M}{\delta^2} (M_{n,q_n} (t-x)^2)(x) \\ & \leq \epsilon + \epsilon |(M_{n,q_n} e_0)(x) - e_0(x)| \\ & \quad + K_3 \sum_{i=0}^2 |(M_{n,q_n} e_i)(x) - e_i(x)|, \end{aligned} \quad (39)$$

where $K_3 = \max\{M/\delta^2, 2Mx/\delta^2, Mx^2/\delta^2\}$. Then for every $\epsilon > 0$ there exist $N = N(\epsilon) > 0$ such that for all $n \in \mathbb{N}$, we get

$$\begin{aligned} & |(M_{n,q_n} f)(x) - f(x)| \\ & \leq \epsilon + \epsilon (r_0 + \epsilon) + K_3 (3\epsilon + r_0 + r_1 + r_2) \leq r + \epsilon_1, \end{aligned} \quad (40)$$

here, $\epsilon_1 = \epsilon(1 + r_0 + \epsilon + 3K_3)$. Since ϵ is arbitrary and small, $r - \lim_{n \rightarrow \infty} q_n = 1$, we get

$$r - \lim_{n \rightarrow \infty} |(M_{n,q_n} f)(x) - f(x)| = 0. \quad (41)$$

This completes the proof. $\square$

Theorem 16. Let a sequence $(q_n)_n$; $q_n \in (0, 1)$ such that $r - \lim_{n \rightarrow \infty} q_n = 1$ and let the sequence of positive linear operators M_{n,q_n}^* ; $n \in \mathbb{N}$ be defined by (12). If $r_i - \lim_{n \rightarrow \infty} |(M_{n,q_n}^* e_i)(x) - e_i| = 0$ for $i = 0, 1, 2$. Then for all functions $f \in CI_{n,q}$, we get

$$r - \lim_{n \rightarrow \infty} |(M_{n,q_n}^* f)(t) - f(t)| = 0, \quad (42)$$

where r is any real number such that $r \geq K_4(r_0 + r_1 + r_2)$ for some $K_4 > 0$.

Proof. The proof is analogous as Theorem 15. $\square$

Theorem 17. Let f be the integrable and bounded in the interval $I_{n,q}$ and let if f'' exists at a point $x \in I_{n,q}$. Let a sequence $(q_n)_n$; $q_n \in (0, 1)$ such that $\lim_{n \rightarrow \infty} q_n = 1$ and let the sequence of positive linear operators M_{n,q_n} ; $n \in \mathbb{N}$ be defined by (10). Then, one gets that

$$\begin{aligned} & \lim_{n \rightarrow \infty} [n]_{q_n} |(M_{n,q_n} f)(x) - f(x)| \\ & = (1 - 3x) f'(x) + x^2 f''(x). \end{aligned} \quad (43)$$

Proof. Let if f'' exists at a point $x \in I_{n,q}$, then by using Taylor's expansion, we write

$$\begin{aligned} f(t) &= f(x) + (t-x) f'(x) + \frac{(t-x)^2}{2} f''(x) \\ & \quad + (t-x)^2 \lambda(t-x), \end{aligned} \quad (44)$$

where $\lambda(t-x) \rightarrow 0$ as $t \rightarrow x$. Applying M_{n,q_n} , we get

$$\begin{aligned} (M_{n,q_n} f)(x) &= f(x) (M_{n,q_n} 1)(x) \\ & \quad + f'(x) (M_{n,q_n} (t-x))(x) \\ & \quad + \frac{f''(x)}{2} (M_{n,q_n} (t-x)^2)(x) \\ & \quad + (M_{n,q_n} (t-x)^2 \lambda(t-x))(x). \end{aligned} \quad (45)$$

By using Theorem 4, and multiplying $[n]_{q_n}$ both sides, we get

$$\begin{aligned} & [n]_{q_n} \left[(M_{n,q_n} f)(x) - f(x) \right] \\ &= f'(x) [n]_{q_n} \left(\frac{(1-3x)[n]_{q_n} - 2x}{[n+2]_{q_n} [n+1]_{q_n}} \right) + \cdots \\ &+ \frac{f''(x) [n]_{q_n}}{2} M_{n,q_n,c} \phi^2(x) + [n]_{q_n} R_{[n]_{q_n}}(t, x). \end{aligned} \quad (46)$$

Here we write,

$$\begin{aligned} & [n]_{q_n} R_{[n]_{q_n}}(t, x) = [n+1]_{q_n}^2 \sum_{k=0}^{\infty} b_{n,k}(q; x) \\ & \cdots \int_0^{[n]_{q_n}/[n+1]_{q_n}} p_{n,k}^*(q_n; q_n t) \phi^2 \lambda \phi d_q t, \\ & \left| [n]_{q_n} R_{[n]_{q_n}}(t, x) \right| \leq [n+1]_{q_n}^2 \sum_{k=0}^{\infty} b_{n,k}(q; x) \\ & \cdots \int_0^{[n]_{q_n}/[n+1]_{q_n}} p_{n,k}^*(q_n; q_n t) |\phi^2 \lambda \phi| d_q t \leq [n]_{q_n} \\ & \cdot \epsilon (M_{n,q_n}(t-x)^2)(x) \\ & + \frac{[n]_{q_n} M}{\delta^2} (M_{n,q_n,c}(t-x)^4)(x) \leq [n]_{q_n} \\ & \cdot \epsilon o\left(\frac{1}{[n]_{q_n}}\right) + \frac{[n]_{q_n} M}{\delta^2} o\left(\frac{1}{[n]_{q_n}^2}\right) \leq \epsilon \\ & + \frac{M}{([n]_{q_n})^{-1/2}} o\left(\frac{1}{[n]_{q_n}}\right) \leq \epsilon + Mo\left(\frac{1}{\sqrt{[n]_{q_n}}}\right); \\ & \text{for } \delta = ([n]_{q_n})^{-1/4}. \end{aligned} \quad (47)$$

Since ϵ is arbitrary and small, $\lim_{n \rightarrow \infty} q_n = 1$ and whenever $n \rightarrow \infty$, we get

$$\left| [n]_{q_n} R_{[n]_{q_n}}(t, x) \right| \rightarrow 0. \quad (48)$$

By using (48), in (46), we get

$$\begin{aligned} & \lim_{n \rightarrow \infty} [n]_{q_n} \left| (M_{n,q_n} f)(x) - f(x) \right| \\ &= (1-3x) f'(x) + x^2 f''(x). \end{aligned} \quad (49)$$

This completes the proof. $\square$

Theorem 18. Let f be the integrable and bounded in the interval $I_{n,q}$ and let if f'' exists at points $x; t \in I_{n,q}$. Let a sequence $(q_n)_n; q_n \in (0, 1)$ such that $\lim_{n \rightarrow \infty} q_n = 1$ and let the sequence of positive linear operators $M_{n,x,q_n}^*; n \in N$ be defined by (12). Then, one gets that

$$\begin{aligned} & \lim_{n \rightarrow \infty} [n]_{q_n} \left| (M_{n,x,q_n}^* f)(t) - f(t) \right| \\ &= (1-3t) f'(x+t) + t^2 f''(x+t). \end{aligned} \quad (50)$$

Proof. The proof is analogous as Theorem 17. $\square$

5. Conclusion

In this way we conclude that weighted statistical approximation theorem, Korovkin type theorems for fuzzy continuous functions, an estimate for the rate of convergence, and some properties are obtained for these operators. For $q = 1$, the operators defined by (8) reduce generalized Baskakov operators. For $q = 1$, the operators defined by (6) reduce generalized Durrmeyer operators (5). By using maple programming, Honey Sharma concludes that Deo's modified operator [10, 11] does not improve the approximation process. In this paper we conclude that, by Deo's modified operators, (10) and (12) improve the approximation process when the value of n is very large, that is, when n tends to infinity.

Conflict of Interests

The authors declare that they have no competing interests.

Authors' Contribution

All authors contributed equally and significantly in writing this paper. All authors read and approved the final paper.

Acknowledgments

The authors are thankful to Director of National Institute of Technology, Raipur (CG), for encouragement. This work was supported and grant funded by the Chhattisgarh Council of Science Technology, Raipur 492001, India. They are also thankful to Director of CCOST, Chhattisgarh, India, for time-to-time eminent support and encouragement.

References

- [1] T. Trif, "Meyer-konig and Zeller operators based on q -integers," *Revue d'Analyse Numérique et de Théorie de l'Approximation*, vol. 29, no. 2, pp. 221–229, 2000.
- [2] O. Dogru and O. Duman, "Statistical approximation of Meyer-konig and Zeller operators based on q -integers," *Publicationes Mathematicae Debrecen*, vol. 68, pp. 199–214, 2006.
- [3] A. Aral and V. Gupta, "On the Durrmeyer type modification of the q -Baskakov type operators," *Nonlinear Analysis. Theory, Methods & Applications*, vol. 72, no. 3-4, pp. 1171–1180, 2010.
- [4] A. Aral and V. Gupta, "Generalized q -Baskakov operators," *Mathematica Slovaca*, vol. 61, no. 4, pp. 619–634, 2011.
- [5] T. Acar, V. Gupta, and A. Aral, "Rate of convergence for generalized Szasz operators," *Bulletin of Mathematical Sciences*, vol. 1, no. 1, pp. 99–113, 2011.
- [6] H. Sharma, "Note on approximation properties of generalized Durrmeyer operators," *Mathematical Sciences*, vol. 6, article 24, 2012.
- [7] H. Sharma and S. J. Aujla, "A certain family of mixed summation integral type Lupas Phillips Bernstein operators," *Mathematical Sciences*, vol. 6, article 26, 2012.
- [8] M. Orku and O. Dogru, "Statistical approximation of a kind of Kantorovich type q -Szasz-Mirakjan operators," *Positivity*, vol. 75, no. 5, pp. 2874–2882, 2012.

- [9] V. A. Baskakov, "An example of a sequence of linear positive operators in the space of continuous functions," *Doklady Akademii Nauk SSSR*, vol. 113, pp. 249–251, 1973.
- [10] N. Deo, M. A. Noor, and M. A. Siddiqui, "On approximation by a class of new Bernstein type operators," *Applied Mathematics and Computation*, vol. 201, no. 1-2, pp. 604–612, 2008.
- [11] N. Deo, "Faster rate of convergence on Srivastava-Gupta operators," *Applied Mathematics and Computation*, vol. 218, no. 21, pp. 10486–10491, 2012.
- [12] N. Deo, N. Bhardwaj, and S. P. Singh, "Simultaneous approximation on generalized Bernstein Durrmeyer operators," *Afrika Matematika*, vol. 24, no. 1, pp. 77–82, 2013.
- [13] H. S. Kasana, G. Prasad, P. N. Agrawal, and A. Sahai, "Modified Szasz operators," in *Proceedings of the International Conference on Mathematical Analysis and Its Applications*, pp. 29–41, Pergamon Press, 1985.
- [14] S. K. Sahoo and S. P. Singh, "Some approximation results on a special class of positive linear operators," *Proceeding of the Mathematical Society, BH University*, vol. 24, pp. 1–9, 2008.
- [15] O. Duman and C. Orhan, "Statistical approximation by positive linear operators," *Studia Mathematica*, vol. 161, no. 2, pp. 187–197, 2004.
- [16] M. Burgin and O. Duman, "Approximations by linear operators in spaces of fuzzy continuous functions," *Positivity*, vol. 15, no. 1, pp. 57–72, 2011.