

ROTHE APPROACH TO THE EXISTENCE OF SOLUTION OF AN ABLATION-TRANSPIRATION COOLING CONTROL SYSTEM¹

GUO BAO ZHU²

*Institute of Systems Science
Academia Sinica
Beijing, CHINA*

ABSTRACT

A one dimensional ablation model with transpiration cooling control and nonlinear effect is studied, which is a distributed parameter control system with moving boundary and both governing equation and boundary conditions involving control variable. A solution is given by a Rothe approximation scheme.

Key words: Ablation-transpiration cooling, distributed parameter control system, moving boundary, Rothe approximation.

AMS (MOS) subject classifications: 49J05, 49L01, 49M99.

1. INTRODUCTION

When a space vehicle flies quickly through the air, the high temperature due to the friction between the front surface of the vehicle and air may cause ablation of the material which will cause damage to the structure of the vehicle. The ablation is even serious for the sideways of launching of the electromagnetic gun. In engineering, a thermal shield must be designed to prevent this event [1-4]. In [1, 2], it is realized by coupling a transpiration cooling control design, which can be demonstrated by a one dimensional version of a solid thermal shield as shown in Figure 1.

¹Received: October, 1992. Revised: March, 1993.

²This work was carried out with the aid of a grant from National Postdoctoral Foundation of China.




Figure 1: Schematic representation of the transpiration cooling control system of a porous medium slab

The thermal shield consists of a porous solid structure of thickness ℓ . The gaseous coolant is inputted at $x = \ell$ and the heated air flow enters at $x = 0$. The coolant flows through the air hole of the slab and enters into the heater air flow $Q(m)$, a specified flow function of the coolant flux $m(t)$ (mass of coolant per unit time flowing through per unit area) on the outer layer of the structure [1, 2]:

$$Q(m) = \bar{q} \Psi, \quad \bar{q} = q_0 \left(1 - \frac{h_w}{h_r}\right), \quad (1)$$

where q_0 is the theoretical heat flow, h_r is the recovery enthalpy, $h_w = c_\rho u_w$ is the enthalpy of outer surface. The blocking coefficient Ψ is different for different coolants:

$$\text{for helium, } \Psi = 1 - 0.724 \frac{h_e}{q_0} m - 0.13 \left(\frac{h_e}{q_0} m\right)^2;$$

$$\text{for water, } \Psi = 1 - 1.04 m \Delta h_i / q_0. \quad (2)$$

When the temperature of the front face exceeds the melting temperature u_m of the material, the outer layer melts and recedes to the new position $x = s(t)$ after time t (Figure 1). Denoting by $u(x, t)$ the temperature of the medium at the space-time point (x, t) and denoting by u_c the temperature of the coolant inside the tank, Yang [1] formulated this process as the following partial differential equation with a moving boundary:

$$\left\{ \begin{array}{l} \frac{\partial u(x, t)}{\partial t} = \alpha^2 \frac{\partial^2 u(x, t)}{\partial x^2} + \beta(t) \frac{\partial u(x, t)}{\partial x} + s'(t) \frac{\ell - x}{\ell - s(t)} \frac{\partial u(x, t)}{\partial x}, \quad s(t) < x < \ell, \\ -\kappa \frac{\partial u(\ell, t)}{\partial x} = m(t) (\Delta h_N - \Delta h_c) = m(t) c_{pL} (u(\ell, t) - u_c), \\ \rho L s'(t) - \kappa \frac{\partial u(s(t), t)}{\partial x} = Q(m(t)), \end{array} \right. \quad (3)$$

where L is the latent heat of melting. Symbols $\Delta h_N = c_{pL}u_N$, $\Delta h_c = c_{pL}u_c$ denote the enthalpy of the coolant at inner face and at tank storing coolant, respectively. The thermal conductivity κ , density ρ and the specific heat c_ρ of the ablation material are constants, c_{pL} is the specific heat of the coolant, $\alpha^2 = \kappa/(\rho c_\rho)$ and $\beta(t) = c_{pL}/(\rho c_\rho)m(t)$.

Since the melted part is broken away (by wind for example) from material immediately after melting, the following Signorini boundary condition is necessary (see [5]):

$$u(s(t), t) \leq u_m, s'(t) \geq 0, [u_m - u(s(t), t)]s'(t) = 0.$$

Observing that $u_w = u(s(t), t)$ and $u_N = u(\ell, t)$, we finally formulate the ablation transpiration cooling control process as the following Stefan-Signorini problem:

$$\left\{ \begin{array}{l} u_t(x, t) = \alpha^2 u_{xx}(x, t) + \beta(t)u_x(x, t) + s'(t)\frac{\ell-x}{\ell-s(t)}u_x(x, t), \quad s(t) < x < \ell, \\ \alpha^2 u_x(\ell, t) + \beta(t)u(\ell, t) = \beta(t)u_c, \quad u_c > 0, \\ u(s(t), t) \leq u_m, \quad s'(t) \geq 0, \quad [u_m - u(s(t), t)]s'(t) = 0, \\ s'(t) = \frac{\kappa}{\rho L}[u_x(s(t), t) - a_1(\beta(t))u(s(t), t) + a_2(\beta(t))], \quad s(0) = 0, \end{array} \right. \quad (4)$$

where

$$a_1(\beta(t)) = \frac{q_0 c_\rho \Psi}{\kappa h_r}, \quad a_2(\beta(t)) = \frac{q_0}{\kappa} \Psi. \quad (5)$$

The initial condition $u(x, 0) = u_0(x)$ and the function $\beta(t)$ should satisfy the consistency condition:

$$\left\{ \begin{array}{l} u_0 \in C^2[0, \ell], \quad u_c \leq u_0 \leq u_m, \quad \beta(t) \in C^1[0, \infty), \quad Q(\beta(t)) \in C^1[0, \infty), \\ u_c \leq u_0(x) \leq u_m, \quad \beta(t) \geq 0, \quad \beta(t) \leq \beta_m, \quad \alpha^2 u'_0(\ell) + \beta(0)u_0(\ell) = u_c \beta(0), \\ u_0(0) \leq u_m, \quad u'_0(0) - a_1(\beta(0))u_0(0) + a_2(\beta(0))u_0(0) \geq 0, \\ [u_0(0) - u_m][u'_0(0) - a_1(\beta(0))u_0(0) + a_2(\beta(0))u_0(0)] = 0. \end{array} \right. \quad (6)$$

To simplify notation, let

$$v(x, t) = u_m - u(x, t), \quad v_0(x) = u_m - u_0(x). \quad (7)$$

Then problem (4) can be written in the form

$$\left\{ \begin{array}{l} v_t(x, t) = \alpha^2 v_{xx}(x, t) + \beta(t)v_x(x, t) + s'(t)\frac{\ell-x}{\ell-s(t)}v_x(x, t), \\ \alpha^2 v_x(\ell, t) + \beta(t)v(\ell, t) = q\beta(t), \quad q = u_m - u_c > 0, \\ v(s(t), t) \geq 0, \quad s'(t) \geq 0, \quad v(s(t), t)s'(t) = 0, \\ v(x, 0) = v_0(x), \\ s'(t) = \frac{\kappa}{\rho L}[-v_x(s(t), t) + a_1(\beta(t))v(s(t), t) + Q(\beta(t))] \end{array} \right. \quad (8)$$

where $Q(\beta(t)) = \frac{q_0}{\kappa}(1 - \frac{c_\rho}{h_r}u_m)\Psi > 0$.

In the paper, the symbol $\|\cdot\|$ is used for the C^0 norm of a continuous function.

2. A FIXED PROBLEM

For any fixed time $T > 0$, and a function

$$s(t) \in C^1[0, T], \quad s(0) = 0, \quad s'(t) \geq 0, \quad s(T) < \ell,$$

the standard transformation

$$y = \frac{x - s(t)}{\ell - s(t)}\ell, \quad \vartheta(y, t) = v(x, t) \quad (10)$$

will map $[s(t), \ell]$ onto $[0, \ell]$ and the fixed problem (8) becomes a fixed domain problem:

$$\left\{ \begin{array}{l} \vartheta_t(y, t) = \alpha^2 \left(\frac{\ell}{\ell - s(t)} \right)^2 \vartheta_{yy}(y, t) + \left[\beta(t) \frac{\ell}{\ell - s(t)} + 2s'(t) \frac{\ell - y}{\ell - s(t)} \right] \vartheta_y(y, t), \\ \alpha^2 \frac{\ell}{\ell - s(t)} \vartheta_y(\ell, t) + \beta(t) \vartheta(\ell, t) = q\beta(t), \\ \vartheta(0, t) \geq 0, \quad -\frac{\ell}{\ell - s(t)} \vartheta_y(0, t) + a_1(\beta(t)) \vartheta(0, t) + Q(\beta(t)) \geq 0, \\ \vartheta(0, t) \left[-\frac{\ell}{\ell - s(t)} \vartheta_y(0, t) + a_1(\beta(t)) \vartheta(0, t) + Q(\beta(t)) \right] = 0. \end{array} \right. \quad (11)$$

Let

$$0 = t_0 < t_1 < t_2 < \dots < t_N = T, \quad t_n - t_{n-1} = \Delta t = \frac{T}{N}$$

be a uniform partition of $[0, T]$. Considering

$$\beta_n = \beta(t_n), \quad s_n = s(t_n),$$

and applying the Rothe approximation scheme to (11), we obtain a boundary value problem as follows:

$$\left\{ \begin{array}{l} \frac{\vartheta_n(y) \vartheta_{n-1}(y)}{\Delta t} = \alpha^2 \left(\frac{\ell}{\ell - s_n} \right)^2 \vartheta_n''(y) + \left[\beta_n \frac{\ell}{\ell - s_n} + 2 \frac{s_n - s_{n-1}}{\Delta t} \frac{\ell - y}{\ell - s_n} \right] \vartheta_n'(y), \\ \alpha^2 \frac{\ell}{\ell - s_n} \vartheta_n'(\ell) + \beta_n \vartheta_n(\ell) = q\beta_n, \quad \vartheta_0(y) = v_0(y), \\ \vartheta_n(0) \geq 0, \quad -\frac{\ell}{\ell - s_n} \vartheta_n'(0) + a_1(\beta_n) \vartheta_n(0) + Q(\beta_n) \geq 0, \\ \vartheta_n(0) \left[-\frac{\ell}{\ell - s_n} \vartheta_n'(0) + a_1(\beta_n) \vartheta_n(0) + Q(\beta_n) \right] = 0. \end{array} \right. \quad (12)$$

Using the transformation

$$x = s_n + \frac{\ell - s_n}{\ell} y, \quad v_n(x) = \vartheta_n(y) \quad (13)$$

we finally obtain the Rothe approximation scheme of the fixed problem (8):

$$\left\{ \begin{array}{l} \frac{v_n(x) - v_{n-1}(\vartheta(x))}{\Delta t} = \alpha^2 v_n''(x) + [\beta_n + 2 \frac{s_n - s_{n-1}}{\Delta t} \frac{\ell - x}{\ell - s_n} v_n'(x)], \quad s_n < x < \ell, \\ \alpha^2 v_n'(\ell) + \beta_n v_n(\ell) = q\beta_n, \\ v_n(s_n) \geq 0, \quad -v_n'(s_n) + a_1(\beta_n)v_n(s_n) + Q(\beta_n) \geq 0, \\ v_n(s_n)[-v_n'(s_n) + a_1(\beta_n)v_n(s_n) + Q(\beta_n)] = 0, \\ v_0(x) = v_m - v_0(x), \quad \vartheta(x) = \frac{\ell - s_{n-1}}{\ell - s_n}x - \frac{s_n - s_{n-1}}{\ell - s_n}\ell. \end{array} \right. \quad (14)$$

Lemma 1: *There exists a unique solution $\{v_n(x), 1 \leq n \leq N\}$ to (14) and*

$$0 \leq v_n(x) \leq q, \quad c_1 \leq v_n'(x) \leq c_2 \quad (15)$$

where

$$c_1 = \min v_0'(x), \quad c_2 = \max\{\|v_0\|, q\beta_M/\alpha^2, q\|a_1\| + \|Q\|\}.$$

Proof: Equation (14) can be solved by the method of invariant imbedding (sweep method) (see [9]):

$$v_n'(x) = R_n(x)v_n(x) + z_n(x) \quad (16)$$

where

$$\left\{ \begin{array}{l} R_n'(x) = -R_n^2(x) - \frac{1}{\alpha^2}[\beta_n + 2 \frac{s_n - s_{n-1}}{\Delta t} \frac{\ell - x}{\ell - s_n}]R_n(x) + \frac{1}{\alpha^2 \Delta t}, \quad R_n(\ell) = -\beta_n/\alpha^2, \\ z_n'(x) = -[R_n(x) + \frac{1}{\alpha^2}(\beta_n + 2 \frac{s_n - s_{n-1}}{\Delta t} \frac{\ell - x}{\ell - s_n})]z_n(x) - \frac{v_{n-1}(\vartheta(x))}{\alpha^2 \Delta t}, \quad z_n(\ell) = q\beta_n/\alpha^2. \end{array} \right. \quad (17)$$

Observe that by defining

$$p(x) = R_n(x) + \frac{1}{\alpha^2}[\beta_n + 2 \frac{s_n - s_{n-1}}{\Delta t} \frac{\ell - x}{\ell - s_n}],$$

where

$$R_n(x) = -\beta_n/\alpha^2 \exp\left(\int_x^\ell p(\tau)d\tau\right) - 1/(\alpha^2 \Delta t) \int_x^\ell \exp\left(\int_x^s p(\tau)d\tau\right) ds < 0,$$

$v_n(x)$ can be determined from (16) with initial condition

$$v_n(s_n) = \max\{0, \frac{Q(\beta_n) - z_n(s_n)}{-R_n(s_n) + a_1(\beta_n)}\}. \quad (18)$$

If $\pm v_n$ attains its maximum at $x_0 \in (s_n, \ell)$, then

$$\pm v_n(x_0) \leq \pm v_{n-1}(\vartheta(x_0)).$$

If v_n attains its negative minimum at $x = \ell$, by the maximum principle (see [10]), $v'_n(\ell) < 0$, and hence $v_n(\ell) > q$ from the boundary condition at $x = \ell$. Therefore, we have a contradiction. If v_n attains its positive maximum at $x = s_n$, then $v'_n(s_n) < 0$ and hence $v_n(s_n) = 0$ by the boundary condition at $x = s_n$. Therefore, we obtain a contradiction. If v_n attains its positive maximum at $x = \ell$, by the maximum principle, $v'_n(\ell) > 0$, and hence $v_n(\ell) < q$ from the boundary condition at $x = \ell$. Notice that if $v_n(s_n) \geq 0$ and $0 \leq v_0(x) \leq q$, then $0 \leq v_n(x) \leq q$. Next, let $\phi_n(x) = v'_n(x)$. Then

$$\left\{ \begin{array}{l} \alpha^2 \phi_n''(x) + [\beta_n + 2 \frac{s_n - s_{n-1}}{\Delta t} \frac{\ell - x}{\ell - s_n}] \phi_n'(x) - [\frac{1}{\Delta t} + 2 \frac{s_n - s_{n-1}}{\Delta t} \frac{1}{\ell - s_n}] \phi_n(x) \\ \quad = \frac{-\phi_{n-1}(\vartheta(x)) \ell - s_{n-1}}{\Delta t \ell - s_n}, \\ 0 \leq \alpha^2 \phi_n(\ell) = \beta_n(q - v_n(\ell)) \leq \beta_M q, \\ 0 \leq \phi_n(s_n) \leq a_1(\beta_n)v_n(s_n) + Q(\beta_n) \leq q \|a_1\| + \|Q\|. \end{array} \right.$$

The second inequality of (15) can be obtained by a direct application of the maximum principle for elliptic ordinary differential equations (see [10]). The proof is complete.

Next, take a family of nonpositive functions $\beta_\epsilon \in C^2(-\infty, \infty)$ with $\beta'_\epsilon \leq 0$, $\epsilon > 0$, and

$$\beta_\epsilon(t) = \begin{cases} 0, & \text{if } t \geq 0, \\ (t + \epsilon)/\epsilon, & \text{if } t < -2\epsilon, \end{cases} \quad (19)$$

and consider a penalized problem of (14):

$$\left\{ \begin{array}{l} \frac{v_{\epsilon n}(x) - v_{\epsilon(n-1)}(\vartheta(x))}{\Delta t} = \alpha^2 v_{\epsilon n}''(x) + [\beta_n + 2 \frac{s_n - s_{n-1}}{\Delta t} \frac{\ell - x}{\ell - s_n}] v_{\epsilon n}'(x), \quad s_n < x < \ell, \\ \alpha^2 v_{\epsilon n}'(\ell) + \beta_n v_{\epsilon n}(\ell) = q \beta_n, \\ -v_{\epsilon n}'(s_n) + a_1(\beta_n)v_{\epsilon n}(s_n) + Q(\beta_n) + \beta_\epsilon(v_{\epsilon n}(s_n)) = 0, \\ v_{\epsilon 0}(x) = v_0(x), \quad \vartheta(x) = \frac{\ell - s_{n-1}}{\ell - s_n} x - \frac{s_n - s_{n-1}}{\ell - s_n} \ell. \end{array} \right. \quad (20)$$

Note: $\vartheta(s_n) = s_{n-1}$, $\vartheta(\ell) = \ell$.

Lemma 2: *There exists a unique solution to (20) and the following estimates hold:*

- (i) $-c_0\epsilon \leq v_{\epsilon n}(x) \leq q$, $c_0 = \max\{2, 1 + \|Q\|\}$;
- (ii) $c_1 \leq v_{\epsilon n}'(x) \leq c_2$

$$c_1 = \max\{\min v'_0(x), (c_0 \|a_1\| + 1 - c_0)\epsilon\},$$

$$c_2 = \max\{q \|a_1\| + \|Q\|, (q\beta_M + c_0\epsilon)/\alpha^2, \max v'_0(x)\};$$

$$(iii) \quad |v_{\epsilon n}(x) - v_{\epsilon(n-1)}(x)| \leq C\Delta t, \quad \|v''_{\epsilon n}\| \leq C,$$

where C is a constant depending only on the C^1 norm of $\beta(t)$, $s(t)$ and $\ell - s(T)$, and the C^2 norm of v_0 .

Proof: Again, we solve Equation (20) by the method of invariant imbedding. The solution is related by the Riccati transformation:

$$v'_{\epsilon n}(x) = R_n(x)v_{\epsilon n}(x) + z_n(x). \tag{21}$$

Substituting $v'_{\epsilon n}(x) = R_n(x)v_{\epsilon n}(x) + z_n(x)$ into the boundary condition of (20) at s_n , we obtain

$$(-R_n(s_n) + a_1(\beta_n))v_{\epsilon n}(s_n) - z_n(s_n) + Q(\beta_n) + \beta'_\epsilon(v_{\epsilon n}(s_n)) = 0.$$

Define

$$f(t) = (-R_n(s_n) + a_1(\beta_n))t - z_n(s_n) + Q(\beta_n) + \beta'_\epsilon(t).$$

Then

$$f'(t) = (-R_n(s_n) + a_1(\beta_n)) + \beta'_\epsilon(t) > 0,$$

$$f(+\infty) = \infty, \quad f(-\infty) = -\infty,$$

and hence there exists a unique t_0 such that $f(t_0) = 0$. Therefore, Equation (20) has a unique solution with $v_{\epsilon n}(s_n) = t_0$.

Now, we shall use mathematical induction. For $n = 0$, the conclusions (i) and (ii) hold. Suppose that it is correct for $k < n$. We shall consider the case of $k = n$. By the maximum principle from [10], if $v_{\epsilon n}$ attains its negative minimum at x_0 , then either $v_{\epsilon n}(x_0) \geq v_{\epsilon(n-1)}(v(x_0))$ or $Q(\beta_n) + \beta'_\epsilon(v_{\epsilon n}(x_0)) \geq 0$ depending on whether x_0 lies on the interior of (s_n, ℓ) or on its boundaries. So

$$\text{either } v_{\epsilon n}(x) \geq -2\epsilon$$

$$\text{or } v_{\epsilon n}(x) \geq -[1 + Q(\beta_n)]\epsilon \geq -[1 + \|Q\|]\epsilon,$$

since $v_0(x) \geq 0$. This is the first part of (i). If $v_{\epsilon n}$ attains its positive maximum at x_0 , then $v_{\epsilon n}(x_0) \leq v_{\epsilon(n-1)}(v(x_0))$, $v_{\epsilon n}(x_0) \leq q$, or $v_{\epsilon n}(x_0) \leq 0$. This is the second part of (i). To prove (ii), let $\psi_n(x) = v'_{\epsilon n}(x)$. Then

$$\left\{ \begin{aligned} & \alpha^2 \psi_n''(x) + [\beta_n + 2 \frac{s_n - s_{n-1}}{\Delta t} \frac{\ell - x}{\ell - s_n}] \psi_n'(x) - [\frac{1}{\Delta t} + 2 \frac{s_n - s_{n-1}}{\Delta t} \frac{1}{\ell - s_n}] \psi_n(x) \\ & \quad = \frac{-\psi_{(n-1)}(\vartheta(x))}{\Delta t} \frac{\ell - s_{n-1}}{\ell - s_n}, \\ & \alpha^2 \psi_n(\ell) = \beta_n(q - v_{\epsilon n}(\ell)), \\ & \psi_n(s_n) = a_1(\beta_n)v_{\epsilon n}(s_n) + Q(\beta_n) + \beta_\epsilon(v_{\epsilon n}(s_n)). \end{aligned} \right. \quad (22)$$

Using a similar argument as above, we get the second part of (ii). Now, we shall prove (iii).

For this purpose, let

$$\eta_n(x) = \pm v_{\epsilon n}''(x) e^{-\chi(x-\ell-\delta)^2}, \quad \delta = \frac{\ell - s_n}{2}, \quad \chi > 0.$$

Then $\eta_n(x)$ satisfies the following equation:

$$\begin{aligned} & \alpha^2 \eta_n''(x) + [4\chi\alpha^2(x-\ell+\delta) + \beta_n + 2 \frac{s_n - s_{n-1}}{\Delta t} \frac{\ell - x}{\ell - s_n}] \eta_n'(x) - [\frac{1}{\Delta t} - 2\chi\alpha^2 \\ & - 4\chi^2\alpha^2(x-\ell+\delta)^2 - 2\chi(x-\ell+\delta)(\beta_n + 2 \frac{s_n - s_{n-1}}{\Delta t} \frac{\ell - x}{\ell - s_n}) + 4 \frac{s_n - s_{n-1}}{\Delta t} \frac{1}{\ell - s_n}] \eta_n(x) \\ & \quad = \mp \frac{1}{\Delta t} \eta_{n-1}(\vartheta(x)) \pm \eta_{n-1}(\vartheta(x)) K(x), \end{aligned} \quad (23)$$

where

$$K(x) = 1 - e^{\chi(\vartheta(x) - \ell - \delta)^2 - \chi(x - \ell - \delta)^2} \left(\frac{\ell - s_{n-1}}{\ell - s_n} \right),$$

and the following boundary conditions:

$$\begin{aligned} & \alpha^2 \eta_n'(\ell) + (\beta_n + \beta_{n-1} + 2\chi\delta\alpha^2) \eta_n(\ell) \\ & = \pm \left[[2 \frac{s_n - s_{n-1}}{\Delta t} \frac{1}{\ell - s_n} - \beta_n \beta_{n-1} / \alpha^2] v_{\epsilon n}'(\ell) - \frac{s_n - s_{n-1}}{\Delta t} \frac{1}{\ell - s_n} v_{\epsilon(n-1)}'(\ell) \right. \\ & \quad \left. + \frac{\beta_n - \beta_{n-1}}{\Delta t} (q - v_{\epsilon n}(\ell)) / \alpha^2 \right] e^{-\chi\delta^2} = K_2(\epsilon), \\ & \alpha^2 \eta_n'(s_n) - [2\chi\alpha^2(\ell - s_n - \delta) + \alpha^2 \beta_\epsilon'(\xi) + \alpha^2 a_1(\beta_{n-1}) - \beta_n - 2 \frac{s_n - s_{n-1}}{\Delta t}] \eta_n(s_n) \\ & = \pm \left[[\frac{s_n - s_{n-1}}{\Delta t} \frac{1}{\ell - s_n} + (a_1(\beta_{n-1}) + \beta_\epsilon'(\xi)) (\beta_n + 2 \frac{s_n - s_{n-1}}{\Delta t}) \right] v_{\epsilon n}'(s_n) \\ & \quad + \left(\beta_\epsilon'(\vartheta) + a_1(\beta_n) + \frac{s_n - s_{n-1}}{\Delta t} \frac{1}{\ell - s_n} \right) v_{\epsilon n}'(s_n) - \frac{s_n - s_{n-1}}{\Delta t} \frac{1}{\ell - s_n} v_{\epsilon(n-1)}'(s_{n-1}) \\ & \quad + \frac{a_1(\beta_n) - a_1(\beta_{n-1})}{\Delta t} v_{\epsilon n}(s_n) + \frac{Q(\beta_n) - Q(\beta_{n-1})}{\Delta t} \right] e^{-\chi(s_n - \ell - \delta)^2} = K_3(\epsilon), \end{aligned}$$

where $-c_0\epsilon \leq \xi \leq q$ and $\chi > 0$ is such that

$$2\chi\delta\alpha^2 \geq 1, \quad 2\chi\alpha^2(\ell - s_n - \delta) - \beta_n - 2 \frac{s_n - s_{n-1}}{\Delta t} \geq 1.$$

Suppose that $\eta_{n-1}(x) \leq C_{n-1}$. Then if η_n attains its maximum in the interior of (s_n, ℓ) , then by the inequality $|K(x)| \leq K_0 \Delta t$, we obtain

$$\eta_n(x) \leq C_{n-1} + K_{10} \Delta t C_{n-1}, \text{ where } K_{10} > 0 \text{ is a constant,}$$

and hence

$$\eta_n(x) \leq (1 + K_{10} \Delta t)^N C_0, \quad \Delta t = T/N.$$

If $\eta_n(x)$ attains its maximum at $x = \ell$ or $x = s_n$ then by (i) and (ii),

$$\eta_n(x) \leq K_2(\epsilon) \leq K_{20}, \quad \eta_n(x) \leq K_3(\epsilon) \leq K_{30}.$$

Since

$$\lim_{t \rightarrow \infty} (1 + K_{10} \Delta t)^N = K_{10} T$$

we, finally, get from (20) that $\eta_n(x) \leq C$ or $|v''_{\epsilon n}| \leq C$. Since

$$\frac{v_{\epsilon n}(x) - v_{\epsilon(n-1)}(x)}{\Delta t} = \alpha^2 v''_{\epsilon n}(x) + [\beta_n + 2 \frac{s_n - s_{n-1}}{\Delta t} \frac{\ell - x}{\ell - s_n}] v'_{\epsilon n}(x) - v'_{\epsilon(n-1)}(\xi) \frac{x - \vartheta}{\Delta t},$$

$$\xi \in (\vartheta(x), x), \quad \frac{x - \vartheta}{\Delta t} = \frac{s_n - s_{n-1}}{\Delta t} \frac{\ell - x}{\ell - s_n},$$

we have $\left| \frac{v_{\epsilon n}(x) - v_{\epsilon(n-1)}(x)}{\Delta t} \right| \leq C$. The proof is complete.

Since the constant C in Lemma 2 is independent on ϵ , by the Ascoli-Arzelà theorem applied for any integer n , there exists a subsequence of $\{v_{\epsilon n}(x), v'_{\epsilon n}(x)\}$, which for easier notation is still denoted by $v_{\epsilon n}$, such that

$$v_{\epsilon n} \rightarrow v_n, \quad v'_{\epsilon n} \rightarrow v'_n, \quad \text{uniformly as } \epsilon \rightarrow 0,$$

and, from equation (20),

$$v''_{\epsilon n} \rightarrow v''_n, \quad \text{uniformly as } \epsilon \rightarrow 0.$$

If $v_n(x_0) > 0$, then by uniform convergence, for sufficiently small ϵ , there exists a constant $c > 0$ such that $v_{\epsilon n}(x_0) \geq c > 0$. Therefore, from (20), $-v'_{\epsilon n}(s_n) + a_1(\beta_n)v_{\epsilon n}(s_n) + Q(\beta_n) = 0$, and hence $-v'_n(s_n) + a_1(\beta_n)v_n(s_n) + Q(\beta_n) = 0$, i.e. $v_n(x)$ is the solution of (14). Since the solution of (14) is unique, the whole sequence of solutions of (20) converges uniformly to the solution of (14) as ϵ goes to zero.

We thus have proved the following lemma.

Lemma 3: *The solution of Equation (14) satisfies the inequality*

$$|v_n(x) - v_{n-1}(x)| \leq C\Delta t, \quad \|v_n''\| \leq C,$$

where C is a constant depending only on the C^1 norm of $\beta(t)$, $s(t)$, $\ell - s(T)$, and the C^2 norm of v_0 .

Define

$$\left\{ \begin{array}{l} s_N(t) = \frac{1}{\Delta t}[(t - t_{n-1})s_n + (t_n - t)s_{n-1}] \text{ for } t \in [t_{n-1}, t_n], \\ v_N(x, t) = \frac{1}{\Delta t}[(t - t_{n-1})v_n(x) + (t_n - t)v_{n-1}(x)] \text{ for } t \in [t_{n-1}, t_n]. \end{array} \right. \quad (24)$$

Lemma 4:

- (a) $0 \leq s_N'(t) \leq \|s'\|$;
- (b) $0 \leq v_N(x, t) \leq q$;
- (c) $|v_{N_x}(x, t)| \leq \max\{\|v_0\|, q\beta_M/\alpha^2, q\|a_1\| + \|Q\|\}$;
- (d) $v_{N_x}(x, t) \geq 0$, if $u_0'(x) \leq 0$;
- (e) $|v_{N_{xx}}(x, t)| \leq C$;
- (f) $|v_{N_x}(x, t_1)| - |v_{N_x}(x, t_2)| \leq C|t_1 - t_2|^{1/2}$ for any $t_1, t_2 \in [t_{n-1}, t_n]$,

where C has the same meaning as in Lemma 3.

Proof: Conditions (a)-(e) are direct consequences of Lemma 1 and Lemma 3, and the definition of $v_N(x, t)$. Now, we shall only consider condition (f). Notice that

$$v_{N_{xt}}(x, t) = \frac{v_n'(x) - v_{n-1}'(x)}{\Delta t}$$

and

$$\begin{aligned} v_{N_{xt}}(x, t) &= \alpha^2 v_n'''(x) + [\beta_n + 2\frac{s_n - s_{n-1}}{\Delta t} \frac{\ell - x}{\ell - s_n}] v_n''(x) - [2\frac{s_n - s_{n-1}}{\Delta t} \frac{1}{\ell - s_n}] v_n'(x) \\ &\quad - v_{n-1}'(\xi) \frac{s_n - s_{n-1}}{\Delta t} \frac{\ell - x}{\ell - s_n} = \alpha^2 v_n'''(x) + G_n(x), \end{aligned}$$

where $|G_n(x)| \leq C$, $\xi \in (\vartheta(x), x)$, $t \in [t_{n-1}, t_n]$. For any $t_{n-1} \leq t_1 < t_2 < t_n$, $\tau = (t_2 - t_1)^{1/2}$,

$$\begin{aligned} \int_x^{x+\tau} [v_{N_x}(y, t_1) - v_{N_x}(y, t_2)] dy &= \int_x^{x+\tau} \int_{t_1}^{t_2} v_{N_{xt}}(y, t) dy dt \\ &= \left[\alpha^2 [v_n''(x+\tau) - v_n''(x)] + \int_x^{x+\tau} G_n(y) dy \right] (t_2 - t_1). \end{aligned}$$

Hence there exists a point $\xi \in (x, x + \tau)$ such that

$$|v_{N_x}(\xi, t_1) - v_{N_x}(\xi, t_2)| \leq C |t_1 - t_2|^{1/2}.$$

Thus

$$\begin{aligned} & |v_{N_x}(x, t_1) - v_{N_x}(x, t_2)| \leq |v_{N_x}(x, t_1) - v_{N_x}(\xi, t_1)| \\ & + |v_{N_x}(x, t_2) - v_{N_x}(\xi, t_2)| + C |t_1 - t_2|^{1/2} \leq (C + 2 \|v_{N_{xx}}\|) |t_1 - t_2|^{1/2}, \end{aligned}$$

and consequently (f) holds. The proof is complete.

Since $s'(t)$ is a continuous function on $[0, T]$,

$$\lim_{N \rightarrow \infty} \|s_N - s\| = 0, \quad \lim_{N \rightarrow \infty} \max_n \left\| \frac{s_n - s_{n-1}}{\Delta t} - s'(t) \right\|_{c[t_{n-1}, t_n]} = 0. \quad (25)$$

By the Ascoli-Arzelà theorem, there exist subsequences of $v_N(x, t)$, $s_N(t)$, which for easier notation are still denoted by v_N , s_N , and a function $\phi(x, t)$ such that

$$\begin{aligned} & s_N \rightarrow s, \quad v_N \rightarrow \phi, \quad v_{N_x} \rightarrow \phi_x \text{ uniformly as } N \rightarrow \infty, \\ & v_{N_t} \rightarrow \phi_t, \quad v_{N_{xx}} \rightarrow \phi_{xx} \text{ in } L^\infty \text{ weak star topology,} \end{aligned} \quad (26)$$

and

$$\begin{aligned} & 0 \leq \phi \leq q, \quad |\phi_x(x, t)| \leq C_0, \quad |\phi_t(x, t)| \leq C, \quad |\phi_{xx}(x, t)| \leq C, \\ & |\phi_x(x, t_1)| - |\phi_x(x, t_2)| \leq C |t_1 - t_2|^{1/2}, \text{ for any } t_1, t_2 \in [0, T], \end{aligned}$$

where C is the constant from Lemma 4.

Notice that

$$\begin{aligned} & -v'_N(s_N(t_n), t_n) + a_1(\beta(t_n))v_N(s_N(t_n), t_n) + Q(\beta(t_n)) \geq 0, \\ & [v_N(s_N(t_n), t_n)][-v'_N(s_N(t_n), t_n) + a_1(\beta(t_n))v_N(s_N(t_n), t_n) + Q(\beta(t_n))] = 0, \\ & \alpha^2 v'_N(\ell, t_n) + \beta(t_n)v_N(\ell, t_n) = q\beta(t_n), \end{aligned}$$

at each point of the partition $\{t_n\}$ associated with N . Since these points are dense in $[0, T]$ as $N \rightarrow \infty$ and the convergence of $v_{N_x}(x, t)$, $v_N(x, t)$, $s_N(t)$ is uniform, it follows immediately that

$$\left\{ \begin{aligned} & -\phi_x(s(t), t) + a_1(\beta(t))\phi(s(t), t) + Q(\beta(t)) \geq 0, \\ & \phi(s(t), t)[- \phi'(s(t), t) + a_1(\beta(t))\phi(s(t), t) + Q(\beta(t))] = 0, \\ & \alpha^2 \phi_x(\ell, t) + \beta(t)\phi(\ell, t) = q\beta(t). \end{aligned} \right. \quad (27)$$

We shall show that $\phi(x, t) = v(x, t)$ is the solution of fixed problem (8). In fact, since

$$\begin{aligned}
v_{Nt}(x, t) &= \alpha^2 v''_n(x) + [\beta_n + \frac{s_n - s_{n-1}}{\Delta t} \frac{\ell - x}{\ell - s_n}] v'_n(x) + K_n(x) \\
&= \alpha^2 v_{Nxx}(x, t_n) + [\beta(t_n) + s'_N(t) \frac{\ell - x}{\ell - s_N(t_n)}] v_{Nx}(x, t_n) + K_n(x), \\
s_N(t_n) &< x < \ell, t \in (t_{n-1}, t_n), n \geq 1,
\end{aligned}$$

where

$$\begin{aligned}
K_n(x) &= [v'_n(x) - v'_{n-1}(\xi)] \frac{s_n - s_{n-1}}{\Delta t} \frac{\ell - x}{\ell - s_n} = [v'_n(x) - v'_{n-1}(x)] \frac{s_n - s_{n-1}}{\Delta t} \frac{\ell - x}{\ell - s_n} \\
&+ v''_{n-1}(\Theta) \frac{s_n - s_{n-1}}{\Delta t} \frac{\ell - x}{\ell - s_n} (x - \xi), \xi, \Theta \in (\vartheta(x), x), \frac{x - \vartheta}{\Delta t} = \frac{s_n - s_{n-1}}{\Delta t} \frac{\ell - x}{\ell - s_n}, \\
|K_n(x)| &\leq C \Delta t^{1/2}.
\end{aligned}$$

For any $f(x, t) \in C^2(\Omega)$, $\Omega = \{(x, t), t \in [0, T], x \in [s(t), \ell]\}$,

$$\begin{aligned}
\sum_{n=1}^N \Delta t \int_{s_N(t_n)}^{\ell} v_{Nt}(x, t) f(x, t_n) dx &= \sum_{n=1}^N \Delta t \int_{s_N(t_n)}^{\ell} \alpha^2 v_{Nxx}(x, t_n) f(x, t_n) dx \\
&+ \sum_{n=1}^N \Delta t \int_{s_N(t_n)}^{\ell} [\beta(t_n) + s'_N(t_n) \frac{\ell - x}{\ell - s_N(t_n)}] v_{Nx}(x, t_n) f(x, t_n) dx + G_N,
\end{aligned}$$

where

$$\begin{aligned}
|G_N| &\leq \left| \sum_{n=1}^N \Delta t \int_{s_N(t_n)}^{\ell} K_n(x) f(x, t_n) dx \right| \leq CT(\ell - s(T)) \|f\| \Delta t^{1/2}, \\
\lim_{N \rightarrow \infty} G_N &= 0.
\end{aligned}$$

By using integration by parts and by uniform convergence of (25) and (26), we can obtain

$$\begin{aligned}
& - \int_{\Omega} \int f_t \phi dx dt + \int_0^T s'(t) [\phi f](s(t), t) dt + \int_{s(T)}^{\ell} [f \phi](x, T) dx - \int_0^{\ell} f(x, 0) v_0(x) dx \\
&= \alpha^2 \int_0^T [[\phi_x f](l, t) - [\phi_x f](s(t), t)] dt - \alpha^2 \int_{\Omega} \phi_x f_x dx dt \\
&+ \int_{\Omega} [\beta(t) + s'(t) \frac{\ell - x}{\ell - s(t)}] \phi_x f dx dt, \tag{28}
\end{aligned}$$

and (28) holds for any continuous function f with first order derivatives bounded on Ω by smooth function approximation.

From Equation (8) it also holds for its solution v that

$$\begin{aligned}
 & - \int_{\Omega} \int_0^T f_t v dx dt + \int_0^T s'(t) [vf](s(t), t) dt + \int_{s(T)}^{\ell} [fv](x, T) dx - \int_0^{\ell} f(x, 0) v_0(x) dx \\
 & = \alpha^2 \int_0^T [[v_x f](l, t) - [v_x f](s(t), t)] dt - \alpha^2 \int_{\Omega} \int v_x f_x dx dt \\
 & \quad + \int_{\Omega} \int [\beta(t) + s'(t) \frac{\ell - x}{\ell - s(t)}] v_x f dx dt. \tag{29}
 \end{aligned}$$

Let $\delta > \|\frac{s'(t)}{\ell - s(t)}\|$, $f = (v - \phi)e^{-\delta t}$ in (28) and $f = (\phi - v)e^{-\delta t}$ in (29). After these substitutions, and then adding (28) and (29), we obtain

$$\begin{aligned}
 & \alpha^2 \int_{\Omega} \int (v - \phi)_x^2 e^{-\delta t} dx dt + 1/2 e^{-\delta T} \int_{s(T)}^{\ell} (v - \phi)^2(x, T) dx \\
 & + 1/2 \int_0^T \beta(t) (v - \phi)^2(l, t) e^{-\delta t} dt + \int_0^T s'(t) (v - \phi)^2(s(t), t) e^{-\delta t} dt \\
 & + 1/2 \int_0^T \beta(t) (v - \phi)^2(s(t), t) e^{-\delta t} dt + 1/2 \int_{\Omega} \int (\delta - \frac{s'(t)}{\ell - s(t)}) (v - \phi)^2 e^{-\delta t} dx dt \\
 & = -\alpha^2 \int_0^T (v - \phi)(v - \phi)_x(s(t), t) dt \leq 0.
 \end{aligned}$$

The last inequality comes from the fact that

$$(v - \phi)(v - \phi)_x(s(t), t) dt \leq 0,$$

by the condition at $x = s(t)$. Hence $\phi = v$. This demonstration implies that the (weak) solution of fixed problem (8) is unique and, hence, that (26) holds for the whole sequence v_N . Thus we have proved the following:

Theorem 1: *Let $s_N(t)$ and $v_N(x, t)$ be defined as in (24), where $v_n(x)$ is produced by the Rothe approximation scheme (14) for fixed problem (8). Then*

$$\begin{aligned}
 & v_N \rightarrow v, v_{Nx} \rightarrow v_x \text{ uniformly as } N \rightarrow \infty, \\
 & v_{Nt} \rightarrow v_t, v_{Nx} \rightarrow v_{xx} \text{ in } L^\infty \text{ weak star topology,}
 \end{aligned}$$

where v is the unique solution of Equation (8), and v satisfies

$$0 \leq v \leq q, |v_x(x, t)| \leq C_0, |v_t(x, t)| \leq C, |v_{xx}(x, t)| \leq C,$$

$$|v_x(x, t_1)| - |v_x(x, t_2)| \leq C |t_1 - t_2|^{1/2} \text{ for any } t_1, t_2 \in [0, T],$$

where C depends only on the C^1 norm of β , s , $\ell - s(T)$, and the C^2 norm of v_0 .

3. EXISTENCE OF THE SOLUTION

This section will contribute to the existence of solution of Equations (8)-(9). For this purpose, we construct the Rothe approximation scheme of Equations (8)-(9) as follows:

$$\left\{ \begin{array}{l} \frac{v_n(x) - v_{n-1}(\vartheta(x))}{\Delta t} = \alpha^2 v_n''(x) + [\beta_n + 2 \frac{s_n - s_{n-1}}{\Delta t} \frac{\ell - x}{\ell - s_n}] v_n'(x), \quad s_n < x < \ell, \\ \alpha^2 v_n'(\ell) + \beta_n v_n(\ell) = q \beta_n, \\ v_n(s_n) \geq 0, \quad -v_n'(s_n) + a_1(\beta_n) v_n(s_n) + Q(\beta_n) \geq 0, \\ v_n(s_n) [-v_n'(s_n) + a_1(\beta_n) v_n(s_n) + Q(\beta_n)] = 0, \quad n = 1, 2, \dots, N, \\ v_0(x) = v_m - v_0(x), \quad \vartheta(x) = \frac{\ell - s_{n-1}}{\ell - s_n} x - \frac{s_n - s_{n-1}}{\ell - s_n} \ell, \\ s_n = s_{n-1} + \frac{\kappa}{\rho L} [-v_{n-1}'(s_{n-1}) + a_1(\beta_{n-1}) v_{n-1}(s_{n-1}) + Q(\beta_{n-1})] \Delta t, \quad s_0 = 0, \end{array} \right. \quad (30)$$

and $s_N(t), v_N(x, t)$ are defined in the same form as in (24) but s_n and v_n are produced by (30).

Arguing as in Lemma 1-4, we have

Lemma 5: *There exists a unique solution to Equation (30) and the corresponding functions $s_N(t)$ and $v_N(x, t)$ satisfy conditions (b)-(f) of Lemma 4. Moreover,*

$$0 \leq s_N'(t) = C_0 \leq 2 \frac{\kappa}{\rho L} \max\{\|v_0\|, q \beta_M / \alpha^2, q \|a_1\| + \|Q\|\}. \quad (31)$$

Now

$$\begin{aligned} s_N(T) &= s_N = \sum_{n=1}^N (s_n - s_{n-1}) \\ &= \frac{\kappa}{\rho L} \sum_{n=1}^N [-v_{n-1}'(s_{n-1}) + a_1(\beta_{n-1}) v_{n-1}(s_{n-1}) + Q(\beta_{n-1})] \Delta t \leq C_0 T. \end{aligned}$$

If T is sufficiently small, then $s_N < \text{const.} < \ell$ for all N . Let $S(t)$ and $\Phi(x, t)$ be the uniform limits of subsequence of $s_N(t)$ and $v_N(x, t)$, which for easier notation are still denoted by s_N and v_N , as N goes to infinity. Observe that

$$\frac{\rho L}{\kappa} s_N(T) = \sum_{n=1}^N (s_n - s_{n-1})$$

$$\begin{aligned}
 &= \sum_{n=1}^N [-v'_{n-1}(s_{n-1}) + a_1(\beta_{n-1})v_{n-1}(s_{n-1}) + Q(\beta_{n-1})]\Delta t \\
 &= \sum_{n=1}^N -v_{Nx}(s_N(t_{n-1}), t_{n-1})\Delta t + \sum_{n=1}^N a_1(\beta_{n-1})v_N(s_N(t_{n-1}), t_{n-1})\Delta t \\
 &\quad + \sum_{n=1}^N Q(\beta_{n-1})\Delta t. \\
 &= \sum_{n=1}^N [\Phi_x(s_N(t_{n-1}), t_{n-1}) - v_{Nx}(s_N(t_{n-1}), t_{n-1})]\Delta t \\
 &\quad + \sum_{n=1}^N [-\Phi_x[s_N(t_{n-1}), t_{n-1}) + \Phi_x(S(t_{n-1}), t_{n-1})]\Delta t \\
 &\quad + \sum_{n=1}^N a_1(\beta_{n-1})[v_N(s_N(t_{n-1}), t_{n-1}) - \Phi(s_N(t_{n-1}), t_{n-1})]\Delta t \\
 &\quad + \sum_{n=1}^N a_1(\beta_{n-1})[\Phi(s_N(t_{n-1}), t_{n-1}) - \Phi(S(t_{n-1}), t_{n-1})]\Delta t \\
 &\quad + \sum_{n=1}^N [-\Phi_x(s(t_{n-1}), t_{n-1}) + a_1(\beta_{n-1})\Phi(S(t_{n-1}), t_{n-1}) + Q(\beta_{n-1})]\Delta t.
 \end{aligned}$$

The last term of the above formula is a Riemann sum. From uniform convergence we have immediately that

$$S(T) = \frac{\kappa}{\rho L} \int_0^T [-\Phi_x(S(t), t) + a_1(\beta(t))\Phi(x, t) + Q(\beta(t))]dt.$$

Considering t as any $t \in (0, T]$ we also have

$$S(t) = \frac{\kappa}{\rho L} \int_0^t [-\Phi_x(S(t), t) + a_1(\beta(t))\Phi(x, t) + Q(\beta(t))]dt. \tag{32}$$

From the same arguments in (26), we also have

$$\left\{ \begin{array}{l} \Phi(S(t), t) \geq 0, \quad -\Phi'(S(t), t) + a_1(\beta(t))\Phi(x, t) + Q(\beta(t)) \geq 0, \\ \Phi(S(t), t)[- \Phi'(S(t), t) + a_1(\beta(t))\Phi(x, t) + Q(\beta(t))] = 0, \\ \alpha^2 \Phi_x(\ell, t) + \beta(t)\Phi(\ell, t) = q\beta(t). \end{array} \right. \tag{33}$$

From (32), $S(t)$ is a C^1 function on $[0, T]$. Using the same argument as in the proof of Theorem 1, we can prove for $S(t)$, $\Phi(x, t)$ is a solution of Equations (8)-(9). Finally, it is easy to see that

$$\|\Phi_{xx}\| \leq C,$$

where C depends only on the C^1 norm of β and $s(t)$, the C^2 norm of $v_0(x)$, and $\ell - s(T)$. Since $S'(t) \leq C_0$, where C_0 is defined in (31), C only depends on the C^1 of β , the C^2 norm of $v_0(x)$, and $\ell - s(T)$. We can extend this solution until $S(T) = \ell$ (e.g. [12]).

Theorem 2:

- (i) *There exists a solution $(s(t), u(x, t)), (x, t) \in [s(t), \ell] \times [0, T]$ to Equation (4) and either $T = \infty, s(t) < \ell$ for all $t \geq 0$; or $T < \infty$ and $s(T) = \ell$.*

This solution can be obtained as the uniform limit of the following Rothe approximation solutions $s_N(t)$ and $u_N(x, t)$:

$$\begin{cases} s_N(t) = \frac{1}{\Delta t}[(t - t_{n-1})s_n + (t_n - t)s_{n-1}], \text{ for } t \in [t_{n-1}, t_n), \\ u_N(x, t) = \frac{1}{\Delta t}[(t - t_{n-1})u_n(x) + (t_n - t)u_{n-1}(x)], \text{ for } t \in [t_{n-1}, t_n), \end{cases} \tag{34}$$

in which s_n and v_n are produced by

$$\begin{cases} s_n = s_{n-1} + \frac{\kappa}{\rho L}[u'_{n-1}(s_{n-1}) - a_1(\beta_{n-1})v_{n-1}(s_{n-1}) + a_2(\beta_{n-1})]\Delta t. \\ u'_n(x) = R_n(x)u_n(x) + z_n(x) \\ u_n(s_n) = \min\{u_m, \frac{a_2(\beta_n) + z_n(s_n)}{-R_n(s_n) + a_1(\beta_n)}\}, \end{cases} \tag{35}$$

where $R_n(x) < 0, z_n(x) \geq 0, \vartheta(x) \equiv \frac{\ell - s_{n-1}}{\ell - s_n}x - \frac{s_n - s_{n-1}}{\ell - s_n}\ell,$

$$\begin{cases} R'_n(x) = -R_n^2(x) - \frac{1}{\alpha^2}[\beta_n + 2\frac{s_n - s_{n-1}}{\Delta t} \frac{\ell - x}{\ell - s_n}]R_n(x) + \frac{1}{\alpha^2 \Delta t}, R_n(\ell) = -\beta_n/\alpha^2, \\ z'_n(x) = -[R_n(x) + \frac{1}{\alpha^2}(\beta_n + 2\frac{s_n - s_{n-1}}{\Delta t} \frac{\ell - x}{\ell - s_n})]z_n(x) - \frac{u_{n-1}(\vartheta(x))}{\alpha^2 \Delta t}, z_n(\ell) = q\beta_n/\alpha^2; \end{cases}$$

- (ii) $u_c \leq u(x, t) \leq u_m;$
- (iii) $u_x(x, t) \leq 0, \text{ if } u'_0(x) \leq 0.$

4. NUMERICAL SIMULATION

The following parameter is taken from a real material with a steel bar as the ablation material. The melting point is $1450^\circ C$, and helium is the coolant.

$$\begin{aligned} \ell &= 0.015m, c_\rho = 0.153kcal/kg^\circ C, \kappa = 0.0063kcal/^\circ C.s.m., \\ c_{\rho L} &= 1.24kcal/kg^\circ C, \rho = 7850kg/m^3, h_r = 3500kcal/kg, \end{aligned}$$

$$h_e = 1728 \text{kcal/kg}, q_0 = 5000 \text{kcal/s.m}^2, L = 65 \text{kcal/kg},$$

$$u_m = 1450^\circ\text{C}, u_c = 315^\circ\text{C}, u_0(x) = 516^\circ\text{C}.$$

The numerical results are calculated through solving (35) by a FORTRAN program. The results demonstrate that if coolant is not added, the steel bar is almost melted at 10 seconds, but in the case of inputting coolant $m = 2$ (the maximum coolant value in our example: $m = 4.38$), the temperature of the outer surface never attains the melting point: the coolant is very effective.

(1) $m = 0, T = 10, N = 30$

t_n	$s_N(t_n)$	$u_N(s_N(t_n), t_n)$
0.3333333	3.823523E-003	
0.6666667	3.823523E-003	237.8863
1.0000000	3.823523E-003	1172.275
1.3333330	4.159586E-003	1450.000
1.6666670	4.159586E-003	1405.850
2.0000000	5.066955E-003	1450.000
2.3333330	5.066955E-003	993.608
2.6666670	5.235254E-003	1450.000
3.0000000	5.714478E-003	1450.000
3.3333330	5.714478E-003	1322.453
3.6666660	6.508625E-003	1450.000
4.0000000	6.508625E-003	1077.579
4.3333330	6.868615E-003	1450.000
4.6666670	6.868615E-003	1412.066
5.0000000	7.886279E-003	1450.000
5.3333330	7.886279E-003	957.884
5.6666670	8.059592E-003	1450.000
6.0000000	8.614007E-003	1450.000
6.3333340	8.614007E-003	1277.720
6.6666670	9.440608E-003	1450.000
7.0000010	9.440608E-003	1090.880
7.3333340	9.941348E-003	1450.000
7.6666680	9.941348E-003	1349.251
8.0000010	1.097733E-002	1450.000
8.3333340	1.097733E-002	985.194
8.6666670	1.140899E-002	1450.000
9.0000000	1.144755E-002	1450.000
9.3333330	1.276071E-002	1450.000
9.6666660	1.276071E-002	953.719
9.9999990	1.355138E-002	1450.000

(2) $m = 2, T = 10, N = 30$

t_n	$s_N(t_n)$	$u_N(s_N(t_n), t_n)$
0.3333333	2.393934E-003	
0.6666667	2.393934E-003	241.9474
1.0000000	2.393934E-003	660.9301
1.3333330	2.393934E-003	805.6894
1.6666670	2.393934E-003	887.0065
2.0000000	2.393934E-003	942.0304
2.3333330	2.393934E-003	983.4641
2.6666670	2.393934E-003	1016.6440
3.0000000	2.393934E-003	1044.0590
3.3333330	2.393934E-003	1068.1590
3.6666660	2.393934E-003	1087.3070
4.0000000	2.393934E-003	1104.1690
4.3333330	2.393934E-003	1117.2270
4.6666670	2.393934E-003	1137.4850
5.0000000	2.393934E-003	1142.5890
5.3333330	2.393934E-003	1146.9740
5.6666670	2.393934E-003	1151.2580
6.0000000	2.393934E-003	1155.2860
6.3333340	2.393934E-003	1158.9660
6.6666670	2.393934E-003	1162.2390
7.0000010	2.393933E-003	1167.0030
7.3333340	2.393933E-003	1174.3570
7.6666680	2.393933E-003	1173.6960
8.0000010	2.393933E-003	1173.3640
8.3333340	2.393933E-003	1173.7020
8.6666670	2.393933E-003	1174.3910
9.0000000	2.393933E-003	1175.2630
9.3333330	2.393934E-003	1176.2150
9.6666660	2.393934E-003	1177.1690
9.9999990	2.393933E-003	1178.0520




Figure 2. Numerical simulation of an ablation steel bar

REFERENCES

- [1] Yang, X.S., Transpiration cooling control of thermal protection, *Acta Automatica Sinica*, 11 (1985), 345-350.
- [2] Yang, X.S., Transpiration cooling with control of water temperature field, *Acta Automatica Sinica*, 17 (1991), 385-394.
- [3] Fair, H.D., Electromagnetic earth-to-space launch, *IEEE Transactions on Magnetics*, 25 (1989), 9-16.
- [4] Palmer, M.R., Electromagnetic space launch: a re-evaluation in light of current technology and launch needs and feasibility of a near term demonstration, *IEEE Transactions on Magnetics*, 25 (1989), No. 1.
- [5] Jiang, L.S., Remarks on the Stefan-Signorini Problem, in *Free Boundary Problems: Vol. III*, Res. Notes in Math. 120, 1985, 13-19.
- [6] Friedman, A., and Jiang, L.S., A Stefan-Signorini Problem, *J. Diff. Eqs.*, 51 (1984), 213-231.
- [7] Friedman, A., *Partial Differential Equations of Parabolic Type*, Prentice-Hall, 1964.
- [8] Fasano, A., Meyer, G.H., and Primicerio, M., On a problem in the Polymer Industry: Theoretical and numerical investigations of swelling, *SIAM J. Math. Anal.*, 17 (1986), 945-960.
- [9] Meyer, G.H., One-dimensional parabolic free boundary problems, *SIAM*, 19 (1977), 17-34.
- [10] Protter, M.H. and Weinberger, H.F., *Maximum Principles in Differential Equations*, Springer-Verlag, 1984.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

