A COUNTEREXAMPLE TO A COMPACT EMBEDDING THEOREM FOR FUNCTIONS WITH VALUES IN A HILBERT SPACE

Accepted for publication to the
Proceedings of the American Mathematical Society
 Providence, RI USA

S. MIGÓRSKI

Jagellonian University, Institute for Information Sciences
ul. Nawojki 11, 30072 Cracow, Poland
E-mail: migorski@softlab.ii.uj.edu.pl
Fax: 4812 341865

Let V and H be two real separable Hilbert spaces with V densely and continuously embedded in H . Identifying H with its dual, we have a Gelfand triple (V, H, V') , where V' is the dual space to V . For a positive T , let $\mathcal{V} = L^2(0, T; V)$, $\mathcal{V}' = L^2(0, T; V')$ denote the spaces of the square summable functions defined on $(0, T)$ with values in V and V' , respectively. Let us define $\mathcal{W} = \{v \in \mathcal{V} : v' \in \mathcal{V}'\}$, where the time derivative $u' = \frac{du}{dt}$ is understood in the weak sense. Equipped with the standard norm $\|v\|_{\mathcal{W}} = \left(\|v\|_{\mathcal{V}}^2 + \|v'\|_{\mathcal{V}'}^2 \right)^{\frac{1}{2}}$, $\mathcal{W}$ becomes a Hilbert space. It is known (cf. [2]) that the embedding

$$\mathcal{W} \subset C(0, T; H) \tag{1}$$

is continuous, where the space of continuous functions from $[0, T]$ into H , is endowed with the supremum norm. E. Nagy proved the following theorem in [3].

Theorem: *If V and H are infinite-dimensional separable Hilbert spaces such that $V \subset H$ densely, continuously and compactly, then embedding (1) is also compact.*

The purpose of the paper is to deliver that this result of Nagy was recently exploited in several papers in connection with the study of evolution equations and inclusions (see, for instance, [1]).

References

- [1] Hu, S., Lakshmikantham, V. and Papageorgiou, N.S., On the solution set of nonlinear evolution inclusions, *Dynamic Systems Appl.* **1** (1992), 71-82.
- [2] Lions, J.L. and Magenes, E., *Problèmes aux limites non homogènes et applications*, vol. 1, Dunod, Paris 1968.
- [3] Nagy, E.V., A theorem on compact embedding for functions with values in an infinite dimensional Hilbert space, *Ann. Univ. Sci. Budapest Eötvös Sect. Math.* **22-23** (1979-80), 243-245.