AN ITERATIVE ALGORITHM ON FIXED POINTS OF RELAXED LIPSCHITZ OPERATORS

RAM U. VERMA

*International Publications, 12046 Coed Drive
Orlando, Florida 32826 USA*

and

*Istituto per la Ricerca di Base
I-86075 Monteroduni (IS), Molise, Italy*

(Received February, 1996; Revised November, 1996)

Fixed points of Lipschitzian relaxed Lipschitz operators based on a generalized iterative algorithm are approximated.

Key words: Iterative Algorithm, Fixed Point, Relaxed Lipschitz Operator.

AMS subject classifications: 47H10.

1. Introduction

Recently, Wittman [6, Theorem 2], using an iterative procedure

$$x_n = (1 - a_n)x_0 + a_nTx_{n-1} \text{ for } n \geq 1, \quad (1)$$

approximated fixed points of nonexpansive mappings $T:K \rightarrow K$ from a nonempty closed convex subset K of a real Hilbert space H into itself, where x_0 is an element of K and $\{a_n\}$ is an increasing sequence in $[0, 1)$ such that

$$\lim_{n \rightarrow \infty} a_n = 1 \text{ and } \sum_{n=1}^{\infty} (1 - a_n) = \infty. \quad (2)$$

This result refines a number of results including [1].

Here our aim is to approximate the fixed points of Lipschitzian relaxed Lipschitz operators in a Hilbert space setting. As such, the iterative algorithm (1) is not suitable for our purpose, so we apply a modified iterative algorithm which reduces to (1).

Let H be a Hilbert space and $\langle u, v \rangle$ and $\|u\|$ denote, respectively, the inner product and norm on H for u, v in H .

An operator $T: H \rightarrow H$ is said to be *relaxed Lipschitz* if, for all u, v in H , there exists a constant $r > 0$ such that

$$\langle Tu - Tv, u - v \rangle \leq -r \|u - v\|^2. \quad (3)$$

The operator T is called *Lipschitz continuous* (or Lipschitzian) if there exists a constant $s > 0$ such that

$$\|Tu - Tv\| \leq s \|u - v\| \text{ for all } u, v \text{ in } H. \quad (4)$$

Next, we consider the main result on the approximation of the fixed points of Lipschitzian relaxed Lipschitz operators using a modified iterative algorithm which contains a number of iterative schemes including those considered by the author [4, 5] as special cases.

2. The Main Result

Theorem 1: Let H be a real Hilbert space and K be a nonempty closed convex subset of H . Let $T: K \rightarrow K$ be a relaxed Lipschitz and Lipschitz continuous operator on K . Let $r \geq 0$ and $s \geq 1$ be constants for relaxed Lipschitzity and Lipschitz continuity of T , respectively. Let $F = \{x \text{ in } K: Tx = x\}$ be nonempty, and let $\{a_n\}$ be a sequence in $[0, 1]$ such that

$$\sum_{n=0}^{\infty} a_n = \infty \text{ for all } n \geq 0. \quad (5)$$

Then for any x_0 in K the sequence $\{x_n\}$ defined by

$$x_{n+1} = (1 - a_n)x_n + a_n[(1 - t)x_n + tTx_n] \text{ for } n \geq 0, \quad (6)$$

$0 < k = ((1 - t)^2 - 2t(1 - t)r + t^2s^2)^{1/2} < 1$ for all t such that $0 < t < 2(1 + r)/(1 + 2r + s^2)$ and $r \leq s$, converges to an element of F .

For $\{a_n\} = 1$, Theorem 1 reduces to:

Corollary 1: Let $T: K \rightarrow K$ be relaxed Lipschitz and Lipschitz continuous. Let $F = \{x \text{ in } K: Tx = x\}$ be a nonempty set. Then, for x_0 in K , the sequence $\{x_n\}$ generated by an iterative algorithm

$$x_{n+1} = (1 - t)x_n + tTx_n \quad (7)$$

for $0 < t < 2(1 + r)/(1 + 2r + s^2)$ converges to a unique fixed point of T .

Proof of Theorem 1: For an element z in F , we have

$$\begin{aligned} \|x_{n+1} - z\| &= \|(1 - a_n)x_n + a_n[(1 - t)x_n + tTx_n] - z\| \\ &\leq (1 - a_n)\|x_n - z\| + a_n\|(1 - t)(x_n - z) + t(Tx_n - Tz)\|. \end{aligned}$$

Using the relaxed Lipschitzity and Lipschitz continuity of T , we find that

$$\begin{aligned} &\|t(Tx_n - Tz) + (1 - t)(x_n - z)\|^2 \\ &= (1 - t)^2\|x_n - z\|^2 + 2t(1 - t)\langle Tx_n - z, x_n - z \rangle + t^2\|Tx_n - z\|^2 \\ &\leq (1 - t)^2\|x_n - z\|^2 - 2t(1 - t)r\|x_n - z\|^2 + t^2s^2\|x_n - z\|^2 \end{aligned}$$

$$= ((1-t)^2 - 2t(1-t)r + t^2s^2) \|x_n - z\|^2.$$

It follows that

$$\begin{aligned} \|x_{n+1} - z\| &\leq (1 - a_n + a_n((1-t)^2 - 2t(1-t)r + t^2s^2)^{1/2}) \|x_n - z\| \\ &= (1 - (1-k)a_n) \|x_n - z\| \\ &\leq \prod_{j=0}^n (1 - (1-k)a_j) \|x_0 - z\|, \end{aligned}$$

where $0 < k = ((1-t)^2 - 2t(1-t)r + t^2s^2)^{1/2} < 1$ for all t such that $0 < t < 2(1+r)/(1+2r+s^2)$ and $r \leq s$.

Since $\sum_{j=0}^{\infty} a_j$ diverges and $k < 1$, $\lim_{n \rightarrow \infty} \sum_{j=0}^n (1 - (1-k)a_j) = 0$ and, as a result,

$\{x_n\}$ converges strongly to z . This completes the proof.

References

- [1] Halpern, B., Fixed points of nonexpanding maps, *Bull. Amer. Math. Soc.* **73** (1967), 957-961.
- [2] Reich, S., Approximating fixed points of nonexpansive maps, *PanAmerican Math J.* **4:2** (1994), 23-28.
- [3] Verma, R.U., Iterative algorithms for approximating fixed points of strongly monotone operators, *Boletín Acad. Cien. Fis. Mat. Natur.* (to appear).
- [4] Verma, R.U., A fixed point theorem involving Lipschitzian generalized pseudo-contractions, *Proc. Royal Irish Acad.* (to appear).
- [5] Verma, R.U., An iterative procedure for approximating fixed points of relaxed monotone operators, *Numer. Functional Anal. Optimiz.* **17** (1996), 1045-1051.
- [6] Wittmann, R., Approximation of fixed points of nonexpansive mappings, *Arch. Math.* **58** (1992), 486-491.