CONVERGENCE OF AN ITERATION LEADING TO A SOLUTION OF A RANDOM OPERATOR EQUATION

BINAYAK S. CHOUDHURY

B.E. College (Deemed University)

Department of Mathematics

Howrah, West Bengal, India 711103

M. RAY

Rashmoni Balika Vidyalaya

Baruipur, West Bengal, India

(Received April, 1996; Revised June, 1998)

In the present paper, we define a random iteration scheme and consider its convergence to a solution of a random operator equation. There is also a related fixed point result.

Key words: Random Equation, Random Operator, Random Iteration, Fixed Point, Hilbert Space.

AMS subject classifications: 47H40, 47H10.

1. Introduction

In recent years, the study of different types of random equations have attracted much attention, some of which may be noted in [1, 5, 6] and [7]. In this paper, we discuss a random operator equation involving two operators in the context of Hilbert spaces. We have also a random fixed point result as a corollary. We also demonstrate our result for the corresponding deterministic case by an example.

Throughout this paper, (Ω, Σ) denotes a measurable space and H stands for a separable Hilbert space.

A function $f: \Omega \rightarrow H$ is said to be *measurable* if $f^{-1}(B) \in \Sigma$ for every Borel subset B of H .

A function $F: \Omega \times H \rightarrow H$ is said to be *H-continuous*, if $F(t, \cdot): H \rightarrow H$ is continuous for all $t \in \Omega$.

A function $F: \Omega \times H \rightarrow H$ is said to be a *random operator*, if $F(\cdot, x): \Omega \rightarrow H$ is measurable for every $x \in H$.

A measurable function $g: \Omega \rightarrow H$ is said to be a *random fixed point* of the random operator $F: \Omega \times H \rightarrow H$, if $F(t, g(t)) = g(t)$ for all $t \in \Omega$.

A measurable function $g: \Omega \rightarrow H$ is said to be a *solution of the random operator equation* $S(t, x(t)) = T(t, x(t))$, where $S, T: \Omega \times H \rightarrow H$ are random operators, if $S(t, g(t)) = T(t, g(t))$ for all $t \in \Omega$.

The following result was established in [4]. We present this result as a lemma.

Lemma 1.1: *Let H be a Hilbert space. Then for any $x, y, z \in H$ and any real λ , the following equality holds:*

$$\begin{aligned} & \| (1 - \lambda)x + \lambda y - z \|^2 \\ &= (1 - \lambda) \| x - z \|^2 + \lambda \| y - z \|^2 - \lambda(1 - \lambda) \| x - y \|^2. \end{aligned} \quad (1.1)$$

We define the random iteration scheme as follows:

Definition 1.2: *Random iteration scheme.* Let $S, T: \Omega \times H \rightarrow H$ be two random operators defined on a Hilbert space H . Let $g_0: \Omega \rightarrow H$ be any measurable function. Define the following sequence of functions $\{g_n\}$

$$g_{n+1}(t) = (1 - \alpha_n)g_n(t) + \alpha_n h_n(t), \quad (1.2)$$

where

$$h_n(t) = (1 - \beta_n)S(t, g_n(t)) + \beta_n T(t, g_n(t)), \quad (1.3)$$

$$0 < \alpha_n, \beta_n < 1 \text{ for all } n = 0, 1, 2, \dots, \quad (1.4)$$

$$\overline{\lim}_{n \rightarrow \infty} \beta_n = M < 1, \quad (1.5)$$

and

$$\sum_{n=1}^{\infty} \alpha_n \beta_n = \infty. \quad (1.6)$$

The construction of the iteration scheme is based on the same idea as that of Ishikawa's random iteration scheme [2]. But the present iteration is not a modification or generalization of that iteration.

A function $T: H \rightarrow H$ is said to satisfy *Tricomi's condition* if

$$Tp = p \text{ implies } \|Tx - p\| \leq \|x - p\|.$$

We define generalized Tricomi's condition for two operators in the following way.

Definition 1.3: *Generalized Tricomi's Condition.* Two functions $S, T: H \rightarrow H$ are said to satisfy *generalized Tricomi's condition* if

$$Sp = Tp \text{ implies } \|Sx - p\| \leq \|x - p\| \quad (1.7)$$

and

$$\|Tx - p\| \leq \|x - p\|. \quad (1.8)$$

2. Main Results

Theorem 2.1: *Let $S, T: \Omega \times H \rightarrow H$, where H is a separable Hilbert space, be two random operators such that*

- (a) S and T are H -continuous, and
 (b) there exists $f: \Omega \rightarrow H$ (not necessarily measurable) such that

$$(1 - \lambda) \|S(t, x) - f(t)\|^2 + \lambda \|T(t, x) - f(t)\|^2 \leq \|x - f(t)\|^2 \quad (2.1)$$

for all $t \in \Omega, x \in H$ and $0 < \lambda < 1$.

Then the random iteration scheme (Definition 1.2), if convergent, converges to a solution of the random operator equation

$$S(t, x(t)) = T(t, x(t)). \quad (2.2)$$

Proof: For any $t \in \Omega$,

$$\begin{aligned} \|g_{n+1}(t) - f(t)\|^2 &= \|(1 - \alpha_n)g_n(t) + \alpha_n h_n(t) - f(t)\|^2 \quad (\text{by (1.2)}) \\ &= (1 - \alpha_n) \|g_n(t) - f(t)\|^2 + \alpha_n \|h_n(t) - f(t)\|^2 \\ &\quad - \alpha_n(1 - \alpha_n) \|g_n(t) - h_n(t)\|^2 \quad (\text{by (1.1)}) \\ &\leq (1 - \alpha_n) \|g_n(t) - f(t)\|^2 \\ &\quad + \alpha_n \|(1 - \beta_n)S(t, g_n(t)) + \beta_n T(t, g_n(t)) - f(t)\|^2 \quad (\text{by (1.4) and (1.3)}) \\ &= (1 - \alpha_n) \|g_n(t) - f(t)\|^2 + \alpha_n \{(1 - \beta_n) \|S(t, g_n(t)) - f(t)\|^2 \\ &\quad + \beta_n \|T(t, g_n(t)) - f(t)\|^2 - \beta_n(1 - \beta_n) \|S(t, g_n(t)) - T(t, g_n(t))\|^2\} \quad (\text{by (1.1)}) \end{aligned}$$

or,

$$\begin{aligned} &\alpha_n \beta_n (1 - \beta_n) \|S(t, g_n(t)) - T(t, g_n(t))\|^2 \\ &\leq (1 - \alpha_n) \|g_n(t) - f(t)\|^2 - \|g_{n+1}(t) - f(t)\|^2 \\ &\quad + \alpha_n \{(1 - \beta_n) \|S(t, g_n(t)) - f(t)\|^2 + \beta_n \|T(t, g_n(t)) - f(t)\|^2\} \\ &\quad \text{for all } t \in \Omega \end{aligned} \quad (2.3)$$

or,

$$\begin{aligned} \alpha_n \beta_n (1 - \beta_n) \|S(t, g_n(t)) - T(t, g_n(t))\|^2 &\leq (1 - \alpha_n) \|g_n(t) - f(t)\|^2 \\ &\quad - \|g_{n+1}(t) - f(t)\|^2 + \alpha_n \|g_n(t) - f(t)\|^2 \quad \text{for all } t \in \Omega \quad (\text{by (2.1)}) \end{aligned}$$

or,

$$\begin{aligned} \alpha_n \beta_n (1 - \beta_n) \|S(t, g_n(t)) - T(t, g_n(t))\|^2 &\leq \|g_n(t) - f(t)\|^2 \\ &\quad - \|g_{n+1}(t) - f(t)\|^2 \quad \text{for all } t \in \Omega. \end{aligned} \quad (2.4)$$

Summing up the inequalities in (2.4) over n , we obtain for all $t \in \Omega$,

$$\sum_{n=0}^{\infty} \alpha_n \beta_n (1 - \beta_n) \|S(t, g_n(t)) - T(g, g_n(t))\|^2 \leq \|g_0(t) - f(t)\|^2 < \infty. \quad (2.5)$$

Let $M < M' < 1$. Then, by (1.5), there exists a positive integer m_0 such that $\beta_m < M'$, that is $1 - \beta_m > 1 - M'$ for all $m > m_0$.

This shows that

$$\sum_{m=m_0}^{\infty} \alpha_m \beta_m (1 - \beta_m) \geq (1 - M') \sum_{m=m_0}^{\infty} \alpha_m \beta_m = \infty. \quad (2.6)$$

(2.5) and (2.6) imply that, for all $t \in \Omega$,

$$\lim_{n \rightarrow \infty} \|S(t, g_n(t)) - T(t, g_n(t))\|^2 = 0. \quad (2.7)$$

Let

$$g_n(t) \rightarrow g(t) \text{ as } n \rightarrow \infty. \quad (2.8)$$

Since g_0 is measurable and H is separable, according to Himmelberg [3], g_n 's are measurable and, therefore, $g: \Omega \rightarrow H$ is measurable.

Again, S and T are H -continuous, which shows that $\lim_{n \rightarrow \infty} S(t, g_n(t)) = S(t, g(t))$ and

$$\lim_{n \rightarrow \infty} T(t, g_n(t)) = T(t, g(t)) \text{ for all } t \in \Omega.$$

By (2.7) and (2.8), we have for all $t \in \Omega$,

$$S(t, g(t)) = T(t, g(t)), \text{ where } g: \Omega \rightarrow H \text{ is a measurable function.} \quad (2.9)$$

This shows that the random iteration scheme if convergent, converges to a solution of (2.2). $\square$

Corollary 2.2: *Let H be a separable Hilbert space and $S, T: \Omega \times H \rightarrow H$ be two random operators such that*

- (a) S, T are H -continuous, and
- (b) *there exists $f: \Omega \rightarrow H$ (not necessarily measurable) such that*

$$\|S(t, x) - f(t)\| \leq \|x - f(t)\| \quad (2.10)$$

and

$$\|T(t, x) - f(t)\| \leq \|x - f(t)\|. \quad (2.11)$$

Then the random iteration scheme if convergent, converges to a solution of $S(t, x(t)) = T(t, x(t))$.

Proof: It is easily seen that (2.10) and (2.11) imply (2.1). The corollary then follows by Theorem 2.1. $\square$

Setting S as the identify random operator, that is, $S(t, x) = x$ for all $t \in \Omega$ and

$x \in H$, we obtain the following fixed point result as a corollary.

Corollary 2.3: *Let H be a separable Hilbert space and T be a random operator which is H -continuous. Assume that there exists $f: \Omega \rightarrow H$ (not necessarily measurable) such that for all $t \in \Omega$*

$$\|T(t, x) - f(t)\| \leq \|x - f(t)\|. \quad (2.12)$$

Then the sequence of functions $\{g_n\}$, where $g_0: \Omega \rightarrow H$, is measurable and

$$g_{n+1}(t) = (1 - \alpha_n)g_n(t) + \alpha_n((1 - \beta_n)g_n(t) + \beta_n T(t, g_n(t))), \quad n = 0, 1, 2, \dots \quad (2.13)$$

for all $t \in \Omega$, where $0 < \alpha_n, \beta_n < 1$, $\overline{\lim}_{n \rightarrow \infty} \beta_n < 1$, and $\sum_{n=1}^{\infty} \alpha_n \beta_n = \infty$ if convergent, converges to a random fixed point of T .

Corollary 2.4: *Let $S, T: H \rightarrow H$ be two operators such that the following holds: there exists a $z \in H$ such that*

$$(1 - \lambda) \|Sx - z\|^2 + \lambda \|Tx - z\|^2 \leq \|x - z\|^2, \quad (2.14)$$

for all $x \in H$ and $0 < \lambda < 1$.

Then the sequence $\{x_n\}$, obtained by the iteration

$$x_0 \in H, \quad (2.15)$$

$$x_{n+1} = (1 - \alpha_n)x_n + \alpha_n((1 - \beta_n)Sx_n + \beta_n Tx_n), \quad (2.16)$$

where

$$0 < \alpha_n, \beta_n < 1 \quad (2.17)$$

$$\overline{\lim}_{n \rightarrow \infty} \beta_n < 1 \quad (2.18)$$

and

$$\sum_{n=1}^{\infty} \alpha_n \beta_n = \infty, \quad (2.19)$$

if convergent, converges to a solution of $Sx = Tx$.

The proof trivially follows from Theorem 2.1. It may be noted that the separability of H was required to ensure that g_n 's are measurable. In the statement of the corollary, H need not be separable.

Theorem 2.5: *Let C be a convex and compact subset of a separable Hilbert space H and $S, T: \Omega \times H \rightarrow C$ be two random operators such that the following conditions are satisfied:*

- (a) S, T are H -continuous,
- (b) there exists $f: \Omega \rightarrow H$ (not necessarily measurable) such that

$$(1 - \lambda) \|S(t, x) - f(t)\|^2 + \lambda \|T(t, x) - f(t)\|^2 \leq \|x - f(t)\|^2 \quad (2.20)$$

for all $t \in \Omega$, $x \in H$ and $0 < \lambda < 1$, and

(c) $S(t, \cdot), T(t, \cdot): C \rightarrow C$ satisfy Generalized Tricomi's condition for all $t \in \Omega$. Then for any measurable function $g_0: \Omega \rightarrow C$, the sequence of functions $\{g_n\}$ constructed by the random iteration scheme ((1.2)-(1.6)) actually converges to a solution of the random operator equation $S(t, x(t)) = T(t, x(t))$.

Proof: By the construction of $\{g_n\}$ it is seen that g_n 's are measurable functions from Ω to C for all $n = 0, 1, 2, \dots$. Proceeding exactly in the same way as in Theorem 2.1, we have as in (2.7) that

$$\lim_{n \rightarrow \infty} \|S(t, g_n(t)) - T(t, g_n(t))\|^2 = 0.$$

Therefore, for a fixed $t \in \Omega$, there exists a subsequence

$$\{g_{n_i}(t)\} \subset \{g_n(t)\} \text{ such that } \lim_{i \rightarrow \infty} \|S(t, g_{n_i}(t)) - T(t, g_{n_i}(t))\| = 0. \quad (2.21)$$

Again, C is compact, therefore, there exists $\{g_{n_{i_k}}(t)\} \subset \{g_{n_i}(t)\}$ such that $\{g_{n_{i_k}}(t)\}$ is convergent.

Let

$$\lim_{k \rightarrow \infty} g_{n_{i_k}}(t) = g(t) \text{ for } t \in \Omega. \quad (2.22)$$

Since S and T are H -continuous random operators, from (2.21), we have for any $t \in \Omega$,

$$S(t, g(t)) = T(t, g(t)). \quad (2.23)$$

For any $t \in \Omega$,

$$\begin{aligned} & \|g_{n+1}(t) - g(t)\|^2 = \|(1 - \alpha_n)g_n(t) + \alpha_n h_n(t) - g(t)\|^2 \\ & = (1 - \alpha_n) \|g_n(t) - g(t)\|^2 + \alpha_n \|h_n(t) - g(t)\|^2 \\ & \quad - (1 - \alpha_n)\alpha_n \|g_n(t) - h_n(t)\|^2 \\ & \leq (1 - \alpha_n) \|g_n(t) - g(t)\|^2 + \alpha_n \|(1 - \beta_n)S(t, g_n(t)) + \beta_n T(t, g_n(t)) - g(t)\|^2 \\ & \quad \text{(by (1.3) and (1.4))} \\ & = (1 - \alpha_n) \|g_n(t) - g(t)\|^2 + \alpha_n \{(1 - \beta_n) \|S(t, g_n(t)) - g(t)\|^2 \\ & \quad + \beta_n \|T(t, g_n(t)) - g(t)\|^2\} - \alpha_n \beta_n (1 - \beta_n) \|S(t, g_n(t)) - T(t, g_n(t))\|^2 \\ & \quad \text{(by (1.1))} \\ & \leq (1 - \alpha_n) \|g_n(t) - g(t)\|^2 + \alpha_n \|g_n(t) - g(t)\|^2 \\ & \quad \text{(by (2.23) and Generalized Tricomi's condition)} \end{aligned}$$

or

$$\|g_{n+1}(t) - g(t)\| \leq \|g_n(t) - g(t)\|. \quad (2.24)$$

(2.22) and (2.24) together imply that

$$g_n \rightarrow g \text{ as } n \rightarrow \infty. \quad (2.25)$$

Since H is separable, g_n 's are measurable [3], and, hence g is also measurable. From (2.23), g is a random solution of $S(t, x(t)) = T(t, x(t))$. This completes the proof. $\square$

We have the following obvious corollary.

Corollary 2.6: *Let $S, T: H \rightarrow C$, where C is a compact and convex subset of a Hilbert space H are such that the following are satisfied:*

- (a) S, T are continuous,
- (b) there exists $z \in H$ such that

$$(1 - \lambda) \|Sx - z\|^2 + \lambda \|Tx - z\|^2 \leq \|x - z\|^2 \quad (2.26)$$

for all $x \in H$ and $0 < \lambda < 1$, and

- (c) S, T satisfy Generalized Tricomi's condition.

Then the sequence, defined as $x_0 \in C$,

$$x_{n+1} = (1 - \alpha_n)x_n + \alpha_n((1 - \beta_n)Sx_n + \beta_nTx_n), \quad n = 0, 1, 2, \dots, \quad (2.27)$$

where

$$0 < \alpha_n, \beta_n < 1, \quad (2.28)$$

$$\overline{\lim}_{n \rightarrow \infty} \beta_n < 1, \quad (2.29)$$

and

$$\sum_{n=1}^{\infty} \alpha_n \beta_n = \infty, \quad (2.30)$$

converges to a solution of the equation $Sx = Tx$.

Example: Let $C = [0, 1]$, $S, T: R \rightarrow [0, 1]$ be defined as

$$\begin{aligned} Sx &= x^2/2 \text{ if } x \in [0, 1] \\ &= 1/2 \text{ if } x > 1 \\ &= 0 \text{ if } x < 0 \end{aligned}$$

and

$$\begin{aligned} Tx &= x^2/4 \text{ if } x \in [0, 1] \\ &= 1/4 \text{ if } x > 1 \\ &= 0 \text{ if } x < 1. \end{aligned}$$

With the choice of $z = 0$, the conditions of Corollary 2.6 are seen to be satisfied. Thus Corollary 2.6 applies to this example.

References

- [1] Bharucha-Reid, A.T., *Random Integral Equations*, Academic Press, New York 1972.
- [2] Choudhury, B.S., Convergence of a random iteration scheme to a random fixed point, *J. Appl. Math. Stoch. Anal.* **8** (1995), 139-142.
- [3] Himmelberg, C.J., Measurable relations, *Fund. Math.* **LXXXVII** (1975), 53-71.
- [4] Ishikawa, S., Fixed points by a new iteration method, *Proc. Amer. Math. Soc.* **44** (1974), 147-150.
- [5] Itoh, S., Nonlinear random equations with monotone operators in Banach spaces, *Math. Ann.* **236** (1978), 133-146.
- [6] Karamolegos, A. and Kravvaritis, D., Nonlinear random operator equations and inequalities in Banach spaces, *Intern. J. Math. and Math. Sci.* **15** (1992), 111-118.
- [7] Kravvaritis, D. and Papageorgiou, N.S., Existence and solutions for nonlinear random operator equations in Banach spaces, *J. Math. Anal. Appl.* **141** (1989), 235-247.