GENERALIZED FLOW INVARIANCE FOR DIFFERENTIAL INCLUSIONS

T. GNANA BHASKAR AND V. LAKSHMIKANTHAM

Received 6 May 2004; Revised 12 May 2005; Accepted 12 May 2005

We introduce a generalized notion of invariance for differential inclusions, using a proximal aiming condition in terms of proximal normals. A set of sufficient conditions for the weak and strong invariance in the generalized sense are presented.

Copyright © 2006 T. G. Bhaskar and V. Lakshmikantham. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

The existence of solutions and flow invariance for differential inclusions are considered in [1] by using a generalized concept of solutions, namely, the Euler solutions of differential equations, without any continuity assumptions. This is done by utilizing a proximal aiming condition in terms of proximal normals. In a recent paper [2], we generalized the concept of proximal normal in the spirit of [3], and then, employing a generalized proximal aiming condition, we proved the existence and flow invariance results for solutions of differential inclusions.

Here in this paper, we consider a generalized notion of invariance, retaining the original notion of proximal normals as in [1], and study the corresponding results for differential inclusions. This generalized notion of flow invariance is useful in studying the solution sets of fuzzy differential equations, which will be considered in a separate paper.

2. Preliminaries

Consider the Cauchy problem

$$x'(t) = f(t, x(t)), \quad x(t_0) = x_0, \quad t_0 \geq 0, \quad (2.1)$$

where $f : [t_0, T] \times \mathbb{R}^n \rightarrow \mathbb{R}^n$ is any function.

2 Generalized flow invariance for differential inclusions

Let $\pi = \{t_0, t_1, \dots, t_N = T\}$ be a partition of $[t_0, T]$. On the interval $[t_0, t_1]$, we consider the differential equation with constant right-hand side

$$x'(t) = f(t_0, x_0), \quad x(t_0) = x_0, \quad (2.2)$$

which has a unique solution, $x(t)$ on $[t_0, t_1]$. Let $x_1 = x(t_1)$. Next, consider, on the interval $[t_1, t_2]$, the IVP

$$x'(t) = f(t_1, x_1), \quad x(t_1) = x_1. \quad (2.3)$$

We take $x_2 = x(t_2) = x(t_2, t_1, x_1)$ as the next node and proceeding in this manner until we get an arc $x_\pi = x_\pi(t)$ defined on all of $[t_0, T]$. The notation x_π is employed to emphasize the role played by the particular partition π in defining x_π which is the *Euler Polygonal arc* corresponding to the partition π . The diameter μ_π of the partition π is given by

$$\mu_\pi := \max \{t_i - t_{i-1} : 1 \leq i \leq N\}. \quad (2.4)$$

By an Euler solution to the IVP (2.1), we mean any arc $x(t)$ which is the uniform limit of the Euler polygonal arcs x_{π_j} , corresponding to some sequence of partitions π_j such that the diameters $\mu_{\pi_j} \rightarrow 0$ as $j \rightarrow \infty$. Clearly, this Euler arc satisfies the initial condition $x(t_0) = x_0$ and the corresponding number N_j of the partition points in π_j tends to infinity.

The following theorem, concerning the existence of Euler solutions for (2.1), is proved in [2].

THEOREM 2.1. *Assume that*

- (1) $f : [t_0, T] \times \mathbb{R}^n \rightarrow \mathbb{R}$ and $\|f(t, x)\| \leq g(t, \|x\|)$, $(t, x) \in [t_0, T] \times \mathbb{R}^n$, where $g : [t_0, T] \times \mathbb{R}_+ \rightarrow \mathbb{R}_+$ is a continuous function, nondecreasing in (t, u) ;
- (2) the maximal solution $r(t) = r(t, t_0, u_0)$ of the scalar differential equation

$$u' = g(t, u), \quad u(t_0) = u_0 \geq 0, \quad (2.5)$$

exists on $[t_0, T]$.

Then, there exists an Euler solution $x(t) = x(t, t_0, x_0)$ of the IVP (2.1) on $[t_0, T]$ which satisfies a Lipschitz condition and any Euler solution of (2.1) has an estimate

$$\|x(t) - x_0\| \leq r(t, t_0, \|x_0\|) - \|x_0\|, \quad t \in [t_0, T]. \quad (2.6)$$

Remark 2.2. We can extend the notion of Euler solution of (2.1) on the interval $[t_0, T]$ to $[t_0, \infty)$ provided we define f and g on $[t_0, \infty)$ instead of $[t_0, T]$, assume that the maximal solution on $r(t)$ exists on $[t_0, \infty)$, and show that Euler solution exists on every $[t_0, T]$, $T \in (t_0, \infty)$.

3. Generalized flow invariance

Let $S(t)$, $t \in [0, \infty)$ be a family of nonempty closed subsets of $\mathbb{R}^n$. Let $x \in \mathbb{R}^n$ be such that $(t, x) \notin \{(t, s) : s \in S(t)\}$, for all $t \geq 0$. Suppose that, for $t \geq 0$, there exists an $s_t \in S(t)$

such that

$$\|x - s_t\| = \|(t, x) - (t, s_t)\| = \inf \{\|x - \tilde{s}\| : \tilde{s} \in S(t)\}. \quad (3.1)$$

The set of all such $s_t \in S(t)$, for each $t \geq 0$, is denoted by $\text{proj}_{S(t)}(x)$. The vector $(t, x - s_t)$ determines a proximal normal direction to $(t, S(t))$ at (t, s_t) . We call any vector η_t of the form $(t, k(x - s_t))$, for any $k \geq 0$, a *proximal normal* (or *P-normal*) to $S(t)$ at s_t , at height t . The set of all η_t obtained in this manner is called a proximal normal cone to $S(t)$ at s_t , at a height t and is denoted by $N_{S(t)}^P(s_t)$. If $s_t \in S(t)$ such that $s_t \notin \text{proj}_{S(t)}(x)$ for all $(t, x) \notin \{(t, s) : s \in S(t)\}$, then we set $N_{S(t)}^P(s_t) = \{0\}$. If $s_t \notin S(t)$, then $N_{S(t)}^P$ is not defined.

Definition 3.1 (generalized flow invariance). The system $\{(S(t), f) : t \geq t_0\}$ is said to be weakly invariant if for all $x_0 \in S(t_0)$, there exists an Euler solution $x(t)$ of (2.1) on $[t_0, \infty)$ such that $x(t_0) = x_0$ and $x(t) \in S(t)$, $t > t_0$.

Note that this implies $(t, x(t)) \in (t, S(t))$, $t \geq t_0$. Also, if $S(t) = S(t_0)$, for all $t \geq t_0$, then the above notion of weak invariance coincides with the one given in [1].

Throughout the rest of the paper, we make the following assumption.

ASSUMPTION 3.2. For all $t > \tau$, $t, \tau \in [t_0, \infty)$ and $z \in \mathbb{R}^n$,

$$d_{S(t)}^2(z) \leq d_{S(\tau)}^2(z) + (t - \tau)^2. \quad (3.2)$$

We can now prove the following result which provides sufficient conditions in terms of the generalized proximal normal for weak invariance of $\{(S(t), f) : t \geq 0\}$.

THEOREM 3.3. Let f and g satisfy the assumptions of Theorem 2.1 on $[t_0, \infty)$ and let $x(t)$ be an Euler solution on $[t_0, \infty)$ of (2.1). Suppose that $x(t)$ lies in an open set $\Omega \subset \mathbb{R}^n$. Assume that for every $(t, z) \in [t_0, \infty) \times \omega$, there exists an $s_t \in \text{proj}_{S(t)}(z)$ such that

$$2\langle f(t, z), (z - s_t) \rangle \leq q(t, d_{S(t)}^2(z)), \quad (3.3)$$

where $q \in C([t_0, \infty) \times \mathbb{R}_+, \mathbb{R})$. Suppose also that the maximal solution $r(t) = r(t, t_0, u_0)$ of the scalar differential equation $u' = q(t, u)$, $u(t_0) = u_0 \geq 0$ exists on $[t_0, \infty)$. Then,

$$d_{S(t)}(x(t)) \leq r(t, t_0, d_{S(t_0)}^2(x_0)). \quad (3.4)$$

If, in addition, $r(t, t_0, 0) \equiv 0$, then $(S(t), f)$, $t \geq t_0$, is weakly invariant.

Proof. Let $x_\pi(t)$ be one polygonal arc in the sequence, converging uniformly to x as per the definition of Euler solution of (2.1). We denote, as before, its nodes at t_i by x_i , $i = 0, 1, \dots, N$, and hence $x(t_0) = x_0$. Let $x_\pi(t)$ be in Ω for all $t_0 \leq t \leq T$, where $T \in (t_0, \infty)$. Accordingly, there exists for each i an element $s_{t_i} \in \text{proj}_{S(t_i)}(x_i)$ such that

$$2\langle f(t_i, x_i), x_i - s_{t_i} \rangle \leq q(t_i, \|x_i - s_{t_i}\|^2). \quad (3.5)$$

4 Generalized flow invariance for differential inclusions

As in Theorem 2.1, letting $\|x'_\pi\| \leq k$, we find

$$\begin{aligned}
 d_{S(t_1)}^2(x_1) &\leq d_{S(t_0)}^2(x_1) + (t_1 - t_0)^2 \\
 &\leq \|x_1 - s_{t_0}\|^2 + (t_1 - t_0)^2 \\
 &\leq (t_1 - t_0)^2 + \|x_1 - x_0\|^2 + \|x_0 - s_{t_0}\|^2 + 2\langle x_1 - x_0, x_0 - s_{t_0} \rangle \\
 &\leq (k^2 + 1)(t_1 - t_0)^2 + d_{S(t_0)}^2(x_0) + 2 \int_{t_0}^{t_1} \langle x'_\pi(t), x_0 - s_{t_0} \rangle dt \\
 &= (k^2 + 1)(t_1 - t_0)^2 + d_{S(t_0)}^2(x_0) + 2 \int_{t_0}^{t_1} \langle f(t, x_0), x_0 - s_{t_0} \rangle dt \\
 &\leq (k^2 + 1)(t_1 - t_0)^2 + d_{S(t_0)}^2(x_0) + q(t_0, d_{S(t_0)}^2(x_0))(t_1 - t_0).
 \end{aligned} \tag{3.6}$$

Since similar estimates hold at any node, we have for $i = 1, 2, \dots, N$,

$$d_{S(t_i)}^2(x_i) \leq d_{S(t_{i-1})}^2(x_{i-1}) + (k^2 + 1)(t_i - t_{i-1})^2 + q(t_{i-1}, d_{S(t_{i-1})}^2(x_{i-1}))(t_i - t_{i-1}). \tag{3.7}$$

And therefore, it follows that

$$\begin{aligned}
 d_{S(t_i)}^2(x_i) &\leq d_{S(t_0)}^2(x_0) + (k^2 + 1) \sum_{j=1}^i (t_j - t_{j-1})^2 + \sum_{j=1}^i q(t_{j-1}, d_{S(t_{j-1})}^2(x_{j-1}))(t_j - t_{j-1}) \\
 &\leq d_{S(t_0)}^2(x_0) + (k^2 + 1)\mu_\pi \sum_{j=1}^i (t_j - t_{j-1})^2 + \sum_{j=1}^i q(t_{j-1}, d_{S(t_{j-1})}^2(x_{j-1}))(t_j - t_{j-1}) \\
 &\leq d_{S(t_0)}^2(x_0) + (k^2 + 1)(T - t_0)\mu_\pi + \sum_{j=1}^i q(t_{j-1}, d_{S(t_{j-1})}^2(x_{j-1}))(t_j - t_{j-1}).
 \end{aligned} \tag{3.8}$$

We now consider the sequence $x_{\pi_j}(t)$ of polygonal arcs converging to $x(t)$. Since the last estimate is true at every node, $\mu_{\pi_j} \rightarrow 0$ as $j \rightarrow \infty$, and the same k applies to each x_π , we deduce in the limit the integral inequality

$$d_{S(t)}^2(x(t)) \leq d_{S(t_0)}^2(x_0) + \int_{t_0}^t q(\tau, d_{S(\tau)}^2(x(\tau))) d\tau, \quad t_0 \leq t \leq T, \tag{3.9}$$

which is the same as

$$d_{S(t)}^2(x(t)) \leq r(t, t_0, d_{S(t_0)}^2(x_0)). \tag{3.10}$$

If $r(t, t_0, 0) \equiv 0$, then assuming $x_0 \in S(t_0)$ implies $x(t) \in S(t)$ for $t \geq t_0$ and therefore the system $(S(t), f)$, $t \geq t_0$, is weakly invariant. The proof is complete. $\square$

4. Weak invariance for differential inclusions

Consider the IVP for the differential inclusion

$$x' \in F(t, x), \quad x(t_0) = x_0, \tag{4.1}$$

where F satisfies the following hypotheses:

- (a) F is a nonempty convex set for each $(t, x) \in \mathbb{R}_+ \times \mathbb{R}^n$;
- (b) F is upper semicontinuous;
- (c) $v \in F(t, x)$ implies that $\|v\| \leq g(t, \|x\|)$, where $g \in C[\mathbb{R}_+^2, \mathbb{R}_+]$, $g(t, w)$ is nondecreasing in w , and the maximal solution $r(t) = r(t, t_0, w_0)$, of the scalar differential equation

$$w' = g(t, w), \quad w(0) = w_0 \geq 0, \quad (4.2)$$

exists on $[0, \infty)$.

We recall the notions of lower and upper Hamiltonians, which are functions from $\mathbb{R} \times \mathbb{R}^n \times \mathbb{R}^n$ as follows:

$$h(t, x, p) = \min_{v \in F(t, x)} \langle p, v \rangle, \quad H(t, x, p) = \max_{v \in F(t, x)} \langle p, v \rangle. \quad (4.3)$$

We are now in a position to discuss the existence and weak invariance of $(S(t), F)$.

THEOREM 4.1. *Assume that for each $t \geq t_0$ and every $s_t \in S(t)$,*

$$h(t, s_t, N_{S(t)}^p(s_t)) \leq 0. \quad (4.4)$$

Suppose further that $g(t, u)$ is subadditive in u , for each t . Then the system $(S(t), F)$, $t \geq t_0$, is weakly invariant.

Proof. For each $t \in [t_0, \infty)$ and $x \in \mathbb{R}^n$, choose $s_t = s_t(x) \in \text{proj}_{S(t)}(x)$, and let v_t in $F(t, s_t)$, minimize over $F(t, s_t)$ the function $v_t \rightarrow \langle v_t, x - s_t \rangle$.

Set $f_p(t, x) = v_t$. Since $x - s_t \in N_{S(t)}^p(s_t)$, we have $\langle f_p(t, x), x - s_t \rangle \leq 0$. This implies that the main assumption of Theorem 3.3 with $q(t, u) = 0$ is satisfied. If $s_0 \in S(t_0)$ is a given element, then for each $t \geq t_0$,

$$\begin{aligned} \|f_p(t, x)\| &= \|v_t\| \leq g(t, \|s_t\|) = g(t, \|s_t - x + x\|) \\ &\leq g(t, \|s_t - x\|) + g(t, \|x\|) \\ &\leq g(t, \|s_0 - x\|) + g(t, |t - t_0|^2) + g(t, \|x\|) \\ &\leq 2g(t, \|x\|) + g(t, \|s_0\|) + g(t, |t - t_0|^2) = \tilde{g}(t, \|x\|). \end{aligned} \quad (4.5)$$

Clearly $\tilde{g}(t, u) \in C([t_0, T] \times \mathbb{R}_+, \mathbb{R}_+)$ and $\tilde{g}(t, u)$ is nondecreasing in (t, u) . Thus $f_p(t, x)$ satisfies the nonlinear growth condition required by Theorem 2.1. Thus, by Theorem 3.3, for any $x(0) = x_0$, we have $x(t) \in S(t)$, on $[t_0, \infty)$.

The proof will be complete if we show that $x(t)$ is a solution of (4.2). Since f_p is not a selection of F , let us define another multifunction as follows:

$$\text{for each } t \geq t_0, \quad F_{S(t)}(t, x) = \text{co} \{F(t, s_t) : s_t \in \text{proj}_{S(t)}(x)\}. \quad (4.6)$$

It can be verified that $f_p(t, x)$ is a selection for $F_{S(t)}(t, x)$, that $F_{S(t)}$ satisfies the hypothesis made at the beginning of this section, and that $F_{S(t)}(t, x) = F(t, x)$ for $x \in S(t)$. Since we know that an Euler solution $x(t)$ of any selection f_p of $F_{S(t)}$ is also a solution of (4.2), it follows that $x'(t) \in F_{S(t)}(t, x(t))$ a.e. Since $F = F_{S(t)}$ on $S(t)$ and $x(t) \in S(t)$, $t \geq t_0$, it

follows that $x(t)$ is a solution of (4.2), and therefore $(S(t), F)$ is weakly invariant. The proof is complete. $\square$

5. Strong invariance

We begin with the following definition.

Definition 5.1. The multifunction F is said to be locally Lipschitz in x , uniformly in t , provided that for all $t \in [t_0, \infty)$, each $x \in \mathbb{R}^n$ admits a neighborhood $U = U(x)$ and a positive constant $K = K(x)$ such that

$$x_1, x_2 \in U \implies F(t, x_2) \subseteq F(t, x_1) + K \|x_1 - x_2\| \bar{B}, \quad (5.1)$$

where $\bar{B}$ is the closed unit ball, centred at 0.

For the remainder of this section, we make the following assumption, which is stronger than Assumption 3.2.

ASSUMPTION 5.2. For all $t > \tau$, $t, \tau \in [t_0, \infty)$, and $z \in \mathbb{R}^n$,

$$d_{S(t)}(z) \leq d_{S(\tau)}(z). \quad (5.2)$$

THEOREM 5.3. Let $(S(t), F)$ be weakly invariant and let F be locally Lipschitz in x . Then there exists a feedback selection g_P for F under which $S(t)$ is invariant.

Proof. Let $f_P(t, x)$ be defined as in Theorem 4.1. Then, $f_P(t, x)$ lies in $F(s_t)$, where $s_t \in \text{proj}_{S(t)}(x)$. Define, for each $t \geq t_0$, $g_P(t, x)$ to be an element in $F(T, x)$ closest to $f_P(T, x)$ so that g_P is a selection for F .

Now, suppose $x_0 \in S(t_0)$ and $[t_0, T]$ is any interval. We will show that any Euler solution $y(t)$ on $[t_0, T]$ from x_0 generated by g_P is such that $y(t) \in S(t)$, $t \in [t_0, T]$. We know there is a bound for $y(t)$ on $[t_0, T]$ such that $\|y(t) - x_0\| < M$. Let K be the Lipschitz constant for F on $B[x_0, M_0]$.

If $\|x - x_0\| < M$, then

$$\begin{aligned} \|s_t - x_0\| &\leq \|s_t - x\| + \|x - x_0\| \\ &= d_{S(t)}(x) + \|x - x_0\| \\ &\leq d_{S(t_0)}(x_0) + |t - t_0| + \|x - x_0\| \\ &\leq 2\|x - x_0\| + |T - t_0| \\ &< M_0. \end{aligned} \quad (5.3)$$

Since $\langle s_P(t, x), x - s_t \rangle \leq 0$, we obtain the following estimate:

$$\begin{aligned} \langle g_P(t, x), x - s_t \rangle &= \langle f_P(t, x), x - s_t \rangle + \langle g_P(t, x) - f_P(t, x), x - s_t \rangle \\ &\leq \|g_P(t, x) - f_P(t, x)\| \|x - s_t\|^2 \\ &= K d_{S(t)}^2(x). \end{aligned} \quad (5.4)$$

Thus, by [1, Exercise 2.2], and an application of Gronwall inequality, we get

$$d_{S(t)}(y(t)) \leq d_{S(t_0)}(x_0)e^{Kt}, \quad t \in [t_0, T]. \quad (5.5)$$

Since $x_0 \in S(t_0)$, this implies that $y(t) \in S(t)$, $t \in [t_0, T]$, $T \in (t_0, \infty)$.

We can now prove the strong invariance of the system $(S(t), F)$. $\square$

THEOREM 5.4. *Let F be locally Lipschitz and suppose that for each $t \geq t_0$ and every $s_t \in S(t)$,*

$$H(t, x, N_{S(t)}(s_t)) \leq 0, \quad \forall S(t). \quad (5.6)$$

Then, $(S(t), F)$, $t \geq t_0$, is strongly invariant.

Proof. Let $y(t)$ be any solution for F on $[t_0, T]$ for each t , with $y(t_0) = x_0 \in S(0)$. As a consequence of Theorem 5.3, there exists an f such that $y(t)$ is an Euler solution of the IVP $x' = f(t, x)$, $x(t_0) = x_0$. As in Theorem 5.3, if $M > 0$ is such that all Euler solutions $x(t)$ of this IVP satisfy $\|x(t) - x_0\| < M$, then $\|s_t - x\| \leq M_0$, where $s_t \in \text{proj}_{S(t)}(x)$. This means that $s_t \in B(x_0, M_0)$.

Let K be the Lipschitz constant for F on $B(x_0, M_0)$ and consider any $x \in B(x_0, M_0)$ and $s_t \in \text{proj}_{S(t)}(x)$. Then, $x - s_t \in N_{S(t)}^P(s_t)$. Since $f(t, x) \in F(t, x)$, there exists $v \in F(t, s_t)$ so that

$$\|v - f(t, x)\| \leq K\|s_t - x\| = Kd_{S(t)}(x). \quad (5.7)$$

This leads us to

$$\langle f(t, x), x - s_t \rangle \leq Kd_{S(t)}^2(x). \quad (5.8)$$

Using an argument similar to Theorem 5.3, we conclude that $y(t) \in S(t)$, $t \in [t_0, T]$, since $x_0 \in S(t_0)$. Since $T \in (t_0, \infty)$, we have that $(S(t), F)$, $t \geq t_0$, is strongly invariant and the proof is complete. $\square$

References

- [1] F. H. Clarke, Yu. S. Ledyev, R. J. Stern, and P. R. Wolenski, *Nonsmooth Analysis and Control Theory*, Graduate Texts in Mathematics, vol. 178, Springer, New York, 1998.
- [2] T. Gnana Bhaskar and V. Lakshmikantham, *Generalized proximal normals and flow invariance for differential inclusions*, Nonlinear Studies **10** (2003), no. 1, 29–37.
- [3] G. S. Ladde and V. Lakshmikantham, *On flow-invariant sets*, Pacific Journal of Mathematics **51** (1974), 215–220.

T. Gnana Bhaskar: Department of Mathematical Sciences, Florida Institute of Technology,
Melbourne, FL 32901, USA
E-mail address: gtenali@fit.edu

V. Lakshmikantham: Department of Mathematical Sciences, Florida Institute of Technology,
Melbourne, FL 32901, USA
E-mail address: lakshmik@fit.edu