

Research Article

Wolfe Type Second Order Nondifferentiable Symmetric Duality in Multiobjective Programming over Cone with Generalized (K, F) -Convexity

A. K. Tripathy

Department of Mathematics, Trident Academy of Technology, Chandaka Industrial Estate, F2/A, Bhubaneswar, Odisha 751024, India

Correspondence should be addressed to A. K. Tripathy; arun.tripathy06@rediffmail.com

Received 2 January 2014; Accepted 6 February 2014; Published 8 April 2014

Academic Editors: F. Ding and F. Zirilli

Copyright © 2014 A. K. Tripathy. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

A new class of second order (K, F) pseudoconvex function is introduced with example. A pair of Wolfe type second order nondifferentiable symmetric dual programs over arbitrary cones with square root term is formulated. The duality results are established under second order (K, F) pseudoconvexity assumption. Also a Wolfe type second order minimax mixed integer programming problem is formulated and the symmetric duality results are established under second order (K, F) pseudoconvexity assumption.

1. Introduction

A mathematical programming with two or more objective functions is called multiobjective programming. Often the several objectives are conflicting in nature. Pareto [1] studied multiobjective problems by reducing them to a single objective one. However, the problems were first explicitly defined and studied by Kuhn and Tucker [2]. They also proposed the definition of proper efficiency which was later modified by Geoffrion [3].

In mathematical programming, a pair of primal and dual programs is called symmetric if the dual of the dual is the primal problem. The duality in linear programming is symmetric. It is not so in nonlinear programming in general. Dorn [4], Dantzig et al. [5], and Mond [6] studied symmetric duality in nonlinear programming assuming the kernel function $f(x, y)$ to be convex in x and concave in y . Subsequently, Mond and Weir [7] presented a distinct pair of symmetric dual nonlinear programs which admits the relaxation of the convexity/concavity assumption to pseudoconvexity/pseudoconcavity. Mond [6] initiated second order symmetric duality of Wolfe type in nonlinear programming and proved the duality theorems under second order convexity. Mangasarian [8] discussed second order duality in

nonlinear programming under inclusion condition. Mond [6, page 93] and Mangasarian [8, page 609] also indicated possible computational advantages of the second order dual over the first order dual. This motivated several authors [3, 6, 9–13] in this field. Yang et al. [13] studied second order multiobjective symmetric dual programs and established the duality relations under F -convexity assumptions. Also Yang et al. [12] formulated a pair of Wolfe type second order nondifferentiable symmetric dual programs containing support function and presented the duality results under F convexity.

Recently, Gulati et al. [14] studied Wolfe and Mond-Weir type second order symmetric duality over arbitrary cones and proved the duality results under generalized convexity assumption. Gulati and Geeta [15] studied Mond-Weir type second order symmetric duality in multiobjective programming over cones and established duality results under pseudoconvexity/ K - F convexity assumption. Gulati and Verma [16] formulated a pair of Wolfe type nondifferentiable multiobjective symmetric duality and established the duality results under invexity assumption. Gupta and Kailey [17] formulated a pair of Wolfe type second order nondifferentiable multiobjective symmetric dual programs

in which the objective function contains support function and proved the duality results under second order F -convexity assumption. Gupta and Kailey [18] presented second order multiobjective symmetric duality involving cone-convex functions. Saini and Gulati [19] presented a pair of Wolfe type nondifferentiable second order symmetric dual programs over arbitrary cones under second order (K, F) -convexity assumption.

In this paper, motivated by Saini and Gulati [19], a new class of second order (K, F) pseudoconvex/second order (K, F) strongly pseudoconvex function is introduced with example. A pair of Wolfe type second order nondifferentiable symmetric dual programs over arbitrary cone containing square root term is formulated. The duality results are established under second order (K, F) pseudoconvexity assumption.

2. Notation and Preliminaries

The following convention for vectors in R^n will be used:

$$\begin{aligned} x < y &\iff x_i < y_i, \quad i = 1, 2, \dots, n, \\ x \leq y &\iff x_i \leq y_i, \quad i = 1, 2, \dots, n, \\ x \leq y &\iff x_i \leq y_i, \quad i = 1, 2, \dots, n, \text{ but } x \neq y. \end{aligned} \tag{1}$$

Definition 1. A set C of R^n is called a cone if, for each $x \in C$ and $\lambda \in R, \lambda \geq 0$, we have $\lambda x \in C$. Moreover, if C is convex, then it is convex cone.

Definition 2. The positive polar cone C^* of C is defined as

$$C^* = \{z \in R^n \mid x^T z \geq 0, \forall x \in C\}. \tag{2}$$

Let $C_1 \subset R^n, C_2 \subset R^m$, and $K \subset R^k$ be closed convex cones with nonempty interiors having polars C_1^*, C_2^* , and K^* , respectively. Let $X \subseteq R^n$ and $Y \subseteq R^m$ be open and $X \times Y \subseteq R^n \times R^m$. Let $C_1 \times C_2 \subseteq X \times Y$.

A general multiobjective nonlinear programming problem can be expressed in the following form.

Primal (P).

$$\begin{aligned} \text{Minimize} \quad & f(x) = \{f_1(x), f_2(x), \dots, f_k(x)\} \\ \text{Subject to} \quad & -g(x) \in Q, \quad x \in X, \\ & \text{where } X \subseteq R^n \text{ is open,} \end{aligned} \tag{3}$$

$f : R^n \rightarrow R^k, g : R^n \rightarrow R^l, Q$ is a closed convex cone with nonempty interior in R^m .

Definition 3. A feasible point \bar{x} is weakly efficient solution of (P) if there exist no other $x \in X$ such that $f(x) - f(\bar{x}) < 0$.

Definition 4. A feasible point \bar{x} is efficient solution of (P) if there exist no other $x \in X$ such that $f(x) - f(\bar{x}) \leq 0$.

Definition 5. A function $F : X \times X \times R^n \rightarrow R$ is sublinear in its third argument if, for all $(x, u) \in X \times X$,

$$(1) F(x, u; a_1 + a_2) \leq F(x, u; a_1) + F(x, u; a_2), \text{ for all } a_1, a_2 \in R^n,$$

$$(2) F(x, u; \alpha a) = \alpha F(x, u; a), \text{ for all } \alpha \in R_+.$$

Definition 6. Let $f : R^n \rightarrow R^k$ be thrice differentiable function. f is said to be second order F -pseudoconvex at $u \in X$, if $(x, p) \in X \times R^n$,

$$\begin{aligned} F_{x,u}(\nabla_u f(u) + \nabla_{uu} f(u) p) &\geq 0 \\ \implies f(x) - f(u) + \frac{1}{2} p^T \nabla_u f(u) p &\geq 0. \end{aligned} \tag{4}$$

Now, we are in position to give definition of second order (K, F) -pseudoconvex function and second order strongly (K, F) pseudoconvex function.

Definition 7. The thrice differentiable function $f = (f_1, f_2, \dots, f_k) : R^n \rightarrow R^k$ is said to be second order (K, F) pseudoconvex at $u \in X$, if $(x, p) \in X \times R^n$,

$$\begin{aligned} &-(F_{x,u}(\nabla_u f_1(u) + \nabla_{uu} f_1(u) p_1), \dots, \\ &F_{x,u}(\nabla_u f_k(u) + \nabla_{uu} f_k(u) p_1)) \notin \text{int } K \\ \implies &-\left(f_1(x) - f_1(u) + \frac{1}{2} p_1^T \nabla_u f_1(u) p_1, \dots, f_k(x) \right. \\ &\left. - f_k(u) + \frac{1}{2} p_k^T \nabla_u f_k(u) p_k\right) \notin \text{int } K. \end{aligned} \tag{5}$$

Definition 8. The thrice differentiable function $f : R^n \rightarrow R^k$ is said to be second order strongly (K, F) pseudoconvex at $u \in X$, if $(x, p) \in X \times R^n$,

$$\begin{aligned} &-(F_{x,u}(\nabla_u f_1(u) + \nabla_{uu} f_1(u) p_1), \dots, \\ &F_{x,u}(\nabla_u f_k(u) + \nabla_{uu} f_k(u) p_1)) \notin \text{int } K, \\ \implies &\left(f_1(x) - f_1(u) + \frac{1}{2} p_1^T \nabla_u f_1(u) p_1, \dots, f_k(x) \right. \\ &\left. - f_k(u) + \frac{1}{2} p_k^T \nabla_u f_k(u) p_k\right) \in \text{int } K. \end{aligned} \tag{6}$$

Definition 9. f is second order (K, F) pseudoconcave, if $-f$ is second order (K, F) pseudoconvex, and f is second order strongly (K, F) pseudoconcave, if $-f$ is second order strongly (K, F) pseudoconvex function.

Example 10. Let

$$\begin{aligned}
 K &= \left\{ (x, y) \mid -4x \leq y \leq -\frac{x}{2}, x > 0 \right\}, \\
 f(x) &= (f_1(x), f_2(x)) = (-x^2 + x, e^{-x}), \quad p = 1, \\
 F_{x,u}(a) &= a(x^3 + u), \\
 (\nabla_u f_1(u), \nabla_u f_2(u)) &= (-2u + 1, -e^{-u}), \\
 (\nabla_{uu} f_1(u), \nabla_{uu} f_2(u)) &= (-2, e^{-u}), \\
 (a_1, a_2) &= (\nabla_u f_1(u) + \nabla_{uu} f_1(u) p, \nabla_u f_2(u) + \nabla_{uu} f_2(u) p) \\
 &= (-2u - 1, 0); \\
 (F_{x,u}(a_1), F_{x,u}(a_2)) &= (-2u - 1, 0)(x^3 + u), \\
 \begin{pmatrix} f_1(x) - f_1(u) + \frac{1}{2} p^T \nabla_{uu} f_1(u) p, \\ f_2(x) - f_2(u) + \frac{1}{2} p^T \nabla_{uu} f_2(u) p \end{pmatrix} &= \left(-x^2 + x + u^2 - u - 1, e^{-x} - \frac{1}{2} e^{-u} \right).
 \end{aligned} \tag{7}$$

Now at

$$(F_{x,u}(a_1), F_{x,u}(a_2)) \geq 0 \implies x \leq 0. \tag{8}$$

So at

$$\begin{pmatrix} f_1(x) - f_1(u) + \frac{1}{2} p^T \nabla_{uu} f_1(u) p, \\ f_2(x) - f_2(u) + \frac{1}{2} p^T \nabla_{uu} f_2(u) p \end{pmatrix} \not\geq 0. \tag{9}$$

So f is not second order F convex function.

Now at

$$\begin{aligned}
 u = 0, x = 1, \quad &-(F_{x,u}(a_1), F_{x,u}(a_2)) = (1, 0) \notin \text{int } K, \\
 \begin{pmatrix} f_1(x) - f_1(u) + \frac{1}{2} p^T \nabla_{uu} f_1(u) p, f_2(x) \\ -f_2(u) + \frac{1}{2} p^T \nabla_{uu} f_2(u) p \end{pmatrix} &= \left(-1, e^{-1} - \frac{1}{2} \right) \notin \text{int } K.
 \end{aligned} \tag{10}$$

So f is not second order strongly (K, F) pseudoconvex. But

$$\begin{aligned}
 -\begin{pmatrix} f_1(x) - f_1(u) + \frac{1}{2} p^T \nabla_{uu} f_1(u) p, f_2(x) \\ -f_2(u) + \frac{1}{2} p^T \nabla_{uu} f_2(u) p \end{pmatrix} &= \left(1, -e^{-1} + \frac{1}{2} \right) \notin \text{int } K.
 \end{aligned} \tag{11}$$

So f is second order (K, F) pseudoconvex.

Now we can define second order (K, F) pseudoconvexity for a multiobjective function:

$$f = (f_1, f_2, \dots, f_k) : R^n \times R^m \longrightarrow R^k. \tag{12}$$

Definition 11. A thrice differentiable function $f = (f_1, f_2, \dots, f_k) : R^n \times R^m \rightarrow R^k$ is said to be second order (K, F) -pseudoconvex at $u \in X$, for fixed v , if there exists sublinear function $F : X \times X \times R^n \rightarrow R, q \in R^n, x \in X, v \in Y$ such that

$$\begin{aligned}
 &-(F_{x,u}(\nabla_u f_1(u, v) + \nabla_{uu} f_1(u, v) q_1), \dots, \\
 &F_{x,u}(\nabla_u f_k(u, v) + \nabla_{uu} f_k(u, v) q_k)) \notin \text{int } K \\
 \implies &-\left(f_1(x, v) - f_1(u, v) + \frac{1}{2} q_1^T \nabla_{uu} f_1(u, v) q_1, \dots, \right. \\
 &\left. f_k(x, v) - f_k(u, v) + \frac{1}{2} q_k^T \nabla_{uu} f_k(u, v) q_k \right) \notin \text{int } K.
 \end{aligned} \tag{13}$$

Definition 12. A thrice differentiable function $f = (f_1, f_2, \dots, f_k) : R^n \times R^m \rightarrow R^k$ is said to be second order (K, G) -pseudoconvex at $y \in Y$, for fixed v , if there exists sublinear function $G : Y \times Y \times R^m \rightarrow R, p \in R^m, x \in X, v \in Y$ such that

$$\begin{aligned}
 &-(G_{v,y}(\nabla_y f_1(x, y) + \nabla_{yy} f_1(x, y) p_1), \dots, \\
 &G_{v,y}(\nabla_y f_k(x, y) + \nabla_{yy} f_k(x, y) p_k)) \notin \text{int } K, \\
 \implies &-\left(f_1(x, v) - f_1(x, y) + \frac{1}{2} p_1^T \nabla_{yy} f_1(x, y) p_1, \dots, \right. \\
 &\left. f_k(x, v) - f_k(x, y) + \frac{1}{2} p_k^T \nabla_{yy} f_k(x, y) p_k \right) \\
 &\notin \text{int } K.
 \end{aligned} \tag{14}$$

Lemma 13 (generalized Schwartz inequality). Let B be a positive semidefinite matrix of order n . Then, for all $x, w \in R^n, x^T B w \leq (x^T B x)^{1/2} (w^T B w)^{1/2}$.

The equality holds if $Bx = \lambda Bw$ for some $\lambda \geq 0$.

3. Wolfe Type Second Order Multiobjective Nondifferentiable Dual Programs

We consider the following pair of second order Wolfe type nondifferentiable multiobjective programming problems with k -objective.

Primal (SWP). Consider

$$\begin{aligned}
 & L(x, y, \lambda, w, p) \\
 = & \text{Minimize } \left\{ f_i(x, y) + (x^T B_i x)^{1/2} \right. \\
 & \quad - y^T [\nabla_y f_i(x, y) + \nabla_{yy} f_i(x, y) p] \\
 & \quad \left. - \frac{1}{2} p_i^T [\nabla_{yy} f_i(x, y) p_i], \quad i = 1, 2, \dots, k \right\} \\
 \text{Subject to } & - \sum_{i=1}^k \lambda_i [\nabla_y f_i(x, y) - D_i w_i + \nabla_{yy} f_i(x, y) p_i] \\
 & \quad \in C_2^*, \\
 & \quad \quad \quad (15) \\
 & w_i^T D_i w_i \leq 1, \quad i = 1, 2, \dots, k, \quad (16) \\
 & x \in C_1, \quad w_i \in R^m, \quad (17) \\
 & \lambda \in \text{int } K^*, \quad \sum_{i=1}^k \lambda_i = 1. \quad (18)
 \end{aligned}$$

Dual (SWD). Consider

$$\begin{aligned}
 & M(u, v, \lambda, z, q) \\
 = & \text{Maximize } \left\{ f_i(u, v) - (v^T D_i v)^{1/2} \right. \\
 & \quad - u^T [\nabla_u f_i(u, v) + \nabla_{uu} f_i(u, v) q_i] \\
 & \quad \left. - \frac{1}{2} q_i^T [\nabla_{uu} f_i(u, v) q_i], \quad i = 1, 2, \dots, k \right\} \\
 \text{Subject to } & \sum_{i=1}^k \lambda_i [\nabla_u f_i(u, v) + B_i z_i + \nabla_{uu} f_i(u, v) q_i] \\
 & \quad \in C_1^*, \\
 & \quad \quad \quad (19) \\
 & z_i^T B_i z_i \leq 1, \quad i = 1, 2, \dots, k, \quad (20) \\
 & v \in C_2, \quad z_i \in R^n, \quad (21) \\
 & \lambda \in \text{int } K^*, \quad \sum_{i=1}^k \lambda_i = 1, \quad (22)
 \end{aligned}$$

where

- (1) $f = (f_1, f_2, \dots, f_k) : R^n \times R^m \rightarrow R^k$ is thrice differentiable function,
- (2) C_1 and C_2 are closed convex cones in R^n and R^m with nonempty interiors, respectively,
- (3) C_1^* and C_2^* are positive polar cones of C_1 and C_2 , respectively,
- (4) K is a closed convex cone in R^k with $\text{int } K \neq \emptyset$ and $R_+^k \subset K$,

(5) q_i, z_i ($i = 1, 2, \dots, k$) are vectors in R^n , and p_i, w_i ($i = 1, 2, \dots, k$) are vectors in R^m ,

(6) B_i and D_i ($i = 1, 2, \dots, k$) are positive semidefinite matrices of order n and m , respectively.

Now we establish the following theorem.

Theorem 14 (weak duality theorem). *Let (x, y, λ, p) be a feasible solution for the primal (WP) and let (u, v, λ, q) be a feasible solution for the dual (WD). Suppose there exist sublinear functional $F : X \times X \times R^n \rightarrow R$ and $G : Y \times Y \times R^m \rightarrow R$ satisfying*

$$(1) F_{x,u}(a) - u^T a \geq 0, \text{ for all } (x, u) \in C_1 \times C_1, a \in C_1^*,$$

$$(2) G_{v,y}(b) - y^T b \geq 0, \text{ for all } (v, y) \in C_2 \times C_2, b \in C_2^*.$$

Furthermore assume that, for each i , $f_i(\cdot, v) + (\cdot)^T B_i z_i$ is second order (K, F) -pseudoconvex at u for fixed v and $f_i(x, \cdot) + (\cdot)^T D_i w_i$ is second order pseudoconcave at y for fixed x :

$$\text{Inf (WP)} - \text{Sup (WD)} \in \text{int } K. \quad (23)$$

Proof. Since (u, v, λ, q) is feasible solution for (WD), from dual constraint (5) we have

$$a = \sum_{i=1}^k \lambda_i [\nabla_u f_i(u, v) + B_i z_i + \nabla_{uu} f_i(u, v) q_i] \in C_1^*. \quad (24)$$

So

$$\begin{aligned}
 u \in C_1 & \implies u^T a \\
 & = u^T \left(\sum_{i=1}^k \lambda_i [\nabla_u f_i(u, v) + B_i z_i + \nabla_{uu} f_i(u, v) q_i] \right) \geq 0.
 \end{aligned} \quad (25)$$

Again hypothesis (1) implies $F_{x,u}(a) \geq 0$. Consider

$$\implies F_{x,u} \left(\sum_{i=1}^k \lambda_i [\nabla_u f_i(u, v) + B_i z_i + \nabla_{uu} f_i(u, v) q_i] \right) \geq 0. \quad (26)$$

Since F is sublinear with respect to third argument,

$$\implies \sum_{i=1}^k \lambda_i F_{x,u} ([\nabla_u f_i(u, v) + B_i z_i + \nabla_{uu} f_i(u, v) q_i]) \geq 0. \quad (27)$$

Since $\lambda \in \text{int } K$, the above inequality can be written as

$$\begin{aligned}
 & - (F_{x,u} ([\nabla_u f_1(u, v) + B_1 z_1 + \nabla_{uu} f_1(u, v) q_1]), \dots, \\
 & \quad F_{x,u} ([\nabla_u f_k(u, v) + B_k z_k + \nabla_{uu} f_k(u, v) q_k])) \notin \text{int } K.
 \end{aligned} \quad (28)$$

So second order (K, F) -pseudoconvexity of $f_i(\cdot, v) + (\cdot)^T B_i z_i$ at u for fixed v implies that

$$\begin{aligned}
 & - \left(\begin{aligned} & f_1(x, v) + (x)^T B_1 z_1 - f_1(u, v) - (u)^T B_1 z_1 \\ & + \frac{1}{2} q_1^T \nabla_{uu} f_1(u, v) q_1, \dots, f_k(x, v) + (x)^T B_k z_k \\ & - f_k(u, v) - (u)^T B_k z_k + \frac{1}{2} q_k^T \nabla_{uu} f_k(u, v) q_k \end{aligned} \right) \notin \text{int } K.
 \end{aligned} \quad (29)$$

This implies that, for $\lambda \in \text{int } K$,

$$\begin{aligned}
 & -\sum_{i=1}^k \lambda_i \left[f_i(x, v) + (x)^T B_i z_i - f_i(u, v) \right. \\
 & \quad \left. - (u)^T B_i z_i - \frac{1}{2} q_i^T \nabla_{uu} f_i(u, v) q_i \right] \not\geq 0 \\
 \implies & \sum_{i=1}^k \lambda_i \left[f_i(x, v) + (x)^T B_i z_i - f_i(u, v) \right. \\
 & \quad \left. - (u)^T B_i z_i - \frac{1}{2} q_i^T \nabla_{uu} f_i(u, v) q_i \right] \geq 0.
 \end{aligned} \tag{30}$$

Similarly (u, v, λ, q) is feasible solution for (WD), so from primal constraint (1) we have

$$b = -\sum_{i=1}^k \lambda_i \left[\nabla_y f_i(x, y) - D_i w_i + \nabla_{yy} f_i(x, y) p_i \right] \in C_2^*. \tag{31}$$

So

$$\begin{aligned}
 y \in C_2 & \implies y^T b \\
 & = -y^T \left(\sum_{i=1}^k \lambda_i \left[\nabla_y f_i(x, y) \right. \right. \\
 & \quad \left. \left. - D_i w_i + \nabla_{yy} f_i(x, y) p_i \right] \right) \geq 0.
 \end{aligned} \tag{32}$$

Again hypothesis (2) implies $G_{v,y}(b) \geq 0$. Consider

$$\begin{aligned}
 \implies & G_{v,y} \left(-\sum_{i=1}^k \lambda_i \left[\nabla_y f_i(x, y) - D_i w_i + \nabla_{yy} f_i(x, y) p_i \right] \right) \\
 & \geq 0.
 \end{aligned} \tag{33}$$

Since G is sublinear with respect to third argument,

$$\begin{aligned}
 \implies & \sum_{i=1}^k \lambda_i G_{v,y} \left(-\left[\nabla_y f_i(x, y) - D_i w_i + \nabla_{yy} f_i(x, y) p_i \right] \right) \\
 & \geq 0.
 \end{aligned} \tag{34}$$

Since $\lambda \in \text{int } K$, the above inequality can be written as

$$\implies -\left(G_{v,y} \left(-\left[\nabla_y f_1(x, y) - D_1 w_1 + \nabla_{yy} f_1(x, y) p_1 \right] \right), \dots \right) \notin \text{int } K. \tag{35}$$

So second order (K, F) -pseudoconcavity of $f_i(x, \cdot) - (\cdot)^T D_i w_i$ at y for fixed x implies that

$$\begin{aligned}
 & -\left(-\left[f_1(x, v) - (v)^T D_1 w_1 - f_1(x, y) + (y)^T D_1 w_1 \right. \right. \\
 & \quad \left. \left. + \frac{1}{2} p_1^T \nabla_{yy} f_1(x, y) p_1 \right], \dots, \right. \\
 & \quad \left. -\left[f_k(x, v) - (v)^T D_k w_k - f_k(x, y) + (y)^T D_k w_k \right. \right. \\
 & \quad \left. \left. + \frac{1}{2} p_k^T \nabla_{yy} f_k(x, y) p_k \right] \right)
 \end{aligned} \tag{36}$$

$\notin \text{int } K$.

This implies that, for $\lambda \in \text{int } K$,

$$\begin{aligned}
 & -\sum_{i=1}^k \lambda_i \left[-\left[f_k(x, v) - (v)^T D_k w_k - f_k(x, y) \right. \right. \\
 & \quad \left. \left. + (y)^T D_k w_k + \frac{1}{2} p_k^T \nabla_{yy} f_k(x, y) p_k \right] \right] \not\geq 0 \\
 \implies & \sum_{i=1}^k \lambda_i \left[-f_i(x, v) + (v)^T D_i w_i + f_i(x, y) \right. \\
 & \quad \left. - (y)^T D_i w_i - \frac{1}{2} p_i^T \nabla_{yy} f_i(x, y) p_i \right] \geq 0.
 \end{aligned} \tag{37}$$

Adding (30) and (37), we get

$$\begin{aligned}
 \implies & \sum_{i=1}^k \lambda_i \left[f_i(x, y) + (x)^T B_i z_i - (y)^T D_i w_i \right. \\
 & \quad \left. - \frac{1}{2} p_i^T \nabla_{yy} f_i(x, y) p_i \right] \\
 & - \sum_{i=1}^k \lambda_i \left[f_i(u, v) - (v)^T D_i w_i + u^T B_i z_i \right. \\
 & \quad \left. - \frac{1}{2} q_i^T \nabla_{uu} f_i(u, v) q_i \right] \geq 0.
 \end{aligned} \tag{38}$$

Now from Schwartz inequality (Lemma 13), (17), and (21), we have

$$\begin{aligned}
 x^T B_i z_i & \leq (x^T B_i x)^{1/2} (z_i^T B_i z_i)^{1/2} \leq (x^T B_i x)^{1/2}, \\
 & \quad i = 1, 2, \dots, k, \\
 v^T D_i w_i & \leq (v^T D_i v)^{1/2} (w_i^T D_i w_i)^{1/2} \leq (v^T D_i v)^{1/2}, \\
 & \quad i = 1, 2, \dots, k.
 \end{aligned} \tag{39}$$

Also from primal constraint (15), we have

$$\sum_{i=1}^k \lambda_i \left[\nabla_y f_i(x, y) - D_i w_i + \nabla_{yy} f_i(x, y) p_i \right] \in C_2^*. \tag{40}$$

For $y \in C_1$, $y^T (\sum_{i=1}^k \lambda_i [\nabla_y f_i(x, y) - D_i w_i + \nabla_{yy} f_i(x, y) p_i]) \geq 0$

$$\begin{aligned} & - \sum_{i=1}^k \lambda_i [y^T D_i w_i] \\ & \geq -y^T \left(\sum_{i=1}^k \lambda_i [\nabla_y f_i(x, y) + \nabla_{yy} f_i(x, y) p_i] \right). \end{aligned} \quad (41)$$

Similarly from dual constraint (19), we have

$$\begin{aligned} & \Rightarrow - \sum_{i=1}^k \lambda_i [u^T B_i z_i] \\ & \geq -u^T \left(\sum_{i=1}^k \lambda_i [\nabla_u f_i(u, v) + \nabla_{uu} f_i(u, v) q_i] \right). \end{aligned} \quad (42)$$

Using (39), (41), and (42) in (38), we obtain that

$$\begin{aligned} & \sum_{i=1}^k \lambda_i \left\{ f_i(x, y) + (x^T B_i x)^{1/2} \right. \\ & \quad \left. - y^T [\nabla_y f_i(x, y) + \nabla_{yy} f_i(x, y) p_i] \right. \\ & \quad \left. - \frac{1}{2} p_i^T [\nabla_{yy} f_i(x, y) p_i] \right\} \\ & - \sum_{i=1}^k \lambda_i \left\{ f_i(u, v) - (v^T D_i v)^{1/2} \right. \\ & \quad \left. - u^T [\nabla_u f_i(u, v) + \nabla_{uu} f_i(u, v) q_i] \right. \\ & \quad \left. - \frac{1}{2} q_i^T [\nabla_{uu} f_i(u, v) q_i] \right\} \geq 0 \\ & \Rightarrow \text{Inf (WP)} - \text{Sup (WD)} \in \text{int } K. \end{aligned} \quad (43)$$

□

Theorem 15 (strong duality). *Let $(\bar{x}, \bar{y}, \bar{\lambda}, \bar{w}, \bar{p})$ be weakly efficient solution of (WP) such that*

- (i) $\nabla_{yy} f_i(\bar{x}, \bar{y})$ is nonsingular,
- (ii) the matrix $\sum_{i=1}^k \lambda_i [\nabla_y (\nabla_{yy} f_i(\bar{x}, \bar{y}) \bar{p})]$ is positive definite,
- (iii) the set $\{\nabla_y f_1 - C_1 w_1, \dots, \nabla_y f_k - C_k w_k\}$ is linearly independent,
- (iv) $\sum_{i=1}^k \bar{\lambda}_i (\nabla_y (\nabla_{yy} f_i \bar{p})) \bar{p} \notin \text{span}\{\nabla_y f_1 - C_1 w_1, \dots, \nabla_y f_k - C_k w_k\} \setminus \{0\}$.

Then there exist $\bar{z}_i \in R^n$ such that $(\bar{x}, \bar{y}, \bar{\lambda}, \bar{z}, \bar{q} = 0)$ is feasible for (WD) and two objective values of (WP) and (WD) are equal. Also, if the hypotheses of Theorem 14 are satisfied for all feasible solution of (WP) and (WD), then $(\bar{x}, \bar{y}, \bar{\lambda}, \bar{z}, \bar{q} = 0)$ is an efficient solution of (WD).

Proof. Since $(\bar{x}, \bar{y}, \bar{\lambda}, \bar{w}, \bar{p})$ is weakly efficient solution of (WD), by the Fritz-John necessary optimality condition on

convex cone domain given in Bazaraa and Goode [20], there exist $\alpha \in K^*$, $\beta \in C_2$, $\gamma, \tau_i \in R_+$ such that the following conditions are satisfied at $(\bar{x}, \bar{y}, \bar{\lambda}, \bar{w}, \bar{p})$:

$$\begin{aligned} & (x - \bar{x})^T \\ & \times \left(\sum_{i=1}^k \alpha_i [\nabla_x f_i + B_i z_i] + \sum_{i=1}^k \lambda_i (\nabla_{yx} f_i)^T (\beta - \gamma \bar{y}) \right. \\ & \quad \left. + \sum_{i=1}^k \lambda_i (\nabla_x (\nabla_{yy} f_i \bar{p}))^T [\beta - \gamma \bar{y} - \alpha_i \bar{p}_i] \right) \end{aligned} \quad (44)$$

$$\geq 0, \quad \forall x \in C_1,$$

$$\begin{aligned} & \sum_{i=1}^k (\alpha_i - \gamma \lambda_i) (\nabla_y f_i - C_i w_i) \\ & + \sum_{i=1}^k \lambda_i (\nabla_{yy} f_i)^T [\beta - \gamma (\bar{y} + \bar{p})] \end{aligned} \quad (45)$$

$$+ \sum_{i=1}^k \lambda_i (\nabla_y (\nabla_{yy} f_i \bar{p}_i))^T \left(\beta - \gamma \bar{y} - \frac{1}{2} \gamma \bar{p}_i \right) = 0,$$

$$\sum_{i=1}^k \lambda_i (\nabla_{yy} f_i)^T (\beta - \gamma \bar{y} - \gamma \bar{p}_i) = 0, \quad (46)$$

$$\sum_{i=1}^k \lambda_i (\nabla_y f_i - C_i w_i + \nabla_{yy} f_i \bar{p}_i)^T \beta = 0, \quad (47)$$

$$x^T B_i z_i = (x^T B_i x)^{1/2}, \quad i = 1, 2, \dots, k, \quad (48)$$

$$\bar{\lambda} = (\bar{\lambda}_1, \bar{\lambda}_2, \dots, \bar{\lambda}_k) \in \text{int } K, \quad (49)$$

$$\alpha_i C_i \bar{y} + (\beta - \lambda \bar{y}) \lambda_i C_i = 2\tau_i C_i w_i, \quad (50)$$

$$\tau_i (\bar{w}_i^T C_i \bar{w}_i - 1) = 0, \quad (51)$$

$$\bar{z}_i^T B_i \bar{z}_i \leq 1, \quad (52)$$

$$(\alpha, \beta, \gamma, \tau) \geq 0, \quad (53)$$

$$(\alpha, \beta, \gamma, \tau) \neq 0. \quad (54)$$

Since $\nabla_{yy} f_i$ is nonsingular, (46) implies that $\beta = \gamma(\bar{y} + \bar{p}_i)$

$$\beta = \gamma(\bar{y} + \bar{p}_i). \quad (55)$$

We claim that $\gamma > 0$. Indeed if $\gamma = 0$, then (55) implies $\beta = 0$, which contradicts (54).

Hence

$$\gamma > 0. \quad (56)$$

Since $\gamma > 0$, using (55) in (45), we get

$$\sum_{i=1}^k \lambda_i [\nabla_y (\nabla_{yy} f_i \bar{p})]^T \bar{p}_i = \frac{-2}{\gamma} \sum_{i=1}^k (\alpha_i - \gamma \lambda_i) [\nabla_y f_i - C_i w_i], \quad (57)$$

which by hypothesis (ii) and (iv) yields

$$\bar{p}_i = 0, \quad i = 1, 2, \dots, k. \tag{58}$$

From (55) and (58), we obtain

$$\beta = \gamma \bar{y}. \tag{59}$$

Using (58) and (59) and hypothesis (iii) in (45), we get

$$\alpha = \gamma \lambda. \tag{60}$$

Again using (58), (59), and (60) in (44), we get

$$(x - \bar{x})^T \sum_{i=1}^k \lambda_i [\nabla_x f_i + B_i z_i] \geq 0, \quad \forall x \in C_1. \tag{61}$$

Let $x \in C_1$. Then $x + \bar{x} \in C_1$ and so (61) implies

$$\begin{aligned} (x)^T \sum_{i=1}^k \lambda_i [\nabla_x f_i + B_i z_i] &\geq 0, \quad \forall x \in C_1 \\ \implies \sum_{i=1}^k \lambda_i [\nabla_x f_i + B_i z_i] &\in C_1^*. \end{aligned} \tag{62}$$

Also from (56), (59), and $\beta \in C_2$, we obtain

$$\bar{y} \in C_2. \tag{63}$$

Thus, from (52), (62), and (63), we obtain that $(\bar{x}, \bar{y}, \bar{z}, \bar{\lambda}, \bar{q} = 0)$ satisfies the dual constraints (19), (20), (21), and (22).

Thus $(\bar{x}, \bar{y}, \bar{z}, \bar{\lambda}, \bar{q} = 0)$ is efficient for (WD).

Let $(2\tau_i/\alpha_i) = t$; then $t \geq 0$. From (50) and (59), we get

$$C_i \bar{y} = t C_i \bar{w}_i. \tag{64}$$

This is a condition of Schwartz inequality:

$$\bar{y} C_i \bar{w}_i = (\bar{y}^T C_i \bar{w}_i)^{1/2} (\bar{w}_i^T C_i \bar{w}_i). \tag{65}$$

In case $\tau_i > 0$, from (51) we get $\bar{w}_i^T C_i \bar{w}_i = 1$. So (65) implies $\bar{y} C_i \bar{w}_i = (\bar{y}^T C_i \bar{w}_i)^{1/2}$.

In case $\tau_i = 0$, we get $t = 0$. So $C_i \bar{y} = 0$. Hence $\bar{y} C_i \bar{w}_i = (\bar{y}^T C_i \bar{w}_i)^{1/2}$. Thus in either case

$$\bar{y} C_i \bar{w}_i = (\bar{y}^T C_i \bar{w}_i)^{1/2}. \tag{66}$$

So using (48) and (66), we obtained that the two objective values are equal; that is,

$$L(\bar{x}, \bar{y}, \bar{w}, \bar{\lambda}, \bar{p} = 0) = M(\bar{x}, \bar{y}, \bar{z}, \bar{\lambda}, \bar{q} = 0). \tag{67}$$

Now we claim that $(\bar{x}, \bar{y}, \bar{z}, \bar{\lambda}, \bar{q} = 0)$ is an efficient solution of (WD). If this would not be the case, then there would exist a feasible solution $(\bar{u}, \bar{v}, \bar{z}, \bar{\lambda}, \bar{q} = 0)$ such that

$$\begin{aligned} M(\bar{x}, \bar{y}, \bar{z}, \bar{\lambda}, \bar{q} = 0) &\leq M(\bar{u}, \bar{v}, \bar{z}, \bar{\lambda}, \bar{q} = 0) \\ \implies L(\bar{x}, \bar{y}, \bar{z}, \bar{\lambda}, \bar{q} = 0) &\leq M(\bar{u}, \bar{v}, \bar{z}, \bar{\lambda}, \bar{q} = 0). \end{aligned} \tag{68}$$

This is a contradiction to weak duality Theorem 14.

Hence $(\bar{x}, \bar{y}, \bar{z}, \bar{\lambda}, \bar{q} = 0)$ is efficient solution. \square

Theorem 16 (converse duality theorem). *Let $(\bar{x}, \bar{y}, \bar{\lambda}, \bar{w}, \bar{p})$ be a weakly efficient solution of (WP) such that*

- (i) $\nabla_{uu} f_i(\bar{u}, \bar{v})$ is nonsingular,
- (ii) the matrix $\sum_{i=1}^k \lambda_i [\nabla_u (\nabla_{uu} f_i(\bar{u}, \bar{v}) \bar{q})]$ is positive definite,
- (iii) the set $\{\nabla_u f_1 + B_1 z_1, \dots, \nabla_u f_k + B_k z_k\}$ is linearly independent,
- (iv) $\sum_{i=1}^k \bar{\lambda}_i (\nabla_u (\nabla_{uu} f_i \bar{q})) \bar{q} \notin \text{span}\{\nabla_u f_1 + B_1 z_1, \dots, \nabla_u f_k + B_k z_k\} \setminus \{0\}$.

Then there exist $\bar{w}_i \in R^m$ such that $(\bar{u}, \bar{v}, \bar{\lambda}, \bar{w}, \bar{p} = 0)$ is feasible for (WD) and two objective values of (WP) and (WD) are equal. Also, if the hypotheses of Theorem 14 are satisfied for all feasible solution of (WP) and (WD), then $(\bar{u}, \bar{v}, \bar{\lambda}, \bar{w}, \bar{p} = 0)$ is an efficient solution of (WD).

Proof. The proof follows on lines of Theorem 15. \square

4. Wolfe Type Minimax Mixed Integer Programming

Let U and V be two arbitrary sets of integers in R^{n_1} ($0 \leq n_1 \leq n$) and R^{m_1} ($0 \leq m_1 \leq m$), respectively. Throughout this section, we constrained some of the components of the vector variables $x \in R^n$ and $y \in R^m$ to belong to arbitrary sets of integers U and V , respectively. Then we write $(x, y) = (x^1, x^2, y^1, y^2)$, where $x^1 = (x_1, x_2, \dots, x_n)$ and $y^1 = (y_1, y_2, \dots, y_m)$. x^2 and y^2 are the vectors of the remaining components of x and y , respectively.

Definition 17. Let s^1, s^2, \dots, s^r be elements of an arbitrary vector space. A vector function $\theta(s^1, s^2, \dots, s^r)$ will be called additively separable with respect to s^1 , if there exist vector function $\theta_1(s^1)$ (independent of s^2, \dots, s^r) and $\theta_2(s^2, \dots, s^r)$ (independent of s^1) such that $\theta(s^1, s^2, \dots, s^r) = \theta_1(s^1) + \theta_2(s^2, \dots, s^r)$.

We consider the following pair of Wolfe type nondifferentiable minimax mixed integer symmetric primal and dual programs:

Primal (WIP). Consider

$$\begin{aligned} \max_{x^1} \min_{x^2, y, w, p} &\left\{ f_i(x, y) + \left((x^2)^T B_i x \right)^{1/2} \right. \\ &- (y^2)^T [\nabla_{y^2} f_i(x, y) + \nabla_{y^2 y^2} f_i(x, y) p] \\ &\left. - \frac{1}{2} p_i^T [\nabla_{y^2 y^2} f_i(x, y) p_i], \quad i = 1, 2, \dots, k \right\} \\ \text{Subject to} &- \sum_{i=1}^k \lambda_i [\nabla_{y^2} f_i(x, y) - D_i w_i + \nabla_{y^2 y^2} f_i(x, y) p_i] \\ &\in C_2^*, \end{aligned}$$

$$\begin{aligned}
 &w_i^T D_i w_i \leq 1, \quad i = 1, 2, \dots, k, \\
 &x^2 \in C_1, \quad x^1 \in U, \quad y^1 \in V, \quad w_i \in R^m, \\
 &\lambda \in \text{int } K^*, \quad \sum_{i=1}^k \lambda_i = 1.
 \end{aligned} \tag{69}$$

Dual (WID). Consider

$$\begin{aligned}
 &\min_{v^1} \max_{u, v^2, z} \left\{ f_i(u, v) - \left((v^2)^T D_i v \right)^{1/2} \right. \\
 &\quad \left. + (u^2)^T [\nabla_{u^2} f_i(u, v) + \nabla_{u^2 u^2} f_i(u, v) q_i] \right. \\
 &\quad \left. - \frac{1}{2} q_i^T [\nabla_{u^2 u^2} f_i(u, v) q_i], \quad i = 1, 2, \dots, k \right\} \\
 &\text{Subject to } \sum_{i=1}^k \lambda_i [\nabla_{u^2} f_i(u, v) + B_i z_i + \nabla_{u^2 u^2} f_i(u, v) q_i] \\
 &\quad \in C_1^*, \\
 &\quad z_i^T B_i z_i \leq 1, \quad i = 1, 2, \dots, k, \\
 &\quad v^2 \in C_2, \quad u^1 \in U, \quad v^1 \in V, \quad z_i \in R^n, \\
 &\quad \lambda \in \text{int } K^*, \quad \sum_{i=1}^k \lambda_i = 1,
 \end{aligned} \tag{70}$$

where

- (1) $f = (f_1, f_2, \dots, f_k) : R^n \times R^m \rightarrow R^k$ is thrice differentiable function,
- (2) C_1 and C_2 are closed convex cones in R^{n-n_1} and R^{m-m_1} with nonempty interiors, respectively,
- (3) C_1^* and C_2^* are positive polar cones of C_1 and C_2 , respectively,
- (4) K is a closed convex cone in R^k with $\text{int } K \neq \emptyset$ and $R_+^k \subset K$,
- (5) q_i, z_i ($i = 1, 2, \dots, k$) are vectors in R^n , and p_i, w_i ($i = 1, 2, \dots, k$) are vectors in R^m ,
- (6) B_i and D_i ($i = 1, 2, \dots, k$) are positive semidefinite matrices of order $n - n_1$ and $m - m_1$, respectively.

Theorem 18 (symmetric duality). Let $(\bar{x}, \bar{y}, \bar{\lambda}, \bar{z}, \bar{p})$ be a weakly efficient solution of (WIP). Also

- (i) $f(x, y)$ is additively separable with respect to x^1 or y^1 ; that is, $f_i(x^1, x^2, y) = f_{i1}(x^1) + f_{i2}(x^2, y)$,
- (ii) $f(x, y)$ is thrice differentiable in x^2 and y^2 ,
- (iii) $\nabla_{y^2 y^2} f(\bar{x}, \bar{y})$ is nonsingular,
- (iv) the vector $\bar{p}^T \nabla_{y^2 y^2} f(\bar{x}, \bar{y}) \bar{p} = 0 \Rightarrow \bar{p} = 0$.
Furthermore, for any feasible solution (x, y, λ, w, p)

in (WIP) and for any feasible solution (u, v, λ, z, q) in (WID), suppose there exist functional $F' : X^1 \times X^1 \times R^{n-n_1} \rightarrow R$ and $G' : Y^1 \times Y^1 \times R^{m-m_1} \rightarrow R$ such that

- (v) $(f_{12}(u^2, v) + (u^2)^T B_1 z_1, \dots, f_{k2}(u^2, v) + (u^2)^T B_k z_k)$ is second order (K, F') pseudoconvex at u^2 with respect to $q \in R^{n-n_1}$ for each (u^1, v) and $(f_{12}(x, y^2) - (y^2)^T D_1 w_1, \dots, f_{k2}(x, y^2) - (y^2)^T D_k w_k)$ is second order (K, G') pseudoconcave at y^2 with respect to $p \in R^{m-m_1}$ for each (x, y^1) ,
- (vi) $F'_{x^2, u^2}(a) - (u^2)^T a \geq 0$, for all $a \in C_1^*$, and $G'_{y^2, y^2}(b) - (y^2)^T b \geq 0$, for all $b \in C_2^*$.

Then there exist $z \in R^{n-n_1}$ such that $(\bar{x}, \bar{y}, \bar{\lambda}, \bar{z}, \bar{q} = 0)$ is efficient solution for dual and optimal values (WIP) and (WID) are equal.

Proof. Let

$$\begin{aligned}
 s &= \max_{x^1} \min_{x^2, y, w} \left\{ f_i(x, y) + \left((x^2)^T B_i x^2 \right)^{1/2} - (y^2)^T \right. \\
 &\quad \times [\nabla_{y^2} f_i(x, y) + \nabla_{y^2 y^2} f_i(x, y) p_i] \\
 &\quad \left. - \frac{1}{2} p_i^T \nabla_{y^2 y^2} f_i(x, y) p_i, \right. \\
 &\quad \left. i = 1, 2, \dots, k; (x, y, w, p) \in S \right\}, \\
 t &= \min_{v^1} \max_{u, v^2, z} \left\{ f_i(u, v) - \left((v^2)^T C_i v^2 \right)^{1/2} \right. \\
 &\quad \left. + (u^2)^T [\nabla_{u^2} f_i(u, v) + \nabla_{u^2 u^2} f_i(u, v) q_i] \right. \\
 &\quad \left. - \frac{1}{2} q_i^T \nabla_{u^2 u^2} f_i(u, v) q_i, \right. \\
 &\quad \left. i = 1, 2, \dots, k; (u, v, z, q) \in T \right\},
 \end{aligned} \tag{71}$$

where S and T are a feasible region of primal (WIP) and dual (WID), respectively.

Since $f_i(x, y)$ is additively separable with respect to x^1 or y^1 (say with respect to x^1) from definition, it follows that $f_i(x, y) = f_{i1}(x^1) + f_{i2}(x^2, y)$.

Therefore

$$\begin{aligned}
 \nabla_{y^2} f_i(x, y) &= \nabla_{y^2} f_{i2}(x^2, y), \quad \nabla_{y^2 y^2} f_i(x, y) \\
 &= \nabla_{y^2 y^2} f_{i2}(x^2, y).
 \end{aligned} \tag{72}$$

So the primal (WIP) can be written as

$$s = \max_{x^1} \min_{x^2, y, w} \left\{ f_{i1}(x^1) + f_{i2}(x^2, y) + \left((x^2)^T B_i x^2 \right)^{1/2} - (y^2)^T \times \left[\nabla_{y^2} f_{i2}(x, y) + \nabla_{y^2 y^2} f_{i2}(x, y) p_i \right] - \frac{1}{2} p_i^T \nabla_{y^2 y^2} f_{i2}(x^2, y) p_i, \right. \\ \left. i = 1, 2, \dots, k; (x, y, w, p) \in S \right\} \quad (73)$$

or

$$s = \max_{x^1} \min_{y^1} \left\{ f_{i1}(x^1) + \varphi_i(y^1); x^1 \in U, y^1 \in V \right\}, \quad (74)$$

where (WIP₀):

$$\varphi_i(y^1) = \min_{x^2, y^2, w} \left\{ f_{i2}(x^2, y) + \left((x^2)^T B_i x^2 \right)^{1/2} - (y^2)^T \times \left[\nabla_{y^2} f_{i2}(x, y) + \nabla_{y^2 y^2} f_{i2}(x, y) p_i \right] - \frac{1}{2} p_i^T \nabla_{y^2 y^2} f_{i2}(x^2, y) p_i \right\} \\ \text{Subject to } - \sum_{i=1}^k \lambda_i \left[\nabla_{y^2} f_{i2}(x^2, y) - C_i w_i + \nabla_{y^2 y^2} f_{i2}(x^2, y) p_i \right] \in C_2^*, \\ (w_i)^T C_i w_i \leq 1, \quad i = 1, 2, \dots, k, \\ \lambda \in \text{int } K^*, \quad \sum_{i=1}^k \lambda_i = 1, \\ x^2 \in C_1, \quad w_i \in R^{m-m_1}, \quad p_i \in R^{m-m_1}. \quad (75)$$

Similarly the dual (WID) can be written as

$$t = \min_{v^1} \max_{u^1} \left\{ f_{i1}(u^1) + \theta(v^1), u^1 \in U, v^1 \in V \right\}, \quad (76)$$

where (WID₀):

$$\theta_i(v^1) = \max_{u^2, v^2, z, q} \left\{ f_{i2}(u^2, v) - \left((v^2)^T C_i v^2 \right)^{1/2} - (u^2)^T \times \left[\nabla_{u^2} f_{i2}(u, v) + \nabla_{u^2 u^2} f_{i2}(u, v) q_i \right] - \frac{1}{2} q_i^T \nabla_{u^2 u^2} f_{i2}(u^2, v) q_i \right\} \\ \text{Subject to } \sum_{i=1}^k \lambda_i \left[\nabla_{u^2} f_{i2}(u^2, v) + B_i z_i + \nabla_{u^2 u^2} f_{i2}(u^2, v) q_i \right] \in C_1^*, \\ (z_i)^T B_i z_i \leq 1, \quad i = 1, 2, \dots, k, \\ \lambda \in \text{int } K^*, \quad \sum_{i=1}^k \lambda_i = 1, \\ v^2 \in C_2, \quad z_i \in R^{n-n_1}, \quad q_i \in R^{n-n_1}. \quad (77)$$

For any given y^1 and v^1 , programs (WIP₀) and (WID₀) are a pair of Wolfe type second order nondifferentiable multiobjective symmetric dual programs studied in Section 3 and hence in view of hypothesis (ii)-(vi), Theorems 14 and 15 become applicable. Therefore $y^1 = \bar{y}^1 = v^1$, and we obtain $\bar{q} = 0$, $\varphi(y^1) = \theta(\bar{y}^1)$. So the two optimal values are equal and $(\bar{x}, \bar{y}, \bar{\lambda}, \bar{z}, \bar{q} = 0)$ is an efficient solution for the dual. □

5. Special Cases

(i) If $B_i = D_i = 0, k = 1$, then the problems (SWP) and (SWD) can be reduced to the problem proposed by Gulati et al. [14] as follows.

Primal (WP).

$$\text{Minimize } f(x, y) - y^T \left[\nabla_y f(x, y) + \nabla_{yy} f(x, y) p \right] - \frac{1}{2} p^T \left[\nabla_{yy} f(x, y) p \right] \\ \text{Subject to } - \left[\nabla_y f_i(x, y) + \nabla_{yy} f_i(x, y) p_i \right] \in C_2^*, \\ x \in C_1. \quad (78)$$

Dual (WD).

$$\text{Maximize } f(u, v) + u^T \left[\nabla_u f(u, v) + \nabla_{uu} f(u, v) q \right] - \frac{1}{2} q^T \left[\nabla_{uu} f(u, v) q \right] \quad (79)$$

$$\text{Subject to } \nabla_u f(u, v) + \nabla_{uu} f(u, v) q \in C_1^*, \\ v \in C_2.$$

(ii) If $k = 1$, $C_1 = R_+^n$, $C_2 = R_+^m$, $(x^T Bx)^{1/2} = s(x | C')$, and $(y^T Dy)^{1/2} = s(y | D')$, where $C' = \{Bx | x^T Bx \leq 1\}$, $D' = \{Dy | y^T Dy \leq 1\}$, then the problems (SWP) and (SWD) can be reduced to the problem proposed by Yang et al. [12].

Primal (WP).

$$\begin{aligned} \text{Minimize } & f(x, y) + s(x | C') \\ & - y^T [\nabla_y f(x, y) + \nabla_{yy} f(x, y) p] \\ & - \frac{1}{2} p^T [\nabla_{yy} f(x, y) p] \end{aligned} \quad (80)$$

$$\begin{aligned} \text{Subject to } & \nabla_y f(x, y) - z + \nabla_{yy} f(x, y) p \leq 0, \\ & x \geq 0, \quad z \in D'. \end{aligned}$$

Dual (WD).

$$\begin{aligned} \text{Maximize } & f(u, v) - s(v | D') \\ & - u^T [\nabla_u f(u, v) + \nabla_{uu} f(u, v) q] \\ & - \frac{1}{2} q^T [\nabla_{uu} f(u, v) q] \end{aligned} \quad (81)$$



$$\begin{aligned} \text{Subject to } & \nabla_u f(u, v) + \nabla_{uu} f(u, v) q \geq 0, \\ & v \geq 0, \quad w \in C'. \end{aligned}$$

Conflict of Interests

The author declares that there is no conflict of interests regarding the publication of this paper.

References

- [1] V. Pareto, *Course D'économie Politique*, Lausanne, Raye, Saint Lucia, 1986.
- [2] H. W. Kuhn and A. W. Tucker, "Nonlinear programming," in *Proceedings of the 2nd Berkeley Symposium on Mathematical Statistics and Probability*, pp. 481–492, University of California Press, Berkeley, Calif, USA, 1951.
- [3] A. M. Geoffrion, "Proper efficiency and the theory of vector maximization," *Journal of Mathematical Analysis and Applications*, vol. 22, pp. 618–630, 1968.
- [4] W. S. Dorn, "A symmetric dual theorem for quadratic programs," *Journal of the Operations Research Society of Japan*, vol. 2, pp. 93–97, 1960.
- [5] G. B. Dantzig, E. Eisenberg, and R. W. Cottle, "Symmetric dual nonlinear programs," *Pacific Journal of Mathematics*, vol. 15, pp. 809–812, 1965.
- [6] B. Mond, "Second order duality for nonlinear programs," *Opsearch*, vol. 11, no. 2-3, pp. 90–99, 1974.
- [7] B. Mond and T. Weir, "Generalized concavity and duality," in *Generalized Concavity in Optimization and Economics*, S. Schaible and W. T. Ziemba, Eds., pp. 263–279, Academic Press, New York, NY, USA, 1981.
- [8] O. L. Mangasarian, "Second- and higher-order duality in nonlinear programming," *Journal of Mathematical Analysis and Applications*, vol. 51, no. 3, pp. 607–620, 1975.
- [9] C. R. Bector and S. Chandra, "Generalized bonvex function and second order duality in mathematical programming," Research Report 85-2, The University of Manitoba, Winnipeg, Canada, 1985.
- [10] G. Devi, "Symmetric duality for nonlinear programming problem involving η -bonvex functions," *European Journal of Operational Research*, vol. 104, no. 3, pp. 615–621, 1998.
- [11] S. Khurana, "Symmetric duality in multiobjective programming involving generalized cone-index functions," *European Journal of Operational Research*, vol. 165, no. 3, pp. 592–597, 2005.
- [12] X. M. Yang, X. Q. Yang, K. L. Teo, and S. H. Hou, "Second order symmetric duality in non-differentiable multiobjective programming with F -convexity," *European Journal of Operational Research*, vol. 164, no. 2, pp. 406–416, 2005.
- [13] X. M. Yang, X. Q. Yang, K. L. Teo, and S. H. Hou, "Multiobjective second-order symmetric duality with F -convexity," *European Journal of Operational Research*, vol. 165, no. 3, pp. 585–591, 2005.
- [14] T. R. Gulati, S. K. Gupta, and I. Ahmad, "Second-order symmetric duality with cone constraints," *Journal of Computational and Applied Mathematics*, vol. 220, no. 1-2, pp. 347–354, 2008.
- [15] T. R. Gulati and Geeta, "Mond-Weir type second-order symmetric duality in multiobjective programming over cones," *Applied Mathematics Letters*, vol. 23, no. 4, pp. 466–471, 2010.
- [16] T. R. Gulati and K. Verma, "Nondifferentiable multiobjective Wolfe type symmetric duality under invexity," in *Proceedings of the International Conference on Soft Computing for Problem Solving*, vol. 1, pp. 347–354, 2012.
- [17] S. K. Gupta and N. Kailey, "Nondifferentiable multiobjective second-order symmetric duality," *Optimization Letters*, vol. 5, no. 1, pp. 125–139, 2011.
- [18] S. K. Gupta and N. Kailey, "Second-order multiobjective symmetric duality involving cone-bonvex functions," *Journal of Global Optimization*, vol. 55, no. 1, pp. 125–140, 2013.
- [19] H. Saini and T. R. Gulati, "Nondifferentiable multiobjective symmetric duality with F -convexity over cones," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 74, no. 5, pp. 1577–1584, 2011.
- [20] M. S. Bazaraa and J. J. Goode, "On symmetric duality in nonlinear programming," *Operations Research*, vol. 21, pp. 1–9, 1973.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

