

Research Article

Some Combinatorial Interpretations and Applications of Fuss-Catalan Numbers

Chin-Hung Lin

Department of Mathematics, National Taiwan University, Taipei 10617, Taiwan

Correspondence should be addressed to Chin-Hung Lin, jlch3554@hotmail.com

Received 1 August 2011; Accepted 25 August 2011

Academic Editors: L. Ji and K. Ozawa

Copyright © 2011 Chin-Hung Lin. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Fuss-Catalan number is a family of generalized Catalan numbers. We begin by two definitions of Fuss-Catalan numbers and some basic properties. And we give some combinatorial interpretations different from original Catalan numbers. Finally we generalize the Jonah's theorem as its applications.

1. Introduction

Catalan numbers $\{c_n\}_{n \geq 0}$ [1] are said to be the sequence satisfying the recursive relation

$$c_{n+1} = c_0 c_n + c_1 c_{n-1} + \cdots + c_n c_0, \quad c_0 = 1. \quad (1.1)$$

It is well known that the n th term of Catalan numbers is $c_n = (1/(n+1))\binom{2n}{n} = (1/(2n+1))\binom{2n+1}{n}$ and $\{c_n\}_{n \geq 0} = \{1, 1, 2, 5, 14, 42, 132, \dots\}$. Also, one of many combinatorial interpretations of Catalan numbers is that c_n is the number of shortest lattice paths from $(0, 0)$ to (n, n) on the 2-dimensional plane such that those paths lie beneath the line $y = x$.

On the other hand, *Fuss-Catalan numbers* $\{c_n^{(s)}\}_{s, n \geq 0}$ were investigated by Fuss [2] and studied by several authors [1, 3–7]. Hence we have the following proposition.

Proposition 1.1. *If n and s are nonnegative integers, the following statements are equivalent:*

(1)

$$c_n^{(s)} = \frac{1}{sn+1} \binom{sn+1}{n}; \quad (1.2)$$

(2)

$$c_{n+1}^{(s)} = \sum_{r_1+r_2+\dots+r_s=n} c_{r_1}^{(s)} \times c_{r_2}^{(s)} \times \dots \times c_{r_s}^{(s)}, \quad c_0^{(s)} = 1; \quad (1.3)$$

(3) $c_n^{(s)}$ is the number of shortest lattice paths from $(0,0)$ to $(n, (s-1)n)$ on the 2-dimensional plane such that those paths lie beneath $y = (s-1)x$.

It is easy to see that in the case when $s = 2$, the sequence of Catalan numbers $\{c_n\}$ is a special case of the family of Fuss-Catalan numbers $\{c_n^{(2)}\}$. Although Fuss-Catalan numbers could be viewed as one kind of generalized Catalan numbers, Fuss finished this work many years before Catalan [8].

The proposition describing Fuss-Catalan numbers could be restated in the language of generating functions.

Proposition 1.2. The generating function $C^{(s)}(x) = \sum_{n \geq 0} c_n^{(s)} x^n$ satisfies the equation $x(A^{(s)})^s(x) = A^{(s)}(x) - 1$, where $A(x)$ is a generating function. That is,

$$x\left(C^{(s)}\right)^s(x) = C^{(s)}(x) - 1. \quad (1.4)$$

There are many combinatorial interpretations of Fuss-Catalan numbers, but most of them are similar to that of Catalan numbers. In order to demonstrate the importance of Fuss-Catalan number, in Section 2 we tried to find some combinatorial interpretations which is different from original Catalan numbers.

Finally, Hilton and Pedersen [9] generalized an identity called Jonah's theorem which involves Catalan numbers. So in Section 3 we restated the identity in Jonah's theorem in the form of Fuss-Catalan numbers.

2. Some Other Interpretations

It is remarkable that the interpretation in Proposition 1.1 illustrates the relation between paths in an $n \times n$ square and Catalan numbers. It is reasonable to consider whether Fuss-Catalan numbers are relevant to paths in an $n \times n \times n$ cube. As the cube in Figure 1, consider the shortest path in it from $(0,0,0)$ to (n,n,n) . There are $(3n)!/n!n!n!$ paths. But it is notable that $(3n)!/n!n!n!$ could be also written as

$$\binom{3n}{n} \binom{2n}{n} \binom{n}{n} = (2n+1)c_n^{(3)} \times (n+1)c_n^{(2)}. \quad (2.1)$$

Maybe by giving some constraints, shown in Figure 2, the number of paths will be precisely $c_n^{(3)} \times c_n^{(2)}$. And here are some results, which consider a more general case on an $n \times n \times (s-1)n$ cuboid.



Figure 1: An $n \times n \times n$ cube.

Figure 2: Constrained regions in Corollary 2.2 and Theorem 2.1.

Theorem 2.1. From $(0,0,0)$ to $(n, n, (s-1)n)$ and under the following constraints:

$$\begin{aligned} (s-1)y - z &\geq 0, \\ x - \frac{1}{s}y - \frac{1}{s}z &\geq 0, \end{aligned} \tag{2.2}$$

there are $c_n^{(s+1)} \times c_n^{(s)}$ shortest paths.

Proof. Let P be a shortest path constrained by the conditions in Theorem 2.1. First consider the projection of P on the yz -plane. The projection could be thought as a shortest path in an $n \times (s-1)n$ right triangle in Proposition 1.1. So there are $c_n^{(s)}$ ways to decide a path on this triangle. Fix one path and use this path to cut the cuboid with the positive direction of the x -axis. The graph is like a ladder in the right side of Figure 3 and could be put on a plane, and then it becomes an $n \times sn$ right triangle. So in this situation, there are $c_n^{(s+1)}$ ways to be chosen. Finally since we may choose the paths in the $n \times (s-1)n$ triangle and that in the $n \times sn$ triangle independently, there are $c_n^{(s+1)} \times c_n^{(s)}$ paths totally. \square




Figure 3: Auxiliary graphs for Theorem 2.1.

Corollary 2.2. From $(0,0,0)$ to (n,n,n) and under the following constraints:

$$\begin{aligned} y - z &\geq 0, \\ x - \frac{1}{2}y - \frac{1}{2}z &\geq 0, \end{aligned} \quad (2.3)$$

there are $c_n^{(3)} \times c_n^{(2)}$ shortest paths.

Proof. This is a special case of Theorem 2.1. □

If now we loosen the condition “the base of the cuboid must be square”, we can get some more general result. And the proof of this theorem is similar to that of Theorem 2.1.

Theorem 2.3. Let m, n, s, t be positive integers and $sn = (t-1)m$. From $(0,0,0)$ to $(m,n,(s-1)n)$ and under the following constraints:

$$\begin{aligned} (s-1)y - z &\geq 0, \\ x - \frac{m}{sn}y - \frac{m}{sn}z &\geq 0, \end{aligned} \quad (2.4)$$

there are $c_m^{(t)} \times c_n^{(s)}$ shortest paths.

3. Generalized Jonah's Theorem

Jonah's theorem [9] is the identity

$$\binom{n+1}{m} = \sum_{i \geq 0} c_i \binom{n-2i}{m-i}, \quad n \in \mathbb{N}_0, \quad m \in \mathbb{N}_0, \quad (3.1)$$

where \mathbb{N}_0 is the set of nonnegative integers and c_i is the i th term of Catalan numbers. Hilton and pedersen [9] proved the new identity

$$\binom{n+1}{m} = \sum_{i \geq 0} c_i \binom{n-2i}{m-i}, \quad n \in \mathbb{R}, \quad m \in \mathbb{N}_0, \quad (3.2)$$

where \mathbb{R} is the set of real numbers. The theorem is proven by lattice paths. Here we try to generalize the identity (3.2) as follows showing the connection with Fuss-Catalan numbers $c_n^{(s)}$:

$$\binom{n+1}{m} = \sum_{i \geq 0} c_i^{(s)} \binom{n-si}{m-i}, \quad n \in \mathbb{R}, \quad m \in \mathbb{N}_0. \quad (3.3)$$

The following lemmas will be needed to prove the identity (3.3).

Lemma 3.1. *For any generating function $f(x)$ with $f(0) = 0$, the equation*

$$f(x)B^s(x) = B(x) - 1 \quad (3.4)$$

has at most one solution of generating function (abbreviated SGF). That is, if $f(x)$ is a generating function with $f(0) = 0$, there is at most one generating function $g(x)$ satisfying

$$f(x)g^s(x) = g(x) - 1. \quad (3.5)$$

Proof. The cases $s = 0$ and $s = 1$ is easy since (3.4) could be solved immediately. So we assume that $s \geq 2$. If $g(x)$ is one SGF of (3.4), we have the identity

$$fB^s - B + 1 = (B - g)(fB^{s-1} + fgB^{s-2} + \cdots + fg^{s-2}B + fg^{s-1} - 1), \quad (3.6)$$

where f, B , and g are the abbreviations of $f(x)$, $B(x)$, and $g(x)$. Since the ring of formal power series is an integral domain, if $B(x)$ has any SGF other than $g(x)$ then the second term in the right hand side must be identically zero. However, if $B(x)$ is a generating function, we have $B(0) < +\infty$ and so $f(0)B^i(0) = f(0)g^{s-1}(0) = 0$ for all proper integer i . Since the second term on the right equals -1 but not 0 as $x = 0$, it could not be identically zero. That is, $g(x)$ is the only SGF. \square

Lemma 3.2. *Let*

$$\begin{aligned} p(x) &= 1 + x, \\ q(x) &= C^{(s)}(x(1+x)^{-s}) = \sum_{i \geq 0} c_i^{(s)} x^i (1+x)^{-is}, \end{aligned} \quad (3.7)$$

where $C^{(s)}$ is the generating function of $c_n^{(s)}$ where s is fixed. Then for $f(x) = x(1+x)^{-s}$, both $p(x)$ and $q(x)$ are SGFs of (3.4). Hence $p(x) = q(x)$. That is,

$$1 + x = \sum_{i \geq 0} c_i^{(s)} x^i (1+x)^{-is}. \quad (3.8)$$

Proof. Naturally $p(x)$ is a generating function; $q(x)$ is also a generating function since it is the linear combination of power of the generating function $f(x)$.

Observe that

$$\begin{aligned} (1) \quad & x(1+x)^{-s} p^s(x) = x(1+x)^{-s} (1+x)^s = x = p(x) - 1; \\ (2) \quad & x(1+x)^{-s} q^s(x) = x(1+x)^{-s} (C^{(s)}(x(1+x)^{-s}))^s = C^{(s)}(x(1+x)^{-s}) - 1 = q(x) - 1, \end{aligned}$$

since

$$x \left(C^{(s)}(x) \right)^s = C^{(s)}(x) - 1 \quad (3.9)$$

by Proposition 1.2. So both $p(x)$ and $q(x)$ are SGFs of (3.4). Finally by Lemma 3.1, there is at most one SGF. Hence $p(x) = q(x)$. \square

Theorem 3.3. For any real number n and integer m , the following identity holds:

$$\binom{n+1}{m} = \sum_{i \geq 0} c_i^{(s)} \binom{n-si}{m-i}, \quad n \in \mathbb{R}, \quad m \in \mathbb{N}_0. \quad (3.10)$$

Proof. By multiplying both sides of (3.8) by $(1+x)^n$, we have

$$(1+x)^{n+1} = c_0^{(s)} (1+x)^n + c_1^{(s)} x(1+x)^{n-s} + \cdots + c_m^{(s)} x^m (1+x)^{n-ms} + \cdots. \quad (3.11)$$

Then we get (3.10) by comparing the coefficients. \square

The following are the special cases of Theorem 3.3:

(i) $s = 0 \Rightarrow$ Pascal's theorem

$$\binom{n+1}{m} = \binom{n}{m} + \binom{n}{m-1}, \quad (3.12)$$

(ii) $s = 1 \Rightarrow$ Chu Shih-Chieh's theorem

$$\binom{n+1}{m} = \binom{n}{m} + \binom{n-1}{m-1} + \cdots + \binom{n-m}{0}. \quad (3.13)$$

(iii) $s = 2 \Rightarrow$ Jonah's theorem.

On the other hand, even when $m \geq n+1$ the identity holds.

Example 3.4. Recall that $\binom{a}{a} = 1$ and $\binom{a}{b} = 0$ when $b > a$.

(i) When $s = 3$, $n = 4$, $m = 5$,

$$\begin{aligned} 1 &= \binom{5}{5} = c_0^{(3)} \binom{4}{5} + c_1^{(3)} \binom{1}{4} + c_2^{(3)} \binom{-2}{3} + c_3^{(3)} \binom{-5}{2} + c_4^{(3)} \binom{-8}{1} + c_5^{(3)} \binom{-11}{0} \\ &= 1 \times 0 + 1 \times 0 + 3 \times (-4) + 12 \times 15 + 55 \times (-8) + 273 \times 1 = 1. \end{aligned} \quad (3.14)$$




Figure 4: The lattice for the proof of Theorem 3.3.

(ii) When $s = 3$, $n = 3$, $m = 5$,

$$\begin{aligned}
 0 &= \binom{4}{5} = c_0^{(3)} \binom{3}{5} + c_1^{(3)} \binom{0}{4} + c_2^{(3)} \binom{-3}{3} + c_3^{(3)} \binom{-6}{2} + c_4^{(3)} \binom{-9}{1} + c_5^{(3)} \binom{-12}{0} \\
 &= 1 \times 0 + 1 \times 0 + 3 \times (-10) + 12 \times 21 + 55 \times (-9) + 273 \times 1 = 0.
 \end{aligned} \tag{3.15}$$

Note 1. Theorem 3.3 can also be proved by lattice paths when n is nonnegative integer and $n - m + 1 \geq (s - 1)m$ (see Figure 4).

Consider the number of shortest path from $(0, 0)$ to $(m, n - m + 1)$, which is $\binom{n+1}{m}$. On the other hand, consider the auxiliary line $L : y = (s - 1)x$. Then every path must pass through L in order to reach the ending point $(m, n - m + 1)$. So we can classify all the paths by the points they pass L for the “first time”. Then there are $c_i^{(s)} \binom{n-si}{m-i}$ paths passing by point $(i, (s - 1)i)$, because before $(i, (s - 1)i)$ the path lies beneath L , and thus there are $c_i^{(s)}$ ways; after $(i, (s - 1)i)$ the path must go upward to $(i, (s - 1)i + 1)$ and then finally reach $(m, n - m + 1)$ without any constraints, and thus there are $\binom{n-si}{m-i}$ ways. So the total number of paths will be the summation of that of each point.

Acknowledgments

This paper was written in the Summer Research Program of Taiwan Academia Sinica in 2009. The author would like to thank Taiwan Academia Sinica for providing the financial support. He would like to thank Professor Peter Shiue very much for several useful ideas, suggestions, and comments to improve this paper. He would also like to thank Professor Sen-Peng Eu for encouragements and giving advices.

References

- [1] R. P. Stanley, *Enumerative Combinatorics*, vol. 2, Cambridge University Press, Cambridge, UK, 1999.
- [2] N. I. Fuss, "Solutio quæstionis, quot modis polygonum n laterum in polygona m laterum, per diagonales resolvi quæat," *Nova Acta Academiae Scientiarum Imperialis Petropolitanae*, vol. 9, pp. 243–251, 1791.
- [3] W. Chu, "A new combinatorial interpretation for generalized Catalan number," *Discrete Mathematics*, vol. 65, no. 1, pp. 91–94, 1987.
- [4] R. L. Graham, D. E. Knuth, and O. Patashnik, *Concrete Mathematics*, Addison-Wesley Publishing Company, Reading, Mass, USA, 1989.
- [5] M. Konvalinka, "Divisibility of generalized Catalan numbers," *Journal of Combinatorial Theory A*, vol. 114, no. 6, pp. 1089–1100, 2007.
- [6] B. E. Sagan, "Proper partitions of a polygon and k -Catalan numbers," *Ars Combinatoria*, vol. 88, pp. 109–124, 2008.
- [7] A. D. Sands, "On generalised catalan numbers," *Discrete Mathematics*, vol. 21, no. 2, pp. 219–221, 1978.
- [8] E. Catalan, "Not sur une équation aux différences finies," *Journal de Mathématiques Pures et Appliquées*, vol. 3, pp. 508–516, 1838.
- [9] P. Hilton and J. Pedersen, "The ballot problem and Catalan numbers," *Nieuw Archief voor Wiskunde*, vol. 8, no. 2, pp. 209–216, 1990.

