

Research Article

The Fundamental Groups of m -Quasi-Einstein Manifolds

Hee Kwon Lee

School of Mathematics, KIAS, 207-43 Cheongnyangni 2-dong, Dongdaemun-gu, Seoul 130-722, Republic of Korea

Correspondence should be addressed to Hee Kwon Lee, heekwon@kias.re.kr

Received 21 November 2011; Accepted 8 December 2011

Academic Editors: A. Borowiec and A. M. Cegarra

Copyright © 2011 Hee Kwon Lee. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

In Ricci flow theory, the topology of Ricci soliton is important. We call a metric quasi-Einstein if the m -Bakry-Emery Ricci tensor is a constant multiple of the metric tensor. This is a generalization of gradient shrinking Ricci soliton. In this paper, we will prove the finiteness of the fundamental group of m -quasi-Einstein with a positive constant multiple.

1. Introduction and Main Results

Ricci flow is introduced in 1982 and developed by Hamilton (cf. [1]):

$$\begin{aligned}\frac{\partial}{\partial t}g &= -2\text{Ric}, \\ g(0) &= g_0.\end{aligned}\tag{1.1}$$

Recently, Perelman supplemented Hamilton's result and solved the Poincaré Conjecture and the Geometrization Conjecture by using a Ricci flow theory. But in higher dimension greater than 4 classification using Ricci flow is still far-off. Most above all the classification of Ricci solitons, which are singularity models, is not completed. But there exist many properties of Ricci solitons. Here we say g is a Ricci soliton if (M, g) is a Riemannian manifold such that the identity

$$\text{Ric} + L_X g = cg\tag{1.2}$$

holds for some constant c and some complete vector field X on M . If $c > 0$, $c = 0$, or $c < 0$, then we call it shrinking, steady, or expanding. Moreover, if the vector field X appearing in (1.2) is the gradient field of a potential function $(1/2)f$, one has $\text{Ric} + \nabla \nabla f = cg$ and says g is a gradient Ricci soliton. In 2008, L6pez and R6o have shown that if (M, g) is a complete manifold with $\text{Ric} + L_X g \geq cg$ and some positive constant c , then M is compact if and only if $\|X\|$ is bounded. Moreover, under these assumptions if M is compact, then $\pi_1(M)$ is finite. Furthermore, Wylie [2] has shown that under these conditions if M is complete, then $\pi_1(M)$ is finite. Moreover, in 2008, Fang et al. (cf. [3]) have shown that a gradient shrinking Ricci soliton with a bounded scalar curvature has finite topological type. By [4, Proposition 1.5.6], Cao and Zhu have shown that compact steady or expanding Ricci solitons are Einstein manifolds. In addition by [4, Corollary 1.5.9 (ii)] note that compact shrinking Ricci solitons are gradient Ricci solitons. So we are interested in shrinking gradient Ricci solitons. In [6, page 354], Eminenti et al. have shown that compact shrinking Ricci solitons have positive scalar curvatures. In [6] Case et al. have shown that an m -quasi-Einstein with $1 \leq m < \infty$ and $c > 0$ has a positive scalar curvature. Let me introduce the definition of m -quasi-Einstein.

Definition 1.1. The triple (M, g, f) is an m -quasi-Einstein manifold if it satisfies the equation

$$\text{Ric} + \text{Hess} f - \frac{1}{m} df \otimes df = cg \quad (1.3)$$

for some $c \in \mathbb{R}$.

Here m -Bakry-Emery Ricci tensor $\text{Ric}_f^m \doteq \text{Ric} + \text{Hess} f - (1/m)df \otimes df$ for $0 < m \leq \infty$ is a natural extension of the Ricci tensor to smooth metric measure spaces (cf. [6, Section 1]). Note that if $m = \infty$, then it reduces to a gradient Ricci soliton. In this paper, we will prove the finiteness of the fundamental group of an m -quasi-Einstein with $c > 0$.

Theorem 1.2. Let (M, g, f) be a complete manifold with $c > 0$ and $\text{Ric} + \text{Hess} f - (1/m)df \otimes df \geq cg$. Then it has a finite fundamental group.

2. The Proof of Theorem 1.2

The proof of Theorem 1.2 is similar to the proofs of [2, 7].

Proof. We will prove it by dividing into two cases.

Case 1. $\|\nabla f\|$ is bounded. We claim that the bounded $\|\nabla f\|$ implies the compactness of M . Let q be a point in M , and consider any geodesic $\gamma : [0, \infty) \rightarrow M$ emanating from q and parametrized by arc length t . Then we have

$$\int_0^T \text{Ric}(\dot{\gamma}, \dot{\gamma}) \geq cT + \frac{1}{m} \int_0^T (df(\dot{\gamma}))^2 - \int_0^T \dot{\gamma}(g(\nabla f, \dot{\gamma})) \geq cT - g(\nabla f, \dot{\gamma})|_0^T. \quad (2.1)$$

Since $g(\nabla f, \dot{\gamma})|_0^T$ is bounded we have that $\int_0^\infty \text{Ric}(\dot{\gamma}, \dot{\gamma}) = \infty$. Hence, the claim is followed by the proof of [4, Theorem 1]. Let (\tilde{M}, \tilde{g}) be the Riemannian universal cover of (M, g) , let $p : (\tilde{M}, \tilde{g}) \rightarrow (M, g)$ be a projection map, and let \tilde{f} be a map $f \circ p$. Since p is a local isometry, then the same inequality holds, that is, $\text{Ric}(\tilde{g}) + \text{Hesse}_{\tilde{g}} \tilde{f} - (1/m)d\tilde{f} \otimes d\tilde{f} \geq c\tilde{g}$. Now, since

$\|\tilde{\nabla} \tilde{f}\|$ is bounded, it is followed from the above argument that \tilde{M} is compact. So $\pi_1(M)$ is finite.

Case 2. $\|\nabla f\|$ is unbounded. We will prove this case by following the proof of [2]. By Case 1, M is noncompact. For any $p \in M$, define

$$H_p \doteq \max\{0, \sup\{\text{Ric}_y(v, v) : y \in B(p, 1), \|v\| = 1\}\}. \quad (2.2)$$

Note that by [7, Lemma 2.2] we have

$$\int_0^r \text{Ric}(\dot{\gamma}, \dot{\gamma}) ds \leq 2(n-1) + H_p + H_q. \quad (2.3)$$

Assume that $d(p, q) > 1$. On the other hand, from the inequality of Theorem 1.2, we have

$$\begin{aligned} \int_0^r \text{Ric}(\dot{\gamma}, \dot{\gamma}) ds &\geq cd(p, q) + \frac{1}{m} \int_0^r (df(\dot{\gamma}))^2 - \int_0^r \dot{\gamma}(g(\nabla f, \dot{\gamma})) \\ &\geq cd(p, q) - \|\nabla f\|_p - \|\nabla f\|_q, \end{aligned} \quad (2.4)$$

since $g(\nabla f, \dot{\gamma}) \leq \|\nabla f\| \|\dot{\gamma}\|$. Hence, we have that for any $p, q \in M$

$$d(p, q) \leq \max\left\{1, \frac{1}{c} \left(2(n-1) + H_p + H_q + \|\nabla f\|_p + \|\nabla f\|_q\right)\right\}. \quad (2.5)$$

Now we will apply a similar argument like Case 1. Fix $\tilde{p} \in \tilde{M}$, and let $h \in \pi_1(M)$ identified as a deck transformation on \tilde{M} . Note that $B(\tilde{p}, 1)$ and $B(h(\tilde{p}), 1)$ are isometric, and thus $H_{\tilde{p}} = H_{h(\tilde{p})}$. Also $\|\tilde{\nabla} \tilde{f}\|_{\tilde{p}} = \|\tilde{\nabla} \tilde{f}\|_{h(\tilde{p})}$. So we conclude that

$$d(\tilde{p}, h(\tilde{p})) \leq \max\left\{1, \frac{2}{c} \left(n-1 + H_{\tilde{p}} + \|\tilde{\nabla} \tilde{f}\|_{\tilde{p}}\right)\right\} \quad (2.6)$$

for any $h \in \pi_1(M)$. Since the right-hand side is independent of h , this proves this case. \square

References

- [1] R. S. Hamilton, "Three-manifolds with positive Ricci curvature," *Journal of Differential Geometry*, vol. 17, no. 2, pp. 255–306, 1982.
- [2] W. Wylie, "Complete shrinking Ricci solitons have finite fundamental group," *Proceedings of the American Mathematical Society*, vol. 136, no. 5, pp. 1803–1806, 2008.
- [3] F.-Q. Fang, J.-W. Man, and Z.-L. Zhang, "Complete gradient shrinking Ricci solitons have finite topological type," *Comptes Rendus Mathématique*, vol. 346, no. 11-12, pp. 653–656, 2008.
- [4] H.-D. Cao and X.-P. Zhu, "A complete proof of the Poincaré and geometrization conjectures—application of the Hamilton-Perelman theory of the Ricci flow," *Asian Journal of Mathematics*, vol. 10, no. 2, pp. 165–492, 2006.
- [5] M. Eminenti, G. La Nave, and C. Mantegazza, "Ricci solitons: the equation point of view," *Manuscripta Mathematica*, vol. 127, no. 3, pp. 345–367, 2008.

- [6] J. Case, Y.-J. Shu, and G. Wei, "Rigidity of quasi-Einstein metrics," *Differential Geometry and its Applications*, vol. 29, no. 1, pp. 93–100, 2011.
- [7] M. Fernández-López and E. García-Río, "A remark on compact Ricci solitons," *Mathematische Annalen*, vol. 340, no. 4, pp. 893–896, 2008.

