

Review Article

Attitudes and Beliefs towards Patients with Hazardous Alcohol Use: A Systematic Review

Neelam Mabood,¹ Hansen Zhou,^{1,2} Kathryn A. Dong,³ Samina Ali,¹
T. Cameron Wild,^{4,5} and Amanda S. Newton^{1,4,6}

¹ Department of Pediatrics, Faculty of Medicine and Dentistry, University of Alberta, Edmonton, AB, Canada T6G 1C9

² Department of Psychology, University of Alberta, Edmonton, AB, Canada T6G 2E9

³ Department of Emergency Medicine, Faculty of Medicine and Dentistry, University of Alberta, Edmonton, AB, Canada T6G 2T4

⁴ Department of Psychiatry, Faculty of Medicine and Dentistry, University of Alberta, Edmonton, AB, Canada T6G 2R7

⁵ School of Public Health, University of Alberta, Edmonton, AB, Canada T6G 1C9

⁶ Department of Pediatrics, Edmonton Clinic Health Academy (ECHA), 11405-87 Avenue, Edmonton, AB, Canada T6G 1C9

Correspondence should be addressed to Amanda S. Newton, mandi.newton@ualberta.ca

Received 20 September 2012; Accepted 4 November 2012

Academic Editors: C. Arreola-Risa, C. Damsa, and W. Ryotaro

Copyright © 2012 Neelam Mabood et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Objective. To describe emergency department (ED) staff attitudes and beliefs towards patients presenting with hazardous alcohol use and their clinical management. **Methods.** A search of MEDLINE, EMBASE, CINAHL, SCOPUS from 1990 to 2010, and reference lists from included studies was conducted. Two reviewers independently screened for inclusion and assessed study quality. One reviewer extracted the data and a second checked for completeness and accuracy. **Results.** Among nine studies four reported varied beliefs on whether screening was worthwhile for identifying hazardous alcohol use (physicians: 42%–88%; nurses: 50%–100%). Physicians in three studies were divided on intervention provision (32%–54% in support of intervention provision) as were nurses in two studies (39% and 64% nurses in support of intervention provision). Referral for treatment was identified in two studies as an important part of ED management (physicians: 62% and 97%; nurses: 95%). Other attitudes and beliefs identified across the studies included concern that asking about alcohol consumption would be seen as obtrusive or offensive, and a perceived lack of time and resources available for providing care and referrals. **Conclusions.** ED staff had varying attitudes towards ED management of patients with hazardous alcohol use. Investigations into improving clinical care for hazardous alcohol use are needed to optimize ED management for these patients.

1. Introduction

Hazardous alcohol use is well known to increase an individual's risk of injury due to violence or accidents [1, 2]. Acute treatment and care for alcohol-associated morbidities are often sought in hospital emergency departments (EDs), with ED patients more likely than primary care patients and the general population to report hazardous alcohol use [3–7]. In 2007, almost 3 million ED visits across the USA (2.3% of all ED visits) were primarily related to alcohol and cost the health care system \$1.3 billion dollars [8].

The high volume of alcohol-related presentations to the ED presents an opportunity for staff to detect hazardous

alcohol use, optimize care decisions, and initiate preventative interventions [4]. Patient management approaches using the SBIRT model (Screening, Brief Intervention, Referral to Treatment) have been evaluated in the ED with mixed impact [9, 10]. The SBIRT model uses a public health approach to universal screening for substance use problems. It can rule out nonproblem substance users and identify the level of risk, and who will benefit from a brief advice. The SBIRT model can be used as a stand-alone treatment or combined with other treatment [11, 12].

A large number of studies have demonstrated short-term effects on reducing alcohol use or associated harms and cost-effectiveness for adult patients; however, several studies have

also reported statistically nonsignificant effects [13–35]. The current position of the U.S. Preventive Services Task Force is that there is at least fair evidence to support the use of screening and brief intervention (BI) for those patients who screen positive for hazardous alcohol consumption [36]. In 2006, the American College of Surgeons mandated alcohol screening and intervention for trauma patients admitted to Levels 1 and 2 trauma centers [37]. Despite this, recent evidence suggests that ED-based SBIRT is largely absent in these settings [38]. This absence may be due, in part, to attitudes and beliefs of ED staff [4, 39]. Translational research for ED-based SBIRT with specific attention to barriers to successful implementation including attitudes and beliefs has been recommended [39]. The objective of this systematic review was to describe ED nurses' and physicians' attitudes and beliefs towards patients with hazardous alcohol use and their clinical management.

2. Methods

2.1. Search Strategies. Guided by input from the research team, a research librarian developed and implemented a systematic search strategy using language (English) and year (1990 to 2010) restrictions. The search was conducted in February 2010 and updated in June 2010. We used the EBSCOhost portal, encompassing the MEDLINE, EMBASE, CINAHL, and Scopus databases to conduct the search. We also searched our keywords in Google Scholar and the reference lists of retrieved studies. In our search we used the following key words and MeSH headings: “alcohol,” “attitudes OR beliefs,” “emergency department,” “Alcohol-Related Disorders,” “Attitudes of Health Personnel,” “Emergency Service or Emergency Medicine,” and “Physician-Patient Relations.”

2.2. Study Selection. Two reviewers independently screened the search results (NM, HZ). The full manuscripts of potentially relevant studies were retrieved if they were identified as relevant by at least one of the reviewers, and then independently confirmed for inclusion by two reviewers (NM, HZ). The same reviewers also independently assessed study inclusion/exclusion. Studies were included at the screening and inclusion/exclusion stages if a primary objective was to determine attitudes and beliefs of ED staff (physicians and nurses) towards patients presenting with hazardous alcohol use or their clinical management in the ED. No restrictions were placed on study design (qualitative or quantitative) or patient age. Studies were excluded if they met any of the following criteria: (1) they were conducted in any language besides English, (2) they studied a non-ED setting, (3) the study of attitudes and beliefs was not a primary objective, or (4) they examined attitudes and beliefs towards polysubstance use or the hazardous use of substances other than alcohol.

2.3. Assessment of Quality. Two reviewers (NM, HZ) assessed study quality. Disagreements were resolved with third party discussion (ASN) until mutual agreement was achieved. The

quality assessment of quantitative studies depended on study design. Observational studies were assessed using questions adapted from Guyatt, Sackett, and Cook's (1993) User Guide to Medical Literature by the Critical Appraisal Skills Program [40]. The questions targeted the study's focus, methods, biases, and results presentation including practicality and applicability. Qualitative studies were assessed using a tool developed for the study by the research team (available from the corresponding author upon request). The tool evaluated studies based on the following domains: validity of the study design, setting, and sampling; informed consent and appropriately addressed ethical issues; methodological reporting; efforts to establish credibility and validity; the dependability and reliability of study data.

2.4. Data Extraction and Synthesis. Data from the final set of studies were extracted using a standardized form that assessed key study characteristics (e.g., year of publication, country), characteristics of the study population and setting, and results specific to ED staff attitudes and beliefs. Data were extracted by one reviewer (NM) and checked for accuracy and completeness by a second reviewer (HZ). Discrepancies were resolved by consensus. Published data and tests of significance reported by study authors were included. In the case of unclear or unreported information in the original studies, primary authors were contacted. Meta-analyses were not conducted due to heterogeneity of study definitions and measurement. A qualitative analysis was conducted, and detailed findings are presented in evidence tables. Results are presented by design: controlled trial, observational studies, and mixed method and qualitative studies.

3. Results

3.1. Description of Included Studies. Figure 1 describes the flow of studies through the selection process. The search strategies identified 352 studies as potentially relevant. After title and abstract review, 37 papers were selected for manuscript retrieval and full review, with 9 studies meeting our inclusion criteria after full-text review: seven observational studies, one qualitative, and one mixed method study [41–49]. General characteristics of the studies are summarized in Table 1. These studies, published between 1998 and 2009, were conducted in the United States ($n = 2$), United Kingdom ($n = 2$), Sweden ($n = 2$), Scotland ($n = 1$), Australia ($n = 1$), and China ($n = 1$). The majority of studies sampled physicians only ($n = 4$), but studies reporting information from both physicians and nurses ($n = 3$) or nurses only ($n = 2$) were also included in the review. Study objectives were similar between studies; they either examined ED health care professionals' attitudes and beliefs towards adult patients with presentations for hazardous alcohol use or towards the SBIRT model.

3.2. Methodological Quality

3.2.1. Observational Studies. The seven observational studies met requirements for many of the critical components

TABLE 1: Study characteristics.

First author (country, year)	Study design	Sample	Participants	
			Gender (% F, M)	Response rate (%)
D'Onofrio (USA, 2002)	Nonrandomized controlled trial	<i>N</i> = 36 Physicians [‡] : 100%	28% F, 72% M	100
Indig (Australia, 2009)	Cross-sectional survey	<i>N</i> = 78 Nurses: 54%; physicians: 46%	Physicians: 44% F, 56% M Nurses: 80% F, 20% M	30
O'Rourke (USA, 2006)	Cross-sectional survey	<i>N</i> = 598 Staff physicians: 66%; residents [‡] : 34%	32% F, 68% M	17
Huntley [§] (UK, 2004)	Cross-sectional survey	<i>N</i> = 127 Physicians: 100%	NS	100
Chung (China, 2003)	Cross-sectional survey	<i>N</i> = 190 Nurses: 100%	84% F, 16% M	32
Anderson (Scotland, 2001)	Cross-sectional survey	<i>N</i> = 96 Physicians: 36%; nurses: 63%	Physicians: 9% F, 91% M Nurses: 83% F, 17% M	57
Graham (USA, 2000)	Cross-sectional survey	<i>N</i> = 257 Physicians: 100%	19% F, 81% M	46
Waller (UK, 1998)	Cross-sectional survey	<i>N</i> = 367 Physicians: 46%; nurses: 54%	Physicians: 15% F, 85% M Nurses: 78% F, 22% M	82
Karlsson (Sweden, 2005)	Mixed method	Interview, <i>N</i> = 9; questionnaire, <i>N</i> = 72 Nurses: 100%	NS	75
Nordqvist (Sweden, 2005)	Qualitative	<i>N</i> = 6 Physicians: 100%	34% F, 66% M	100

[‡] Medical residents; NS: not specified; [§]Huntley also reported qualitative results from a separate study; M: male; F: female.




FIGURE 1: Selection of studies.

identified by the Critical Appraisal Skills Programme [40]. Four studies were limited by a lack of power calculation [41, 43, 46, 47]. Two studies were limited by volunteer bias resultant from convenience sampling [46, 47]. Study strengths included providing a clear focus and presenting results with adequate data analyses [41, 43, 46, 47]. Four studies also had clear discussions addressing the research/clinical implications of the findings with references to other studies [41, 43, 46, 47]. Study applicability was addressed by two studies [46, 47]. Three studies were prone to methodological weaknesses such as volunteer bias resultant from convenience sampling [43, 44]. Three studies did not provide a sample size calculation [42, 44, 45]. There was no description of analytical procedures in one study [42]. One study did not use validated measures [45]. Across all observational studies, many had limited applicability to North American emergency care settings, as they were conducted in European and Asian settings utilizing different health care systems and models [41, 44, 45, 47].

3.2.2. Mixed Method and Qualitative Studies. Both the mixed method and qualitative studies in this paper were clear in research focus, but did not provide sufficient explanation to justify the specific qualitative methodology used [48, 49]. However, sampling strategies and data collection methodology were carried out appropriately in both studies. The studies did not describe informed consent procedures or the issue of participant anonymity. Use of multiple data sources and data triangulation enhanced the credibility and validity of one study's findings [48].

3.3. Attitudes and Beliefs of ED Staff towards SBIRT. Figures 2 to 4 present the attitudes and beliefs of ED staff towards SBIRT for patients presenting with hazardous alcohol use. ED staff attitudes towards screening varied (Figure 2). Across two studies, from 42% to 88% of physicians and from 50% to 100% of nurses believed that it was worthwhile to identify hazardous alcohol use in the ED [41, 47]. One study reported that 60% of surveyed physicians and residents believed screening for hazardous alcohol use would improve treatment success [46]. Beliefs regarding who was responsible for screening differed. Huntley et al. reported almost all of surveyed physicians believed that it was their responsibility to screen for alcohol use (98%) while Indig et al. reported only 50% of physicians and 35% of nurses believed that they were responsible for screening [45, 47]. Support for BI for hazardous alcohol use was variable amongst ED staff (Figure 3). Both the Anderson and Waller studies reported that 65% of physicians and 71–75% of nurses believed it was worthwhile to perform BI [41, 43]. Two studies reported less support (51% and 54%) by physicians in the use of BI [42, 46]. A minority of ED staff in Indig et al.'s study believed they were responsible for providing BI (32% of physicians and 39% of nurses) [47]. Indig also reported that 62% of physicians and 95% of nurses believed it was worthwhile to make referrals for specialist treatment (also reported by Huntley et al.), although only 53% of physicians and 51%

of nurses believed they were responsible for referring to a specialist (Figure 4) [45, 47].

3.4. Attitudes and Beliefs Related to the Treatability of Hazardous Alcohol Use. The majority of ED staff felt that something could be done in the ED setting to assist patients with hazardous alcohol use. Only a minority of ED staff in the Waller study believed that the ED setting could do little to assist patients with hazardous alcohol use (33% of physicians, 17% of nurses), a belief also held by a similar percentage of physicians (15%) and very few nurses (2%) in Indig's study [41, 47]. Attitudes and beliefs related to the treatability of hazardous alcohol use were conflicting across the studies. O'Rourke et al. reported that 75% of surveyed physicians believed that alcohol use disorders were treatable, although 80% of those surveyed also believed that current treatments did not work [46]. Graham et al. also reported similar conflicting attitudes and beliefs with 77% of surveyed physicians who agreed that alcohol use disorders were treatable, but also believed that they were difficult to treat (97%) [42]. In contrast, almost all physicians (97%) in the Huntley study believed that treatment could be successful [45].

3.5. Attitudes and Beliefs regarding the Health Care Provider-Patient Relationship. Several studies identified reluctance on the part of ED health care providers to engage in treatment for hazardous alcohol use. Anderson et al. reported that 35% of surveyed physicians and 53% of nurses were hesitant to ask patients about alcohol consumption and believed formal training was required to be able to respond to patients with hazardous alcohol use (53% of physicians, 78% of nurses) [43]. Physicians (46%) and nurses (48%) in Waller's study reported reluctance to question or interact with such patients [41]. Among nurses, Chung et al. reported mixed attitudes towards patients with hazardous alcohol use (mean = 106, SD 13; minimum/maximum score of 27/189 indicating negative/positive attitude) [44]. The belief that patients with hazardous alcohol use lacked motivation to change was cited by physicians (12% and 88%) and nurses (15% and 85%) in two studies [43, 47]. In a qualitative study, Nordqvist et al. found that physicians did not trust the reliability of patients' responses and believed that asking about alcohol use could make patients feel guilty. Physicians in this study also believed that there was little chance of patients reducing their drinking as a result of their interaction with medical staff and did not believe that BI was effective [49]. In Karlsson's study, 11% of nurses believed BI would negatively impact the relationship with patients and that patients would object to participating [48]. Nurses also believed that the subject of alcohol consumption was too sensitive to be discussed in a brief ED visit, with 61% of nurses believing that patients would respond negatively to questions about alcohol use. These concerns also surfaced in other studies. Indig's study found that 24% of physicians and 39% of nurses believed questions about alcohol use were offensive while Andersen et al. reported that 56% of physicians and 90% of nurses surveyed believed that patients found questions about




FIGURE 2: Attitudes and beliefs towards screening for hazardous alcohol use in the ED.

alcohol use offensive and intrusive [43, 47]. Nurses (72%) and physicians (61%) in Waller's study also held similar beliefs [41].

4. Discussion

This paper identified significant variation in ED physicians' and nurses' attitudes and beliefs towards patients with hazardous alcohol use and their management, which may help explain variations in SBIRT model use in the ED. Findings highlight the need to address key issues that underpinned the attitudes and beliefs: perceived time constraints and a lack of resources by ED, concerns that patients would respond negatively to SBIRT, communicating emerging evidence on the SBIRT model to ED staff, and identifying and responding to ED staff learning needs.

The studies in this paper identified that while ED physicians and nurses believed it is worthwhile to screen for hazardous alcohol use, provide BI, and refer for further treatment, far fewer believed they were professionally responsible for these aspects of clinical care and management. The low reports of ED-based SBIRT may be related, in part, to barriers cited across studies in this paper including a perceived lack of time to provide BI, a lack of resources for SBIRT implementation (including specialist staff and support services), and the patient's intoxicated state.

The relationship of these barriers to attitudes and beliefs, however, was not examined in the studies. The issues of who is prepared and best able to conduct SBIRT in the ED, and whether SBIRT is appropriate for this clinical setting remain debated. ED-based studies have employed different strategies for SBIRT delivery in the ED including training ED staff (physician, nurse/nurse practitioner, social worker, and emergency medical technician) and health promotion advocates to augment ED staff roles [20, 50]. Another study used designated mental health nurses to conduct post-ED, follow-up appointments [51]. Time constraints and workload concerns voiced by physicians and nurses may be addressed by introducing specialized SBIRT care providers available for ED or post-ED care. ED-based SBIRT providers may address concerns raised by ED staff about patient reactions to SBIRT while post-ED care can address intoxication as a barrier to ED delivery.

The low reports of ED-based SBIRT as a clinical responsibility identified by this paper could also be the result of providers' beliefs that screening would not improve treatment success (believed by 40% of physicians in one study), current treatments do not work (believed by 80% of physicians in one study), or that hazardous alcohol use/substance use disorders are difficult to treat (believed by 97% of physicians one study). According to Nordqvist et al., physicians need to be assured that performing BI is effective and worth the time [49]. Establishing effective




FIGURE 3: Attitudes and beliefs towards brief intervention (BI) for hazardous alcohol use in the ED.

and timely mechanisms to communicate the growing body of evidence supporting ED-based screening and BI to ED staff may be necessary to address the negative beliefs cited in a large number of studies. Financially compensating physicians for the time it takes to perform SBIRT in the ED, providing adequate supports (personnel, easy access to treatment and referral mechanisms), as well as personalized audit and feedback may also help to improve physician adherence with current recommendations. Further, as new studies address existing methodological limitations (e.g., standardizing outcome measures, establishing effectiveness among patient subgroups such as sex and age) and clinical gaps (e.g., practical screening tools and interventions that complement the busy and hectic nature of an ED) establishing mechanisms to inform ED staff of new evidence in this field will keep staff appraised as to whether SBIRT is not only an effective model, but a feasible model for the ED [39].

A number of issues related to adequate training were cited in the studies. ED staff in two studies cited training as a needed facilitator for patient care and management, and a high percentage of staff in another study identified

a lack of confidence in performing screening (50% of physicians and 71% of nurses), BI (71% of physicians and 73% of nurses), and referral (44% of physicians and 59% of nurses) [41, 43, 47]. Several studies have shown an increase in screening and BI delivery in primary and general care settings and in the ED following a variety of educational and training modalities [52–56]. Confidence has also improved for some staff, but not all [55, 56]. Significant changes have not been reported for attitudes and beliefs and readiness to change clinical behaviors [55]. In-person training has been suggested as more effective than web-based resources while educational support and training are suggested to be more impactful with staff who enter training already therapeutically committed to working with patients with hazardous alcohol consumption [57, 58]. Widespread adoption of the SBIRT model in EDs will require more than a small number of studies demonstrating the effects of training programs. An extensive structured literature review conducted by Williams et al. looked at implementation programs of nine countries having geographically diverse clinical settings and research infrastructures [59]. The paper




FIGURE 4: Attitudes and beliefs towards referral to treatment for hazardous alcohol use in the ED.

found different implementation programs and looked at their success/failure in implementing robust SBIRT strategies [59]. This paper, and similar papers indicate the need for change in the policies related to SBIRT in the ED and related clinical settings [20, 39]. Policies and guidelines set at the institutional level and the presence of faculty/administrators that promote and train for the usage of the SBIRT model may also warrant further attention as do studies focusing on strategies to increase the feasibility of SBIRT model use in the ED vis-à-vis cost analyses evaluating the short-term and long-term costs and benefits of SBIRT implementation [39].

5. Limitations of the Study

This review has several limitations. We assessed the methodological quality of studies based on published methods and did not contact corresponding authors to verify the methods used. As a result, some studies may have been adequately conducted, but the methods were poorly reported. We also did not include studies that explored attitudes and beliefs as a minor objective. As a result, two studies were screened but excluded from our review, and others may have been missed in our search strategy [55, 56]. We chose not to include studies that did not have a primary focus of attitudes and beliefs because the depth and breadth of the investigation may have differed from those studies with it as a primary objective. No pediatric studies were identified in this review. Given that underage drinking is widespread and ample evidence exists that underage hazardous alcohol use leads to adult substance use disorders and persistent dysfunction, the role of the ED in addressing hazardous alcohol use by adolescents could play a critical role in identifying those youth who could benefit from treatment and the initial management of hazardous alcohol use through BI and referral [60–71].

There were also limitations in this review that stemmed from the included studies themselves. The majority of the studies in this review were weak to moderate in quality. Studies that reduce the biases and methodological weaknesses observed in the current body of literature are

needed. This includes eliminating biases from convenience or selective sampling through randomized sampling procedures or population sampling and reducing the likelihood of volunteer bias through low response rates (6 out of 9 studies had response rates <80%). Future qualitative and mixed methods studies require justification for the chosen methodology to demonstrate that they study answered the research questions and evidence needs to be provided that the researcher took steps to ensure that the conclusions reached are dependable and confirmable. It would be helpful to explore the variations in attitudes and beliefs and create an opportunity to evaluate whether they are substantial barriers to the SBIRT model in the ED.

6. Conclusion

Detection of hazardous alcohol consumption followed by BI in the ED setting has large potential benefits due to the wide population that can be captured and a growing body of empirical evidence favoring ED-based SBIRT. This review suggests that attitudes and beliefs of ED physicians and nurses may be key barriers to the widespread uptake of SBIRT for hazardous alcohol use in the ED setting.

Disclosure

Dr. A. S. Newton holds a New Investigator Award from the Canadian Institutes of Health Research (CIHR). Dr. T. C. Wild is a Health Scholar with Alberta Innovates-Health Solutions.



References

- [1] C. J. Cherpitel, Y. Ye, and J. Bond, "Alcohol and injury: multi-level analysis from the Emergency Room Collaborative Alcohol Analysis Project (ERCAAP)," *Alcohol and Alcoholism*, vol. 39, no. 6, pp. 552–558, 2004.
- [2] C. J. Cherpitel, "Alcohol and injuries: a review of international emergency room studies since 1995," *Drug and Alcohol Review*, vol. 26, no. 2, pp. 201–214, 2007.

- [3] G. D'Onofrio and L. C. Degutis, "Preventive care in the emergency department: screening and brief intervention for alcohol problems in the emergency department: a systematic review," *Academic Emergency Medicine*, vol. 9, no. 6, pp. 627–638, 2002.
- [4] G. D'Onofrio and L. C. Degutis, "Screening and brief intervention in the emergency department," *Alcohol Research and Health*, vol. 28, no. 2, pp. 63–72, 2004.
- [5] A. J. McDonald, N. Wang, and C. A. Camargo, "US Emergency Department Visits for Alcohol-Related Diseases and Injuries between 1992 and 2000," *Archives of Internal Medicine*, vol. 164, no. 5, pp. 531–537, 2004.
- [6] P. Nilsen, M. Holmqvist, C. Nordqvist, and P. Bendtsen, "Frequency of heavy episodic drinking among nonfatal injury patients attending an emergency room," *Accident Analysis and Prevention*, vol. 39, no. 4, pp. 757–766, 2007.
- [7] C. J. Cherpitel and Y. Ye, "Drug use and problem drinking associated with primary care and emergency room utilization in the US general population: data from the 2005 national alcohol survey," *Drug and Alcohol Dependence*, vol. 97, no. 3, pp. 226–230, 2008.
- [8] L. M. Wier, K. Levit, E. Stranges et al., *HCUP Facts and Figures: Statistics on Hospital-Based Care in the United States*, Agency for Healthcare Research and Quality, 2008, <http://www.hcup-us.ahrq.gov/reports.jsp>.
- [9] J. W. Bray, A. J. Cowell, and J. M. Hinde, "A systematic review and meta-analysis of health care utilization outcomes in alcohol screening and brief intervention trials," *Medical Care*, vol. 49, no. 3, pp. 287–294, 2011.
- [10] A. Havard, A. Shakeshaft, and R. Sanson-Fisher, "Systematic review and meta-analyses of strategies targeting alcohol problems in emergency departments: interventions reduce alcohol-related injuries," *Addiction*, vol. 103, no. 3, pp. 368–376, 2008.
- [11] T. F. Babor, B. G. McRee, P. A. Kassebaum, P. L. Grimaldi, K. Ahmed, and J. Bray, "Screening, Brief Intervention, and Referral to Treatment (SBIRT): toward a public health approach to the management of substance abuse," *Substance Abuse*, vol. 28, no. 3, pp. 7–30, 2007.
- [12] S. Sterling, T. Valkanoff, and A. Hinman, "Integrating substance use treatment into adolescent health care," *Current Psychiatry Reports*, vol. 14, pp. 453–461, 2012.
- [13] S. Bazargan-Hejazi, E. Bing, M. Bazargan et al., "Evaluation of a brief intervention in an inner-city emergency department," *Annals of Emergency Medicine*, vol. 46, no. 1, pp. 67–76, 2005.
- [14] E. Bernstein, J. Bernstein, and S. Levenson, "Project ASSERT: an ED-based intervention to increase access to primary care, preventive services, and the substance abuse treatment system," *Annals of Emergency Medicine*, vol. 30, no. 2, pp. 182–189, 1997.
- [15] F. C. Blow, K. L. Barry, M. A. Walton et al., "The efficacy of two brief intervention strategies among injured, at-risk drinkers in the emergency department: impact of tailored messaging and brief advice," *Journal of Studies on Alcohol*, vol. 67, no. 4, pp. 568–578, 2006.
- [16] R. H. Aseltine, "The impact of screening, brief intervention and referral for treatment in emergency department patients' alcohol use: a 3-, 6-and 12-month follow-up," *Alcohol and Alcoholism*, vol. 45, no. 6, Article ID agq058, pp. 514–519, 2010.
- [17] F. E. Vaca, D. Winn, C. L. Anderson, D. Kim, and M. Arcila, "Six-month follow-up of computerized alcohol screening, brief intervention, and referral to treatment in the emergency department," *Substance Abuse*, vol. 32, no. 3, pp. 144–152, 2011.
- [18] M. J. Mello, R. Longabaugh, J. Baird, T. Nirenberg, and R. Woolard, "DIAL: a telephone brief intervention for high-risk alcohol use with injured emergency department patients," *Annals of Emergency Medicine*, vol. 51, no. 6, pp. 755–764, 2008.
- [19] R. Longabaugh, R. F. Woolard, T. D. Nirenberg et al., "Evaluating the effects of a brief motivational intervention for injured drinkers in the emergency department," *Journal of Studies on Alcohol*, vol. 62, no. 6, pp. 806–816, 2001.
- [20] Academic ED SBIRT Research Collaborative, "The impact of screening, brief intervention, and referral for treatment on emergency department patients' alcohol use," *Annals of Emergency Medicine*, vol. 50, no. 6, pp. 699–710.e6, 2007.
- [21] A. Spirito, P. M. Monti, N. P. Barnett et al., "A randomized clinical trial of a brief motivational intervention for alcohol-positive adolescents treated in an emergency department," *Journal of Pediatrics*, vol. 145, no. 3, pp. 396–402, 2004.
- [22] P. M. Monti, S. M. Colby, N. P. Barnett et al., "Brief intervention for harm reduction with alcohol-positive older adolescents in a hospital emergency department," *Journal of Consulting and Clinical Psychology*, vol. 67, no. 6, pp. 989–994, 1999.
- [23] M. A. Walton, S. T. Chermack, J. T. Shope et al., "Effects of a brief intervention for reducing violence and alcohol misuse among adolescents: a randomized controlled trial," *JAMA*, vol. 304, no. 5, pp. 527–535, 2010.
- [24] P. M. Désy, P. K. Howard, C. Perhats, and S. Li, "Alcohol screening, brief intervention, and referral to treatment conducted by emergency nurses: an impact evaluation," *Journal of Emergency Nursing*, vol. 36, no. 6, pp. 538–545, 2010.
- [25] S. Estee, T. Wickizer, L. He, M. F. Shah, and D. Mancuso, "Evaluation of the washington state screening, brief intervention, and referral to treatment project: cost outcomes for medicaid patients screened in hospital emergency departments," *Medical Care*, vol. 48, no. 1, pp. 18–24, 2010.
- [26] J.-B. Daeppen, J. Gaume, P. Bady et al., "Brief alcohol intervention and alcohol assessment do not influence alcohol use in injured patients treated in the emergency department: a randomized controlled clinical trial," *Addiction*, vol. 102, no. 8, pp. 1224–1233, 2007.
- [27] G. D'Onofrio, M. V. Pantalon, L. C. Degutis et al., "Brief intervention for hazardous and harmful drinkers in the emergency department," *Annals of Emergency Medicine*, vol. 51, no. 6, pp. 742–750.e2, 2008.
- [28] C. J. Cherpitel, J. Moskalewicz, G. Swiatkiewicz, Y. Ye, and J. Bond, "Screening, Brief Intervention, and Referral to Treatment (SBIRT) in a polish emergency department: three-month outcomes of a randomized, controlled clinical trial," *Journal of Studies on Alcohol and Drugs*, vol. 70, no. 6, pp. 982–990, 2009.
- [29] C. J. Cherpitel, R. A. Korcha, J. Moskalewicz, G. Swiatkiewicz, Y. Ye, and J. Bond, "Screening, brief intervention, and referral to treatment (SBIRT): 12-month outcomes of a randomized controlled clinical trial in a polish emergency department," *Alcoholism*, vol. 34, no. 11, pp. 1922–1928, 2010.
- [30] A. Spirito, H. Sindelar-Manning, S. M. Colby et al., "Individual and family motivational interventions for alcohol-positive adolescents treated in an emergency department: results of a randomized clinical trial," *Archives of Pediatrics and Adolescent Medicine*, vol. 165, no. 3, pp. 269–274, 2011.

- [31] J. Bernstein, T. Heeren, E. Edward et al., "A brief motivational interview in a pediatric emergency department, plus 10-day telephone follow-up, increases attempts to quit drinking among youth and young adults who screen positive for problematic drinking," *Academic Emergency Medicine*, vol. 17, no. 8, pp. 890–902, 2010.
- [32] R. F. Maio, J. T. Shope, F. C. Blow et al., "A randomized controlled trial of an emergency department-based interactive computer program to prevent alcohol misuse among injured adolescents," *Annals of Emergency Medicine*, vol. 45, no. 4, pp. 420–429, 2005.
- [33] R. J. Tait, G. K. Hulse, and S. I. Robertson, "Effectiveness of a brief-intervention and continuity of care in enhancing attendance for treatment by adolescent substance users," *Drug and Alcohol Dependence*, vol. 74, no. 3, pp. 289–296, 2004.
- [34] B. D. Johnston, F. P. Rivara, R. M. Drosch, C. Dunn, and M. K. Copass, "Behavior change counseling in the emergency department to reduce injury risk: a randomized, controlled trial," *Pediatrics*, vol. 110, no. 2, pp. 267–274, 2002.
- [35] R. J. Tait, G. K. Hulse, S. I. Robertson, and P. C. Sprivulis, "Emergency department-based intervention with adolescent substance users: 12-Month outcomes," *Drug and Alcohol Dependence*, vol. 79, no. 3, pp. 359–363, 2005.
- [36] U.S. Preventive Services Task Force, "Screening and behavioral counseling interventions in primary care to reduce alcohol misuse: recommendation statement," *Annals of Internal Medicine*, vol. 140, no. 7, pp. 554–556, 2004.
- [37] American College of Surgeons, *Resources for Optimal Care for the Injured Patient*, American College of Surgeons, Chicago, Ill, USA, 2006.
- [38] R. M. Cunningham, S. R. Harrison, M. P. McKay et al., "National survey of emergency department alcohol screening and intervention practices," *Annals of Emergency Medicine*, vol. 55, no. 6, pp. 556–562, 2010.
- [39] R. M. Cunningham, S. L. Bernstein, M. Walton et al., "Alcohol, tobacco, and other drugs: future directions for screening and intervention in the emergency department," *Academic Emergency Medicine*, vol. 16, no. 11, pp. 1078–1088, 2009.
- [40] Solutions for Public Health, *Critical Appraisal Skills Programme Making Sense of the Evidence: 12 Questions to Help You Make Sense of a Cohort Study*, Solutions for Public Health, 2011, http://www.sph.nhs.uk/acl_users/credentials.cookie_auth/require_login?came_from=http%3A//www.sph.nhs.uk/sph-files/casp-appraisal-tools/cohort%252012%2520questions.pdf.
- [41] S. Waller, B. Thom, S. Harris, and M. Kelly, "Perceptions of alcohol-related attendances in accident and emergency departments in England: a national survey," *Alcohol and Alcoholism*, vol. 33, no. 4, pp. 354–361, 1998.
- [42] D. M. Graham, R. F. Maio, F. C. Blow, and E. M. Hill, "Emergency physician attitudes concerning intervention for alcohol abuse/dependence delivered in the emergency department: a brief report," *Journal of Addictive Diseases*, vol. 19, no. 1, pp. 45–53, 2000.
- [43] S. Anderson, D. R. Eadie, A. M. MacKintosh, and S. Haw, "Management of alcohol misuse in Scotland: the role of A&E nurses," *Accident and Emergency Nursing*, vol. 9, no. 2, pp. 92–100, 2001.
- [44] J. Y. M. Chung, J. T. S. Chan, R. S. D. Yeung, R. C. H. Wan, and S. T. Ho, "Nurses' attitude toward alcoholic patients in accident and emergency department in Hong Kong," *Hong Kong College of Emergency Medicine*, vol. 10, no. 2, pp. 104–112, 2003.
- [45] J. S. Huntley, R. Patton, and R. Touquet, "Attitudes towards alcohol of emergency department doctors trained in the detection of alcohol misuse," *Annals of the Royal College of Surgeons of England*, vol. 86, no. 5, pp. 329–333, 2004.
- [46] M. O'Rourke, L. D. Richardson, I. Wilets, and G. D'Onofrio, "Alcohol-related problems: emergency physicians' current practice and attitudes," *Journal of Emergency Medicine*, vol. 30, no. 3, pp. 263–268, 2006.
- [47] D. Indig, J. Copeland, K. M. Conigrave, and I. Rotenko, "Attitudes and beliefs of emergency department staff regarding alcohol-related presentations," *International Emergency Nursing*, vol. 17, no. 1, pp. 23–30, 2009.
- [48] A. Karlsson, K. Johansson, C. Nordqvist, and P. Bendtsen, "Feasibility of a computerized alcohol screening and personalized written advice in the ED: opportunities and obstacles," *Accident and Emergency Nursing*, vol. 13, no. 1, pp. 44–53, 2005.
- [49] C. Nordqvist, K. Johansson, K. Lindqvist, and P. Bendtsen, "Alcohol prevention measures at an emergency department: physicians' perspectives," *Public Health*, vol. 119, no. 9, pp. 789–791, 2005.
- [50] G. D'Onofrio and L. C. Degutis, "Integrating Project assert: a screening, intervention, and Referral to treatment program for unhealthy alcohol and drug use into an urban emergency department," *Academic Emergency Medicine*, vol. 17, no. 8, pp. 903–911, 2010.
- [51] M. J. Crawford, R. Patton, R. Touquet et al., "Screening and referral for brief intervention of alcohol-misusing patients in an emergency department: a pragmatic randomised controlled trial," *The Lancet*, vol. 364, no. 9442, pp. 1334–1339, 2004.
- [52] M. Fleming, D. Olsen, H. Stathes et al., "Virtual reality skills training for health care professionals in alcohol screening and brief intervention," *Journal of the American Board of Family Medicine*, vol. 22, no. 4, pp. 387–398, 2009.
- [53] J. E. Hettema, J. L. Sorensen, M. Uy, and S. Jain, "Motivational enhancement therapy to increase resident physician engagement in substance abuse education," *Substance Abuse*, vol. 30, no. 3, pp. 244–247, 2009.
- [54] E. F. S. Kaner, C. A. Lock, B. R. McAvoy, N. Heather, and E. Gilvarry, "A RCT of three training and support strategies to encourage implementation of screening and brief alcohol intervention by general practitioners," *British Journal of General Practice*, vol. 49, no. 446, pp. 699–703, 1999.
- [55] G. D'Onofrio, E. S. Nadel, L. C. Degutis et al., "Improving emergency medicine residents' approach to patients with alcohol problems: a controlled educational trial," *Annals of Emergency Medicine*, vol. 40, no. 1, pp. 50–62, 2002.
- [56] E. Bernstein, J. Bernstein, J. Feldman et al., "An evidence based alcohol screening, brief intervention and referral to treatment (SBIRT) curriculum for emergency department (ED) providers improves skills and utilization," *Substance Abuse*, vol. 28, no. 4, pp. 79–92, 2007.
- [57] D. P. Alford, J. M. Richardson, S. E. Chapman, C. E. Dubé, R. W. Schadt, and R. Saitz, "A web-based Alcohol Clinical Training (ACT) curriculum: is in-person faculty development necessary to affect teaching?" *BMC Medical Education*, vol. 8, article 11, 2008.
- [58] P. Anderson, E. Kaner, S. Wutzke et al., "Attitudes and managing alcohol problems in general practice: an interaction analysis based on findings from who collaborative study," *Alcohol and Alcoholism*, vol. 39, no. 4, pp. 351–356, 2004.

- [59] E. C. Williams, M. L. Johnson, G. T. Lapham et al., "Strategies to implement alcohol screening and brief intervention in primary care settings: a structured literature review," *Psychology of Addictive Behaviors*, vol. 25, no. 2, pp. 206–214, 2011.
- [60] United States Department of Health, *The NSDUH Report: Quantity and Frequency of Alcohol Use Among Underage Drinkers*, Substance Abuse and Mental Health Services Administration, Office of Applied Studies, 2010, <http://oas.samhsa.gov/2k8/underage/underage.cfm>.
- [61] L. D. Johnston, P. M. O'Malley, J. G. Bachman, and J. E. Schulenberg, *Monitoring the Future: National Survey Results on Drug Use, 1975–2004: Volume I. Secondary School Students*, National Institute on Drug Abuse, Bethesda, Md, USA, 2005.
- [62] Y. A. Bonomo, G. Bowes, C. Coffey, J. B. Carlin, and G. C. Patton, "Teenage drinking and the onset of alcohol dependence: a cohort study over seven years," *Addiction*, vol. 99, no. 12, pp. 1520–1528, 2004.
- [63] R. M. Viner and B. Taylor, "Adult outcomes of binge drinking in adolescence: findings from a UK national birth cohort," *Journal of Epidemiology and Community Health*, vol. 61, no. 10, pp. 902–907, 2007.
- [64] E. J. D'Amico, P. L. Ellickson, R. L. Collins, S. Martino, and D. J. Klein, "Processes linking adolescent problems to substance-use problems in late young adulthood," *Journal of Studies on Alcohol*, vol. 66, no. 6, pp. 766–775, 2005.
- [65] P. L. Ellickson, J. S. Tucker, and D. J. Klein, "Ten-year prospective study of public health problems associated with early drinking," *Pediatrics*, vol. 111, no. 5, part 1, pp. 949–955, 2003.
- [66] B. F. Grant and D. A. Dawson, "Age at onset of alcohol use and its association with DSM-IV alcohol abuse and dependence: results from the national longitudinal alcohol epidemiologic survey," *Journal of Substance Abuse*, vol. 9, no. 1, pp. 103–110, 1997.
- [67] D. J. DeWit, E. M. Adlaf, D. R. Offord, and A. C. Ogborne, "Age at first alcohol use: a risk factor for the development of alcohol disorders," *American Journal of Psychiatry*, vol. 157, no. 5, pp. 745–750, 2000.
- [68] R. A. Zucker, J. E. Donovan, A. S. Masten, M. E. Mattson, and H. B. Moss, "Early developmental processes and the continuity of risk for underage drinking and problem drinking," *Pediatrics*, vol. 121, no. 4, pp. S252–S272, 2008.
- [69] P. Rohde, P. M. Lewinsohn, J. R. Seeley, D. N. Klein, J. A. Andrews, and J. W. Small, "Psychosocial functioning of adults who experienced substance use disorders as adolescents," *Psychology of Addictive Behaviors*, vol. 21, no. 2, pp. 155–164, 2007.
- [70] B. F. Grant, F. S. Stinson, and T. C. Harford, "Age at onset of alcohol use and DSM-IV alcohol abuse and dependence: a 12-year follow-up," *Journal of Substance Abuse*, vol. 13, no. 4, pp. 493–504, 2001.
- [71] T. Pitkänen, "Problem drinking and psychological well-being: a five-year follow-up study from adolescence to young adulthood," *Scandinavian Journal of Psychology*, vol. 40, no. 3, pp. 197–207, 1999.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

