STRONG ASYMPTOTICS FOR L_p EXTREMAL POLYNOMIALS OFF A COMPLEX CURVE

RABAH KHALDI

Received 29 September 2003 and in revised form 15 June 2004

We study the asymptotic behavior of $L_p(\sigma)$ extremal polynomials with respect to a measure of the form $\sigma = \alpha + \gamma$, where α is a measure concentrated on a rectifiable Jordan curve in the complex plane and γ is a discrete measure concentrated on an infinite number of mass points.

1. Introduction

Let F be a compact subset of the complex plane $\mathbb{C}$ and let B be a metric space of functions defined on F . We suppose that B contains the set of monic polynomials. Then the extremal or general Chebyshev polynomial T_n of degree n is a monic polynomial that minimizes the distance between zero and the set of all monic polynomials of degree n , that is,

$$\text{dist}(T_n, 0) = \min \{ \text{dist}(Q_n, 0) : Q_n(z) = z^n + a_{n-1}z^{n-1} + \cdots + a_0 \} = m_n(B). \quad (1.1)$$

Recently, a series of results concerning the asymptotic of the extremal polynomials was established for the case of $B = L_p(F, \sigma)$, $1 \leq p \leq \infty$, where σ is a Borel measure on F ; see, for example, [3, 7, 8, 12]. When $p = 2$, we have the special case of orthogonal polynomials with respect to the measure σ . A lot of research work has been done on this subject; see, for example, [1, 4, 5, 9, 11, 13]. The case of the spaces $L_p(F, \sigma)$, where $0 < p < \infty$ and F is a closed rectifiable Jordan curve with some smoothness conditions, was studied by Geronimus [2]. An extension of Geronimus's result has been given by Kaliaguine [3] who found asymptotics when $0 < p < \infty$ and the measure σ has a decomposition of the form

$$\sigma = \alpha + \gamma, \quad (1.2)$$

where α is a measure supported on a closed rectifiable Jordan curve E as defined in [2] and γ is a discrete measure with a finite number of mass points.

In this paper, we generalize Kaliaguine's work [3] in the case where $1 \leq p < \infty$ and the support of the measure σ is a rectifiable Jordan curve E plus an infinite discrete set of

mass points which accumulate on E . More precisely, $\sigma = \alpha + \gamma$, where the measure α and its support E are defined as in [3], that is,

$$d\alpha(\xi) = \rho(\xi)|d\xi|, \quad \rho \geq 0, \rho \in L^1(E, |d\xi|); \quad (1.3)$$

γ is a discrete measure concentrated on $\{z_k\}_{k=1}^\infty \subset \text{Ext}(E)$ ($\text{Ext}(E)$ is the exterior of E), that is,

$$\gamma = \sum_{k=1}^{+\infty} A_k \delta(z - z_k), \quad A_k > 0, \sum_{k=1}^{+\infty} A_k < \infty. \quad (1.4)$$

Note that the result of the special case $p = 2$ is also a generalization of [4]. More precisely, in the proof of Theorem 4.3, we show that condition [4, page 265, (17)] imposed on the points $\{z_k\}_{k=1}^\infty$ is redundant.

2. The $H^p(\Omega, \rho)$ spaces ($1 \leq p < \infty$)

Let E be a rectifiable Jordan curve in the complex plane, $\Omega = \text{Ext}(E)$, $G = \{z \in \mathbb{C}, |z| > 1\}$ (∞ belongs to Ω and G).

We denote by Φ the conformal mapping of Ω into G with $\Phi(\infty) = \infty$ and $1/C(E) = \lim_{z \rightarrow \infty} (\Phi(z)/z) > 0$, where $C(E)$ is the logarithmic capacity of E . We denote $\Psi = \Phi^{-1}$.

Let ρ be an integrable nonnegative weight function on E satisfying the Szegő condition

$$\int_E (\log \rho(\xi)) |\Phi'(\xi)| |d\xi| > -\infty. \quad (2.1)$$

Condition (2.1) allows us to construct the so-called Szegő function D associated with the curve E and the weight function ρ :

$$D(z) = \exp \left\{ -\frac{1}{2p\pi} \int_{-\pi}^{+\pi} \frac{w + e^{it}}{w - e^{it}} \log \left(\frac{\rho(\xi)}{|\Phi'(\xi)|} \right) dt \right\} \quad (w = \Phi(z), \xi = \Psi(e^{it})) \quad (2.2)$$

such that

- (i) D is analytic in Ω , $D(z) \neq 0$ in Ω , and $D(\infty) > 0$;
- (ii) $|D(\xi)|^{-p} |\Phi'(\xi)| = \rho(\xi)$ a.e. on E , where $D(\xi) = \lim_{z \rightarrow \xi} D(z)$.

We say that $f \in H^p(\Omega, \rho)$ if and only if f is analytic in Ω and $f_0 \Psi / D_0 \Psi \in H^p(G)$.

For $1 \leq p < \infty$, $H^p(\Omega, \rho)$ is a Banach space. Each function $f \in H^p(\Omega, \rho)$ has limit values a.e. on E and

$$\|f\|_{H^p(\Omega, \rho)}^p = \int_E |f(\xi)|^p \rho(\xi) |d\xi| = \lim_{R \rightarrow 1^+} \frac{1}{R} \int_{E_R} \frac{|f(z)|^p}{|D(z)|^p} |\Phi'(z) dz|, \quad (2.3)$$

where $E_R = \{z \in \Omega : |\Phi(z)| = R\}$.

LEMMA 2.1 [3]. *If $f \in H^p(\Omega, \rho)$, then for every compact set $K \subset \Omega$, there is a constant C_K such that*

$$\sup \{ |f(z)| : z \in K \} \leq C_K \|f\|_{H^p(\Omega, \rho)}. \quad (2.4)$$

3. The extremal problems

Let $1 \leq p < \infty$; we denote $\sigma_l = \alpha + \sum_{k=1}^l A_k \delta(z - z_k)$ and by $\mu(\rho)$, $\mu(l)$, $\mu^\infty(\rho)$, $m_{n,p}(\rho)$, $m_{n,p}(l)$, and $m_{n,p}(\sigma)$ the extremal values of the following problems, respectively:

$$\mu(\rho) = \inf \{ \|\varphi\|_{H^p(\Omega, \rho)}^p : \varphi \in H^p(\Omega, \rho), \varphi(\infty) = 1 \}, \quad (3.1)$$

$$\mu(l) = \inf \{ \|\varphi\|_{H^p(\Omega, \rho)}^p : \varphi \in H^p(\Omega, \rho), \varphi(\infty) = 1, \varphi(z_k) = 0, k = 1, 2, \dots, l \}, \quad (3.2)$$

$$\mu^\infty(\rho) = \inf \{ \|\varphi\|_{H^p(\Omega, \rho)}^p : \varphi \in H^p(\Omega, \rho), \varphi(\infty) = 1, \varphi(z_k) = 0, k = 1, 2, \dots \}, \quad (3.3)$$

$$m_{n,p}(\rho) = \min \{ \|Q_n\|_{L_p(\alpha)} : Q_n(z) = z^n + \dots \}, \quad (3.4)$$

$$m_{n,p}(l) = \min \{ \|Q_n\|_{L_p(\sigma_l)} : Q_n(z) = z^n + \dots \}, \quad (3.5)$$

$$m_{n,p}(\sigma) = \min \{ \|Q_n\|_{L_p(\sigma)} : Q_n(z) = z^n + \dots \}. \quad (3.6)$$

As usual,

$$\|f\|_{L_p(\sigma)} := \left(\int_E |f(\xi)|^p d\sigma(\xi) \right)^{1/p}. \quad (3.7)$$

We denote by φ^* and ψ^∞ the extremal functions of problems (3.1) and (3.3), respectively.

Let $T_{n,p}^l(z)$ and $T_{n,p}(z)$ be the extremal polynomials with respect to the measures σ_l and σ , respectively, that is,

$$\|T_{n,p}^l\|_{L_p(\sigma_l)} = m_{n,p}(l), \quad \|T_{n,p}\|_{L_p(\sigma)} = m_{n,p}(\sigma). \quad (3.8)$$

LEMMA 3.1. *Let $\varphi \in H^p(\Omega, \rho)$ such that $\varphi(\infty) = 1$ and $\varphi(z_k) = 0$ for $k = 1, 2, \dots$, and let*

$$B_\infty(z) = \prod_{k=1}^{+\infty} \frac{\Phi(z) - \Phi(z_k)}{\Phi(z)\Phi(z_k) - 1} \frac{|\Phi(z_k)|^2}{\Phi(z_k)} \quad (3.9)$$

be the Blaschke product. Then

- (i) $B_\infty \in H^p(\Omega, \rho)$, $B_\infty(\infty) = 1$, $|B_\infty(\xi)| = \prod_{k=1}^{+\infty} |\Phi(z_k)|$ ($\xi \in E$);
- (ii) $\varphi/B_\infty \in H^p(\Omega, \rho)$ and $(\varphi/B_\infty)(\infty) = 1$.

Proof. This lemma is proved for $p = 2$ in [1]. The proof is based on the fact that if $f \in H^2(U)$, where $U = \{z \in \mathbb{C}, |z| < 1\}$, and B is the Blaschke product formed by the zeros of f , then $f/B \in H^2(U)$. It remains true in $H^p(U)$ for $1 \leq p < \infty$; see [6, 10]. $\square$

LEMMA 3.2. An extremal function ψ^∞ of problem (3.3) is given by $\psi^\infty = \varphi^* B_\infty$; in addition,

$$\mu^\infty(\rho) = \prod_{k=1}^{+\infty} (|\Phi(z_k)|)^p \mu(\rho). \quad (3.10)$$

Proof. If $\varphi \in H^p(\Omega, \rho)$, $\varphi(\infty) = 1$ and $\varphi(z_k) = 0$ for $k = 1, 2, \dots$. Then by Lemma 2.1, we have $f = \varphi/B_\infty \in H^p(\Omega, \rho)$, $f(\infty) = 1$, and $|B_\infty(\xi)| = \prod_{k=1}^{+\infty} |\Phi(z_k)|$ for $\xi \in E$. These lead to

$$\|f\|^p = \left(\prod_{k=1}^{+\infty} |\Phi(z_k)| \right)^{-p} \|\varphi\|^p. \quad (3.11)$$

Thus

$$\mu(\rho) \leq \left(\prod_{k=1}^{+\infty} |\Phi(z_k)| \right)^{-p} \mu^\infty(\rho). \quad (3.12)$$

On the other hand, since the function $\psi^\infty = \varphi^* B_\infty \in H^p(\Omega, \rho)$, $\varphi(\infty) = 1$ and $\varphi(z_k) = 0$ for $k = 1, 2, \dots$, we get

$$\mu^\infty(\rho) \leq \|\psi^\infty\|^p = \left(\prod_{k=1}^{+\infty} |\Phi(z_k)| \right)^p \mu(\rho). \quad (3.13)$$

Finally, the lemma follows from (3.12) and (3.13). $\square$

4. The main results

Definition 4.1. A measure $\sigma = \alpha + \gamma$ is said to belong to a class A if the absolutely continuous part α and the discrete part γ satisfy conditions (1.3), (1.4), and (2.1) and Blaschke's condition, that is,

$$\sum_{k=1}^{+\infty} (|\Phi(z_k)| - 1) < \infty. \quad (4.1)$$

We denote $\lambda_n = \Phi^n - \Phi_n$, where Φ_n is the polynomial part of the Laurent expansion of Φ^n in the neighborhood of infinity.

Definition 4.2 [2]. A rectifiable curve E is said to be of class Γ if $\lambda_n(\xi) \rightarrow 0$ uniformly on E .

THEOREM 4.3. Let a measure $\sigma = \alpha + \gamma$ satisfy conditions (1.3), (1.4) and Blaschke's condition (4.1); then

$$\lim_{l \rightarrow +\infty} m_{n,p}(l) = m_{n,p}(\sigma). \quad (4.2)$$

Proof. The extremal property of $T_{n,p}(z_k)$ gives

$$\begin{aligned} (m_{n,p}(\sigma))^p &\leq \int_E |T_{n,p}^l(\xi)|^p \rho(\xi) |d\xi| + \sum_{k=1}^l A_k |T_{n,p}^l(z_k)|^p + \sum_{k=l+1}^{+\infty} A_k |T_{n,p}^l(z_k)|^p \\ &= (m_{n,p}(l))^p + \sum_{k=l+1}^{+\infty} A_k |T_{n,p}^l(z_k)|^p. \end{aligned} \quad (4.3)$$

On the other hand, from the extremal property of $T_{n,p}^l(z_k)$, we can write

$$\begin{aligned} m_{n,p}(l) &\leq \left(\int_E |T_{n,p}(\xi)|^p \rho(\xi) |d\xi| + \sum_{k=1}^l A_k |T_{n,p}(z_k)|^p \right)^{1/p} \\ &\leq m_{n,p}(\sigma) = C_n < \infty. \end{aligned} \quad (4.4)$$

Note that C_n does not depend on l ; so for all $l = 1, 2, 3, \dots$,

$$\left(\int_E |T_{n,p}^l(\xi)|^p \rho(\xi) |d\xi| \right)^{1/p} < C_n. \quad (4.5)$$

This implies that there is a constant C'_n independent of l such that for all $l = 1, 2, 3, \dots$,

$$\max \{ |T_{n,p}^l(z)|^p : |z| \leq 2 \} < C'_n. \quad (4.6)$$

Using (4.6) in (4.3) for large enough l with (4.4), we get

$$(m_{n,p}(l))^p \leq (m_{n,p}(\sigma))^p \leq (m_{n,p}(l))^p + C'_n \sum_{k=l+1}^{+\infty} A_k. \quad (4.7)$$

Letting $l \rightarrow \infty$, we obtain

$$\lim_{l \rightarrow \infty} m_{n,p}(l) = m_{n,p}(\sigma). \quad (4.8)$$

□

THEOREM 4.4. Let $1 \leq p < \infty$, $E \in \Gamma$, and let $\sigma = \alpha + \gamma$ be a measure which belongs to A . In addition, for all n and l ,

$$m_{n,p}(l) \leq \left(\prod_{k=1}^l |\Phi(z_k)| \right) m_{n,p}(\rho). \quad (4.9)$$

Then the monic orthogonal polynomials $T_{n,p}(z)$ with respect to the measure σ have the following asymptotic behavior:

- (i) $\lim_{n \rightarrow \infty} (m_{n,p}(\sigma)/(C(E))^n) = (\mu^\infty(\rho))^{1/p}$;
- (ii) $\lim_{n \rightarrow \infty} \|T_{n,p}/[C(E)\Phi]^n - \psi^\infty\|_{H^p(\Omega,\rho)} = 0$;
- (iii) $T_{n,p}(z) = [C(E)\Phi(z)]^n [\psi^\infty(z) + \varepsilon_n(z)]$,

where $\varepsilon_n(z) \rightarrow 0$ uniformly on compact subsets of Ω and ψ^∞ is an extremal function of problem (3.3).

Remark 4.5. For $p = 2$ and E the unit circle, condition (4.9) is proved (see [5, Theorem 5.2]). In this case, this condition can be written as $\gamma_n/\gamma_n^l \leq \prod_{k=1}^l |z_k|$, where $\gamma_n^l = 1/m_{n,2}(l)$ and $\gamma_n = 1/m_{n,2}(\rho)$ are, respectively, the leading coefficients of the orthonormal polynomials associated to the measures σ_l and α .

Proof of Theorem 4.4. Taking the limit when l tends to infinity in (4.9) and using Theorem 4.3, we get

$$\frac{m_{n,p}(\sigma)}{(C(E))^n} \leq \left(\prod_{k=1}^{+\infty} |\Phi(z_k)| \right) \frac{m_{n,p}(\rho)}{(C(E))^n}. \quad (4.10)$$

On the other hand, it is proved in [2] that

$$\lim_{n \rightarrow \infty} \frac{m_{n,p}(\rho)}{(C(E))^n} = (\mu(\rho))^{1/p}. \quad (4.11)$$

Using (4.10), (4.11), and Lemma 3.2, we obtain

$$\limsup_{n \rightarrow \infty} \frac{m_{n,p}(\sigma)}{(C(E))^n} \leq \left(\prod_{k=1}^{+\infty} |\Phi(z_k)| \right) (\mu(\rho))^{1/p} = (\mu^\infty(\rho))^{1/p}. \quad (4.12)$$

It is well known that (see [3, page 231])

$$\forall l > 0, \quad \mu(l) = \mu(\rho) \left(\prod_{k=1}^l |\Phi(z_k)| \right)^p. \quad (4.13)$$

We also have (see [3, Theorem 2.2])

$$\lim_{n \rightarrow \infty} \frac{m_{n,p}(l)}{(C(E))^n} = (\mu(l))^{1/p}. \quad (4.14)$$

From (4.4), we deduce that

$$\forall l > 0, \quad \frac{m_{n,p}(\sigma)}{(C(E))^n} \geq \frac{m_{n,p}(l)}{(C(E))^n}. \quad (4.15)$$

By passing to the limit when n tends to infinity in (4.15) and taking into account (4.13) and (4.14), we get

$$\forall l > 0, \quad \liminf_{n \rightarrow \infty} \frac{m_{n,p}(\sigma)}{(C(E))^n} \geq \left(\prod_{k=1}^l |\Phi(z_k)| \right) (\mu(\rho))^{1/p}. \quad (4.16)$$

Finally, by using [Lemma 3.2](#), we obtain

$$\liminf_{n \rightarrow \infty} \frac{m_{n,p}(\sigma)}{(C(E))^n} \geq \left(\prod_{k=1}^{+\infty} |\Phi(z_k)| \right) (\mu(\rho))^{1/p} = (\mu^\infty(\sigma))^{1/p}. \quad (4.17)$$

Inequalities (4.12) and (4.17) prove [Theorem 4.4\(i\)](#).

We obtain (ii) by proceeding as in [[3](#), pages 234, 235].

To prove (iii), we consider the function

$$\varepsilon_n = \frac{T_{n,p}}{[C(E)\Phi]^n} - \psi^\infty \quad (4.18)$$

which belongs to the space $H^p(\Omega, \rho)$. Then by applying [Lemma 2.1](#), we obtain

$$\begin{aligned} \sup \left\{ \left| \frac{T_{n,p}(z)}{[C(E)\Phi(z)]^n} - \psi^\infty(z) \right| : z \in K \right\} \\ = \sup \{ |\varepsilon_n(z)| : z \in K \} \leq C_K \|\varepsilon_n\|_{H^p(\Omega, \rho)} \longrightarrow 0 \end{aligned} \quad (4.19)$$

for all compact subsets K of Ω . This achieves the proof of the theorem. $\square$

References

- [1] R. Benzine, *Asymptotic behavior of orthogonal polynomials corresponding to a measure with infinite discrete part off a curve*, J. Approx. Theory **89** (1997), no. 2, 257–265.
- [2] Y. L. Geronimus, *On some extremal problems in the space $L_\sigma^{(p)}$* , Mat. Sb. (N.S.) **31(73)** (1952), 3–26 (Russian).
- [3] V. Kaliaguine, *On asymptotics of L_p extremal polynomials on a complex curve ($0 < p < \infty$)*, J. Approx. Theory **74** (1993), no. 2, 226–236.
- [4] R. Khaldi and R. Benzine, *On a generalization of an asymptotic formula of orthogonal polynomials*, Int. J. Appl. Math. **4** (2000), no. 3, 261–274.
- [5] ———, *Asymptotics for orthogonal polynomials off the circle*, J. Appl. Math. **2004** (2004), no. 1, 37–53.
- [6] P. Koosis, *Introduction to H_p Spaces*, London Mathematical Society Lecture Note Series, vol. 40, Cambridge University Press, Cambridge, 1980.
- [7] X. Li and K. Pan, *Asymptotics for L_p extremal polynomials on the unit circle*, J. Approx. Theory **67** (1991), no. 3, 270–283.
- [8] D. S. Lubinsky and E. B. Saff, *Strong asymptotics for L_p extremal polynomials ($1 < p \leq \infty$) associated with weights on $[-1, 1]$* , Approximation Theory (Tampa, Fla, 1985–1986), Lecture Notes in Math., vol. 1287, Springer-Verlag, Berlin, 1987, pp. 83–104.
- [9] F. Peherstorfer and P. Yuditskii, *Asymptotics of orthonormal polynomials in the presence of a denumerable set of mass points*, Proc. Amer. Math. Soc. **129** (2001), no. 11, 3213–3220.
- [10] W. Rudin, *Real and Complex Analysis*, McGraw-Hill, New York, 1966.
- [11] G. Szegő, *Orthogonal Polynomials*, 4th ed., American Mathematical Society Colloquium Publications, vol. 23, American Mathematical Society, Rhode Island, 1975.
- [12] J. P. Tiran and C. Dettaille, *Chebyshev polynomials on a circular arc in the complex plane*, preprint, 1990, Namur University, Belgium.

- [13] H. Widom, *Extremal polynomials associated with a system of curves in the complex plane*, Advances in Math. **3** (1969), 127–232.

Rabah Khaldi: Department of Mathematics, University of Annaba, P.O. Box 12, 23000 Annaba, Algeria

E-mail address: rkhadi@yahoo.fr