Research Article

On Penalty and Gap Function Methods for Bilevel Equilibrium Problems

Bui Van Dinh¹ and Le Dung Muu²

¹ Faculty of Information Technology, Le Quy Don University, Hanoi, Vietnam

² Control and Optimization Department, Institute of Mathematics, VAST, 18 Hoang Quoc Viet, Cau Giay, Hanoi 10307, Vietnam

Correspondence should be addressed to Le Dung Muu, ldmuu@math.ac.vn

Received 9 June 2011; Accepted 15 August 2011

Academic Editor: Ya Ping Fang

Copyright © 2011 B. Van Dinh and L. D. Muu. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We consider bilevel pseudomonotone equilibrium problems. We use a penalty function to convert a bilevel problem into one-level ones. We generalize a pseudo- ∇ -monotonicity concept from ∇ -monotonicity and prove that under pseudo- ∇ -monotonicity property any stationary point of a regularized gap function is a solution of the penalized equilibrium problem. As an application, we discuss a special case that arises from the Tikhonov regularization method for pseudomonotone equilibrium problems.

1. Introduction

Let C be a nonempty closed-convex subset in $\mathbb{R}^n$, and let $f, g : C \times C \rightarrow \mathbb{R}$ be two bifunctions satisfying $f(x, x) = g(x, x) = 0$ for every $x \in C$. Such a bifunction is called an equilibrium bifunction. We consider the following bilevel equilibrium problem (BEP for short):

$$\text{find } \bar{x} \in S_g \text{ such that } f(\bar{x}, y) \geq 0, \quad \forall y \in S_g, \quad (1.1)$$

where $S_g = \{u \in C : g(u, y) \geq 0, \quad \forall y \in C\}$, that is, S_g is the solution set of the equilibrium problems

$$\text{find } u \in C \text{ such that } g(u, y) \geq 0, \quad \forall y \in C. \quad (1.2)$$

As usual, we call problem (1.1) the upper problem and (1.2) the lower one. BEPs are special cases of mathematical programs with equilibrium constraints. Sources for such problems can be found in [1–3]. Bilevel monotone variational inequality, which is a special case of problem

(1.1), was considered in [4, 5]. Moudafi in [6] suggested the use of the proximal point method for monotone BEPs. Recently, Ding in [7] used the auxiliary problem principle to BEPs. In both papers, the bifunctions f and g are required to be monotone on C . It should be noticed that under the pseudomonotonicity assumption on g the solution-set S_g of the lower problem (1.2) is a closed-convex set whenever $g(x, \cdot)$ is lower semicontinuous and convex on C for each x . However, the main difficulty is that, even the constrained set S_g is convex, it is not given explicitly as in a standard mathematical programming problem, and therefore the available methods (see, e.g., [8–14] and the references therein) cannot be applied directly.

In this paper, first, we propose a penalty function method for problem (1.1). Next, we use a regularized gap function for solving the penalized problems. Under certain pseudo- ∇ -monotonicity properties of the regularized bifunction, we show that any stationary point of the gap function on the convex set C is a solution to the penalized subproblem. Finally, we apply the proposed method to the Tikhonov regularization method for pseudomonotone equilibrium problems.

2. A Penalty Function Method

Penalty function method is a fundamental tool widely used in optimization to convert a constrained problem into unconstrained (or easier constrained) ones. This method was used to monotone variational inequalities in [5] and equilibrium problems in [15]. In this section, we use the penalty function method in the bilevel problem (1.1). First, let us recall some well-known concepts on monotonicity and continuity (see, e.g., [16]) that will be used in the sequel.

Definition 2.1. The bifunction $\phi : C \times C \rightarrow \mathbb{R}$ is said to be as follows:

- (a) strongly monotone on C with modulus $\beta > 0$ if

$$\phi(x, y) + \phi(y, x) \leq -\beta \|x - y\|^2, \quad \forall x, y \in C, \quad (2.1)$$

- (b) monotone on C if

$$\phi(x, y) + \phi(y, x) \leq 0, \quad \forall x, y \in C, \quad (2.2)$$

- (c) pseudomonotone on C if

$$\forall x, y \in C : \phi(x, y) \geq 0 \implies \phi(y, x) \leq 0, \quad (2.3)$$

- (d) upper semicontinuous at x with respect to the first argument on C if

$$\overline{\lim_{z \rightarrow x}} \phi(z, y) \leq \phi(x, y), \quad \forall y \in C, \quad (2.4)$$

(e) lower semicontinuous at y with respect to the second argument on C if

$$\lim_{w \rightarrow y} \phi(x, w) \geq \phi(x, y), \quad \forall x \in C. \quad (2.5)$$

Clearly, (a) $\Rightarrow$ (b) $\Rightarrow$ (c).

Definition 2.2 (see [17]). The bifunction $\phi : C \times C \rightarrow \mathbb{R}$ is said to be *coercive* on C if there exists a compact subset $B \subset \mathbb{R}^n$ and a vector $y_0 \in B \cap C$ such that

$$\phi(x, y_0) < 0, \quad \forall x \in C \setminus B. \quad (2.6)$$

Theorem 2.3 (see [18, Proposition 2.1.14]). Let $\phi : C \times C \rightarrow \mathbb{R}$ be a equilibrium bifunction such that $\phi(\cdot, y)$ is upper semicontinuous on C for each $y \in C$ and $\phi(x, \cdot)$ is lower semicontinuous, convex on C for each $x \in C$. Suppose that C is compact or ϕ is coercive on C , then there exists at least one $x^* \in C$ such that $\phi(x^*, y) \geq 0$ for every $y \in C$.

The following proposition tells us about a relationship between the coercivity and the strong monotonicity.

Proposition 2.4. Suppose that the equilibrium bifunction ϕ is strongly monotone on C , and $\phi(x, \cdot)$ is convex, lower semicontinuous with respect to the second argument for all $x \in C$, then for each $y \in C$, there exists a compact set B such that $y \in B$ and $\phi(x, y) < 0$ for all $x \in C \setminus B$.

Proof. Suppose by contradiction that the conclusion does not hold, then there exists an element $y_0 \in C$ such that for every compact set B there is an element $x_B \in C \setminus B$ such that $\phi(x_B, y_0) \geq 0$. Take $B := B_r$ as the closed ball centered at y_0 with radius $r > 1$. Then there exists $x_r \in C \setminus B_r$ such that $\phi(x_r, y_0) \geq 0$. Let x be the intersection of the line segment $[y_0, x_r]$ with the unit sphere $S(y_0; 1)$ centered at y_0 and radius 1. Hence, $x_r = y_0 + t(r)(x - y_0)$, where $t(r) > r$. By the strong monotonicity of ϕ , we have

$$\phi(y_0, x_r) \leq -\phi(x_r, y_0) - \beta \|x_r - y_0\|^2 \leq -\phi(x_r, y_0) - \beta t(r)^2 \|x - y_0\|^2. \quad (2.7)$$

Since $\phi(y_0, \cdot)$ is convex on C , it follows that

$$\phi(y_0, x) \leq \frac{1}{t(r)} \phi(y_0, x_r) + \frac{t(r) - 1}{t(r)} \phi(y_0, y_0), \quad (2.8)$$

which implies that $\phi(y_0, x) \leq -\beta t(r) \|x - y_0\|^2 \leq -\beta r$. Thus,

$$\phi(y_0, x) \longrightarrow -\infty \quad \text{as } r \longrightarrow \infty. \quad (2.9)$$

However, since $\phi(y_0, \cdot)$ is lower semicontinuous on C , by the well-known Weierstrass Theorem, $\phi(y_0, \cdot)$ attains its minimum on the compact set $S(y_0; 1) \cap C$. This fact contradicts (2.9). $\square$

From this proposition, we can derive the following corollaries.

Corollary 2.5 (see [18]). *If the bifunction ϕ is strongly monotone on C , and $\phi(x, \cdot)$ is convex, lower semicontinuous with respect to the second argument for all $x \in C$, then ϕ is coercive on C .*

Corollary 2.6. *Suppose that the bifunction f is strongly monotone on C , and $f(x, \cdot)$ is convex, lower semicontinuous with respect to the second argument for all $x \in C$. If the bifunction g is coercive on C then, for every $\epsilon > 0$, the bifunction $g + \epsilon f$ is uniformly coercive on C , for example, there exists a point $y_0 \in C$ and a compact set B both independent of ϵ such that*

$$g(x, y_0) + \epsilon f(x, y_0) < 0, \quad \forall x \in C \setminus B. \quad (2.10)$$

Proof. From the coercivity of g , we conclude that there exists a compact B_1 and $y_0 \in C$ such that $g(x, y_0) < 0$ for all $x \in C \setminus B_1$. Since f is strongly monotone, convex, lower semicontinuous on C , by choosing $y = y_0$, from Proposition 2.4, there exists a compact B_2 such that $f(x, y_0) < 0$ for all $x \in C \setminus B_2$. Set $B = B_1 \cup B_2$, then B is compact and $g(x, y_0) + \epsilon f(x, y_0) < 0$ for all $x \in C \setminus B$. $\square$

Remark 2.7. It is worth to note that if both f, g are coercive and pseudomonotone on C , then the function $f + g$ is not necessary coercive or pseudomonotone on C .

To see this, let us consider the following bifunctions.

Example 2.8. Let $f(x, y) := (x_1 y_2 - x_2 y_1) e^{x_1}$, $g(x, y) := (x_2 y_1 - x_1 y_2) e^{x_2}$, and $C = \{(x_1, x_2) : x_1 \geq -1, (1/10)(x_1 - 9) \leq x_2 \leq 10x_1 + 9\}$ then we have

- (i) $f(x, y), g(x, y)$ are pseudomonotone and coercive on C ,
- (ii) for all $\epsilon > 0$ the bifunctions $f_\epsilon(x, y) = g(x, y) + \epsilon f(x, y)$ are neither pseudomonotone nor coercive on C .

Indeed,

- (i) if $f(x, y) \leq 0$, then $f(y, x) \geq 0$, thus f is pseudomonotone on C . By choosing $y^0 = (y_1^0, 0), (0 < y_1^0 \leq 1)$ and $B = \{(x_1, x_2) : x_1^2 + x_2^2 \leq r\} (r > 1)$, we have $f(x, y^0) = -x_2 y_1^0 e^{x_1} < 0$ for all $y \in C \setminus B$, which means that f is coercive on C . Similarly, we can see that g is coercive on C ,
- (ii) by definition of f , we have that

$$f_\epsilon(x, y) = (x_2 y_1 - x_1 y_2)(e^{x_2} - \epsilon e^{x_1}), \quad \forall \epsilon > 0. \quad (2.11)$$

Take $x(t) = (t, 2t)$, for all $y(t) = (2t, t)$, then $f_\epsilon(x(t), y(t)) = 3t^2(e^{2t} - \epsilon e^t) > 0$, whereas $f_\epsilon(y(t), x(t)) = -3t^2(e^t - \epsilon e^{2t}) > 0$ for t is sufficiently large. So f_ϵ is not pseudomonotone on C .

Now, we show that the bifunction $f_\epsilon(x, y) = (x_2 y_1 - x_1 y_2)(e^{x_2} - \epsilon e^{x_1})$ is not coercive on C . Suppose, by contradiction, that there exist a compact set B and $y^0 = (y_1^0, y_2^0) \in B \cap C$ such that $f_\epsilon(x, y^0) < 0$ for all $x \in C \setminus B$, then, by coercivity of f_ϵ , it follows, $y_1^0, y_2^0 > 0$ and $y_1^0 \neq y_2^0$. With $x(t) = (t, kt), (t > 0)$, we have $f_\epsilon(x(t), y^0) = t(ky_1^0 - y_2^0)(e^{kt} - \epsilon e^t)$. However

- (i) if $y_1^0 > y_2^0$, then from $1 < k < 10$ it follows that $x(t) \in C$ and $f_\epsilon(x(t), y^0) > 0$ for t is sufficiently large, which contradicts with coercivity,

- (ii) if $y_1^0 < y_2^0$, then, by choosing $1/10 < k < 1$, we obtain $x(t) \in C$ and $f_\epsilon(x(t), y^0) > 0$ for t is large enough. But this cannot happen because of the coercivity of f_ϵ .

Now, for each fixed $\epsilon > 0$, we consider the penalized equilibrium problem $\text{PEP}(C, f_\epsilon)$ defined as

$$\text{find } \bar{x}_\epsilon \in C \text{ such that } f_\epsilon(\bar{x}_\epsilon, y) := g(\bar{x}_\epsilon, y) + \epsilon f(\bar{x}_\epsilon, y) \geq 0, \quad \forall y \in C. \quad (2.12)$$

By $\text{SOL}(C, f_\epsilon)$, we denote the solution set of $\text{PEP}(C, f_\epsilon)$.

Theorem 2.9. *Suppose that the equilibrium bifunctions f, g are pseudomonotone, upper semicontinuous with respect to the first argument and lower semicontinuous, convex with respect to the second argument on C , then any cluster point of the sequence $\{x_k\}$ with $x_k \in \text{SOL}(C, f_{\epsilon_k})$, $\epsilon_k \rightarrow 0$ is a solution to the original bilevel problem (1.1). In addition, if f is strongly monotone and g is coercive on C , then for each $\epsilon_k > 0$ the penalized problem $\text{PEP}(C, f_{\epsilon_k})$ is solvable, and any sequence $\{x_k\}$ with $x_k \in \text{SOL}(C, f_{\epsilon_k})$ converges to the unique solution of the bilevel problem (1.1) as $k \rightarrow \infty$.*

Proof. Let $\{x_k\}$ be any sequence with $x_k \in \text{SOL}(C, f_{\epsilon_k})$, and let $\bar{x}$ be any of its cluster points. Without loss of generality, we may assume that $x_k \rightarrow \bar{x}$ as $k \rightarrow \infty$. Since $x_k \in \text{SOL}(C, f_{\epsilon_k})$, one has

$$g(x_k, y) + \epsilon_k f(x_k, y) \geq 0, \quad \forall y \in C. \quad (2.13)$$

For any $z \in S_g$, we have $g(z, y) \geq 0$, for all $y \in C$ and in particular, $g(z, x_k) \geq 0$. Then, by the pseudomonotonicity of g , we have $g(x_k, z) \leq 0$. Replacing y by z in (2.13), we obtain

$$g(x_k, z) + \epsilon_k f(x_k, z) \geq 0, \quad (2.14)$$

which implies that

$$\epsilon_k f(x_k, z) \geq -g(x_k, z) \geq 0 \implies f(x_k, z) \geq 0. \quad (2.15)$$

Let $k \rightarrow \infty$, by upper semicontinuity of f , we have $f(\bar{x}, z) \geq 0$ for all $z \in S_g$.

To complete the proof, we need only to show that $\bar{x} \in S_g$. Indeed, for any $y \in C$, we have

$$g(x_k, y) + \epsilon_k f(x_k, y) \geq 0, \quad \forall y \in C. \quad (2.16)$$

Again, by upper semicontinuity of f and g , we obtain in the limit, as $\epsilon_k \rightarrow 0$, that $g(\bar{x}, y) \geq 0$ for all $y \in C$. Hence, $\bar{x} \in S_g$.

Now suppose, in addition, that f is strongly monotone on C . By Corollary 2.6, f_{ϵ_k} is uniformly coercive on C . Thus, problem $\text{PEP}(C, f_{\epsilon_k})$ is solvable and, for all $\epsilon_k > 0$, the solution sets of these problems are contained in a compact set B . So any infinite sequence $\{x_k\}$ of the solutions has a cluster point, say, $\bar{x}$. By the first part, $\bar{x}$ is a solution of (1.1). Note that, from the assumption on g , the solution set S_g of the lower equilibrium ($\text{EP}(C, g)$) is a closed, convex, compact set. Since f is lower semicontinuous and convex with respect to the second

argument and is strongly monotone on C , the upper equilibrium problem $EP(S_g, f)$ has a unique solution. Using again the first part of the theorem, we can see that $x_k \rightarrow \bar{x}$ as $k \rightarrow \infty$ $\square$

Remark 2.10. In a special case considered in [6], where both f and g are monotone, the penalized problem (PEP) is monotone too. In this case, (PEP) can be solved by some existing methods (see, e.g., [6, 11–14, 19]) and the references therein. However, when one of these two bifunctions is pseudomonotone, the penalized problem (PEP), in general, does not inherit any monotonicity property from f and g . In this case, problem (PEP) cannot be solved by the above-mentioned existing methods.

3. Gap Function and Descent Direction

A well-known tool for solving equilibrium problem is the gap function. The regularized gap function has been introduced by Taji and Fukushima in [20] for variational inequalities, and extended by Mastroeni in [11] to equilibrium problems. In this section, we use the regularized gap function for the penalized equilibrium problem (PEP). As we have mentioned above, this problem, even when g is pseudomonotone and f is strongly monotone, is still difficult to solve.

Throughout this section, we suppose that both f and g are lower semicontinuous, convex on C with respect to the second argument. First, we recall (see, e.g., [11]) the definition of a gap function for the equilibrium problem.

Definition 3.1. A function $\varphi : C \rightarrow \mathbb{R} \cup \{+\infty\}$ is said to be a gap function for (PEP) if

- (i) $\varphi(x) \geq 0$, for all $x \in C$,
- (ii) $\varphi(\bar{x}) = 0$ if and only if $\bar{x}$ is a solution for (PEP).

A gap function for (PEP) is $\varphi(x) = -\min_{y \in C} f_e(x, y)$. This gap function may not be finite and, in general, is not differentiable. To obtain a finite, differentiable gap function, we use the regularized gap function introduced in [20] and recently used by Mastroeni in [11] to equilibrium problems. From Proposition 2.2 and Theorem 2.1 in [11], the following proposition is immediate.

Proposition 3.2. Suppose that $l : C \times C \rightarrow \mathbb{R}$ is a nonnegative differentiable, strongly convex bifunction on C with respect to the second argument and satisfies

- (a) $l(x, x) = 0$ for all $x \in C$,
- (b) $\nabla_y l(x, x) = 0$ for all $x \in C$.

Then the function

$$\varphi_\epsilon(x) = -\min_{y \in C} [g(x, y) + \epsilon[f(x, y) + l(x, y)]] \quad (3.1)$$

is a finite gap function for (PEP). In addition, if f and g are differentiable with respect to the first argument and $\nabla_x f(x, y), \nabla_x g(x, y)$ are continuous on C , then $\varphi_\epsilon(x)$ is continuously differentiable on C and

$$\nabla \varphi_\epsilon(x) = -\nabla_x g(x, y_\epsilon(x)) - \epsilon \nabla_x [f(x, y_\epsilon(x)) + l(x, y_\epsilon(x))] = -\nabla_x g_\epsilon(x, y_\epsilon(x)), \quad (3.2)$$

where

$$\begin{aligned} g_\epsilon(x, y) &= g(x, y) + \epsilon[f(x, y) + l(x, y)], \\ y_\epsilon(x) &= \arg \min_{y \in C} \{g_\epsilon(x, y)\}. \end{aligned} \quad (3.3)$$

Note that the function $l(x, y) := (1/2)\langle M(y - x), y - x \rangle$, where M is a symmetric positive definite matrix of order n that satisfies the assumptions on l .

We need some definitions on ∇ -monotonicity.

Definition 3.3. A differentiable bifunction $h : C \times C \rightarrow \mathbb{R}$ is called as follows:

(a) strongly ∇ -monotone on C if there exists a constant $\tau > 0$ such that,

$$\langle \nabla_x h(x, y) + \nabla_y h(x, y), y - x \rangle \geq \tau \|y - x\|^2, \quad \forall x, y \in C, \quad (3.4)$$

(b) strictly ∇ -monotone on C if

$$\langle \nabla_x h(x, y) + \nabla_y h(x, y), y - x \rangle > 0, \quad \forall x, y \in C, \quad x \neq y, \quad (3.5)$$

(c) ∇ -monotone on C if

$$\langle \nabla_x h(x, y) + \nabla_y h(x, y), y - x \rangle \geq 0, \quad \forall x, y \in C, \quad (3.6)$$

(d) strictly pseudo- ∇ -monotone on C if

$$\langle \nabla_x h(x, y), y - x \rangle \leq 0 \implies \langle \nabla_y h(x, y), y - x \rangle > 0, \quad \forall x, y \in C, \quad x \neq y, \quad (3.7)$$

(e) pseudo- ∇ -monotone on C if

$$\langle \nabla_x h(x, y), y - x \rangle \leq 0 \implies \langle \nabla_y h(x, y), y - x \rangle \geq 0, \quad \forall x, y \in C. \quad (3.8)$$

Remark 3.4. The definitions (a), (b), and (c) can be found, for example, in [8, 11]. The definitions (d) and (e), to our best knowledge, are not used before. From the definitions, we have

$$(a) \implies (b) \implies (c) \implies (e), \quad (a) \implies (b) \implies (d) \implies (e). \quad (3.9)$$

However, (c) may not imply (d) and vice versa as shown by the following simple examples.

Example 3.5. Consider the bifunction $h(x, y) = e^{x^2}(y^2 - x^2)$ defined on $C \times C$ with $C = \mathbb{R}$. This bifunction is not ∇ -monotone on C , because

$$\langle \nabla_x h(x, y) + \nabla_y h(x, y), y - x \rangle = 2e^{x^2}(y - x)^2(x^2 + xy + 1) \quad (3.10)$$

is negative for $x = -1$, $y = 3$. However, $h(x, y)$ is strictly pseudo- ∇ -monotone. Indeed, we have

$$\begin{aligned}\langle \nabla_x h(x, y), y - x \rangle &= 2xe^{x^2}(y^2 - x^2 - 1)(y - x) \leq 0 \iff x(y^2 - x^2 - 1)(y - x) \leq 0, \\ \langle \nabla_y h(x, y), y - x \rangle &= 2ye^{x^2}(y - x) > 0 \iff y(y - x) > 0.\end{aligned}\quad (3.11)$$

It is not difficult to verify that

$$x(y^2 - x^2 - 1)(y - x) \leq 0 \implies y(y - x) > 0, \quad \text{as } x \neq y. \quad (3.12)$$

Hence this function is strictly pseudo- ∇ -monotone but is not ∇ -monotone.

Vice versa, considering the bifunction $h(x, y) = (y - x)^T M (y - x)$ defined on $\mathbb{R}^n \times \mathbb{R}^n$, where M is a matrix of order $n \times n$, we have the following:

(i) h is ∇ -monotone, because

$$\begin{aligned}\langle \nabla_x h(x, y) + \nabla_y h(x, y), y - x \rangle \\ = \langle -(y - x)^T (M + M^T) + (y - x)^T (M + M^T), y - x \rangle = 0, \quad \forall x, y.\end{aligned}\quad (3.13)$$

Clearly, h is not strictly- ∇ -monotone,

(ii) h is strictly pseudo ∇ -monotone if and only if

$$\langle \nabla_x h(x, y), y - x \rangle = -\langle (y - x)^T (M + M^T), y - x \rangle \leq 0 \quad (3.14)$$

implies

$$\langle \nabla_y h(x, y), y - x \rangle = (y - x)^T (M + M^T), y - x > 0, \quad \forall x, y, \quad x \neq y. \quad (3.15)$$

The latter inequality equivalent to $M + M^T$ is a positive definite matrix of order $n \times n$.

Remark 3.6. As shown in [8] when $h(x, y) = \langle T(x), y - x \rangle$ with T a differentiable monotone operator on C , h is monotone on C if and only if T is monotone on C , and in this case, monotonicity of h on C coincides with ∇ -monotonicity of h on C .

The following example shows that pseudomonotonicity may not imply pseudo- ∇ -monotonicity.

Example 3.7. Let $h(x, y) = -ax(y - x)$, defined on $\mathbb{R}_+ \times \mathbb{R}_+$, ($a > 0$). It is easy to see that

$$h(x, y) \geq 0 \implies h(y, x) \leq 0, \quad \forall x, y \geq 0. \quad (3.16)$$

Thus, h is pseudomonotone on $\mathbb{R}_+$.

We have

$$\langle \nabla_x h(x, y), y - x \rangle = -a(y - x)(y - 2x) < 0, \quad \forall y > 2x > 0. \quad (3.17)$$

But

$$\langle \nabla_y h(x, y), y - x \rangle = -ax(y - x) < 0, \quad \forall y > 2x > 0. \quad (3.18)$$

So h is not pseudo- ∇ -monotone on $\mathbb{R}_+$.

From the definition of the gap function φ_ϵ , a global minimal point of this function over C is a solution to problem (PEP). Since φ_ϵ is not convex, its global minimum is extremely difficult to compute. In [8], the authors have shown that under the strict ∇ -monotonicity a stationary point is also a global minimum of gap function. By a counterexample, the authors in [8] also pointed out that the strict ∇ -monotonicity assumption cannot be relaxed to ∇ -monotonicity. The following theorem shows that the stationary property is still guaranteed under the strict pseudo- ∇ -monotonicity.

Theorem 3.8. *Suppose that g_ϵ is strictly pseudo- ∇ -monotone on C . If $\bar{x}$ is a stationary point of φ_ϵ over C , that is,*

$$\langle \nabla \varphi_\epsilon(\bar{x}), y - \bar{x} \rangle \geq 0, \quad \forall y \in C. \quad (3.19)$$

then $\bar{x}$ solves (PEP).

Proof. Suppose that $\bar{x}$ does not solve (PEP), then $y_\epsilon(\bar{x}) \neq \bar{x}$.

Since $\bar{x}$ is a stationary point of φ_ϵ on C , from the definition of φ_ϵ , we have

$$\langle \nabla \varphi_\epsilon(\bar{x}), y - \bar{x} \rangle = -\langle \nabla_x g_\epsilon(x, y_\epsilon(x)), y_\epsilon(x) - x \rangle \geq 0. \quad (3.20)$$

By strict pseudo- ∇ -monotonicity of g_ϵ , it follows that

$$\langle \nabla_y g_\epsilon(\bar{x}, y_\epsilon(\bar{x})), y_\epsilon(\bar{x}) - \bar{x} \rangle > 0. \quad (3.21)$$

On the other hand, since $y_\epsilon(\bar{x})$ minimizes $g_\epsilon(x, \cdot)$ over C , we have

$$\langle \nabla_y g_\epsilon(\bar{x}, y_\epsilon(\bar{x})), y_\epsilon(\bar{x}) - \bar{x} \rangle \leq 0, \quad (3.22)$$

which is in contradiction with (3.21). $\square$

To compute a stationary point of a differentiable function over a closed-convex set, we can use the existing descent direction algorithms in mathematical programming (see, e.g., [8, 21]). The next proposition shows that if $y(x)$ is a solution of the problem $\min_{y \in C} g_\epsilon(x, y)$, then $y(x) - x$ is a descent direction on C of φ_ϵ at x . Namely, we have the following proposition.

Proposition 3.9. *Suppose that g_ϵ is strictly pseudo- ∇ -monotone on C and x is not a solution to Problem (PEP), then*

$$\langle \nabla \varphi_\epsilon(x), y_\epsilon(x) - x \rangle < 0. \quad (3.23)$$

Proof. Let $d_\epsilon(x) = y_\epsilon(x) - x$. Since x is not a solution to (PEP), then $d_\epsilon(x) \neq 0$. Suppose that, by contradiction, $d_\epsilon(x)$ is not a descent direction on C of φ_ϵ at x , then

$$\langle \nabla \varphi_\epsilon(x), y_\epsilon(x) - x \rangle \geq 0 \iff -\langle \nabla_x g_\epsilon(x, y_\epsilon(x)), y_\epsilon(x) - x \rangle \geq 0, \quad (3.24)$$

which, by strict pseudo- ∇ -monotonicity of g_ϵ , implies

$$\langle \nabla_y g_\epsilon(x, y_\epsilon(x)), y_\epsilon(x) - x \rangle > 0. \quad (3.25)$$

On the other hand, since $y_\epsilon(x)$ minimizes $g_\epsilon(x, \cdot)$ over C , by the well-known optimality condition, we have

$$\langle \nabla_y g_\epsilon(x, y_\epsilon(x)), y_\epsilon(x) - x \rangle \leq 0, \quad (3.26)$$

which contradicts (3.25). $\square$

Proposition 3.10. *Suppose that $g(x, \cdot)$ is strictly convex on C for every $x \in C$ and g is strictly pseudo- ∇ -monotone on C . If $x \in C$ is not a solution of (PEP), then there exists $\bar{\epsilon} > 0$ such that $y_\epsilon(x) - x$ is a descent direction of φ_ϵ on C at x for all $0 < \epsilon \leq \bar{\epsilon}$.*

Proof. By contradiction, suppose that the statement of the proposition does not hold, then there exist $\epsilon_k \searrow 0$ and $x \in C$ such that

$$\langle \nabla \varphi_{\epsilon_k}(x), y_{\epsilon_k}(x) - x \rangle \geq 0 \iff -\langle \nabla_x g_{\epsilon_k}(x, y_{\epsilon_k}(x)), y_{\epsilon_k}(x) - x \rangle \geq 0. \quad (3.27)$$

Since $g_\epsilon(x, \cdot)$ is strictly convex differentiable on C , by Theorem 2.1 in [9], the function $\epsilon \mapsto y_\epsilon(x)$ is continuous with respect to ϵ , thus $y_{\epsilon_k}(x)$ tends to $y_0(x)$ as $\epsilon_k \rightarrow 0$, where $y_0(x) = \arg \min_{y \in C} g(x, y)$. Since $g_{\epsilon_k}(x, y) = g(x, y) + \epsilon_k f(x, y)$ is continuously differentiable, letting $\epsilon_k \rightarrow 0$ in (3.27), we obtain

$$-\langle \nabla_x g(x, y_0(x)), y_0(x) - x \rangle \geq 0. \quad (3.28)$$

By strict pseudo- ∇ -monotonicity of g , it follows that

$$\langle \nabla_y g(x, y_0(x)), y_0(x) - x \rangle > 0. \quad (3.29)$$

On the other hand, since $y_{\epsilon_k}(x)$ minimizes $g_{\epsilon_k}(x, \cdot)$ over C , we have

$$\langle \nabla_y g_{\epsilon_k}(x, y_{\epsilon_k}(x)), y_{\epsilon_k}(x) - x \rangle \leq 0. \quad (3.30)$$

Taking the limit, we obtain

$$\langle \nabla_y g(x, y_0(x)), y_0(x) - x \rangle \leq 0, \quad (3.31)$$

which contradicts (3.29). $\square$

To illustrate Theorem 3.8, let us consider the following examples.

Example 3.11. Consider the bifunctions $g(x, y) = e^{x^2}(y^2 - x^2)$ and $f(x, y) = 10^{x^2}(y^2 - x^2)$ defined on $\mathbb{R} \times \mathbb{R}$. It is not hard to verify that,

- (i) $g(x, y), f(x, y)$ are monotone, strictly pseudo- ∇ -monotone on $\mathbb{R}$,
- (ii) for all $\epsilon > 0$ the bifunction $g(x, y) + \epsilon f(x, y)$ is monotone and strictly pseudo- ∇ -monotone on $\mathbb{R}$ and satisfying all of the assumptions of Theorem 3.8.

Example 3.12. Let $f(x, y) = -x^2 - xy + 2y^2$ and $g(x, y) = -3x^2y + xy^2 + 2y^3$ defined on $\mathbb{R}_+ \times \mathbb{R}_+$ it is easy to see that,

- (i) g, f are pseudomonotone, strictly ∇ -monotone on $\mathbb{R}_+$,
- (ii) for all $\epsilon > 0$ the bifunction $g(x, y) + \epsilon f(x, y)$ is pseudomonotone and strictly ∇ -monotone on $\mathbb{R}_+$ and satisfying all of the assumptions of Theorem 3.8.

4. Application to the Tikhonov Regularization Method

The Tikhonov method [22] is commonly used for handling ill-posed problems. Recently, in [23] the Tikhonov method has been extended to the pseudomonotone equilibrium problem

$$\text{Find } x^* \in C \text{ such that } g(x^*, y) \geq 0, \quad \forall y \in C, \quad (\text{EP}(C, g))$$

where, as before, C is a closed-convex set in $\mathbb{R}^n$ and $g : C \rightarrow \mathbb{R}$ is a pseudomonotone bifunction satisfying $g(x, x) = 0$ for every $x \in C$.

In the Tikhonov regularization method considered in [23], problem $(\text{EP}(C, g))$ is regularized by the problems

$$\text{find } x^* \in C \text{ such that } g_\epsilon(x^*, y) := g(x^*, y) + \epsilon f(x^*, y) \geq 0, \quad \forall y \in C, \quad (\text{EP}(C, g_\epsilon))$$

where f is an equilibrium bifunction on C and $\epsilon > 0$ and play the role of the regularization bifunction and regularization parameter, respectively.

In [23], the following theorem has been proved.

Theorem 4.1. Suppose that $f(\cdot, y), g(\cdot, y)$ are upper semicontinuous and $f(x, \cdot), g(x, \cdot)$ are lower semicontinuous convex on C for each $x, y \in C$ and that g is pseudomonotone on C . Suppose further that f is strongly monotone on C satisfying the condition

$$\exists \delta > 0: \quad |f(x, y)| \leq \delta \|x - x^g\| \|y - x\|, \quad \forall x, y \in C, \quad (4.1)$$

where $x^g \in C$ (plays the role of a guess solution) is given.

Then the following three statements are equivalent:

- (a) the solution set of $(EP(C, g_\epsilon))$ is nonempty for each $\epsilon > 0$ and $\lim_{\epsilon \rightarrow 0^+} x(\epsilon)$ exists, where $x(\epsilon)$ is arbitrarily chosen in the solution set of $(EP(C, g_\epsilon))$,
- (b) the solution set of $(EP(C, g_\epsilon))$ is nonempty for each $\epsilon > 0$ and $\lim_{\epsilon \rightarrow 0^+} \sup \|x(\epsilon)\| < \infty$, where $x(\epsilon)$ is arbitrarily chosen in the solution set of $(EP(C, g_\epsilon))$,
- (c) the solution set of $(EP(C, g))$ is nonempty.

Moreover, if any one of these statements holds, then $\lim_{\epsilon \rightarrow 0^+} x(\epsilon)$ is equal to the unique solution of the strongly monotone equilibrium problem $EP(S_g, f)$, where S_g denotes the solution set of the original problem $(EP(C, g))$.

Note that, when g is monotone on C , the regularized subproblems are strongly monotone and therefore, they can be solved by some existing methods. When g is pseudomonotone, the subproblems, in general, are no longer strongly monotone, even not pseudomonotone. So solving them becomes a difficult task. However, the problem of finding the limit point of the sequences of iterates leads to the unique solution of problem $EP(S_g, f)$.

In order to apply the penalty and gap function methods described in the preceding sections, let us take, for instant,

$$f(x, y) = \langle x - x^g, y - x \rangle. \quad (4.2)$$

Clearly, f is both strongly monotone and strongly ∇ -monotone with the same modulus 1. Moreover, f satisfies the condition (4.1). Therefore, the problem of finding the limit point in the above Tikhonov regularization method can be formulated as the bilevel equilibrium problem

$$\text{find } x \in S_g \text{ such that } f(x^*, y) \geq 0, \quad \forall y \in S_g, \quad (4.3)$$

which is of the form (1.1). Now, for each fixed $\epsilon_k > 0$, we consider the penalized equilibrium problem $PEP(C, f_{\epsilon_k})$ defined as

$$\text{find } \bar{x}_k \in C \text{ such that } f_{\epsilon_k}(\bar{x}_k, y) := g(\bar{x}_k, y) + \epsilon_k f(\bar{x}_k, y) \geq 0, \quad \forall y \in C. \quad (4.4)$$

As before, by $SOL(C, f_{\epsilon_k})$, we denote the solution set of $PEP(C, f_{\epsilon_k})$.

Applying Theorems 2.9 and 3.8, we obtain the following result.

Theorem 4.2. Suppose that the bifunction g satisfies the following conditions:

- (i) $g(x, \cdot)$ is convex, lower semicontinuous for all $x \in C$,
- (ii) g is pseudomonotone and coercive on C .

Then for any $\epsilon_k > 0$, the penalized problem $PEP(C, f_{\epsilon_k})$ is solvable, and any sequence $\{x_k\}$ with $x_k \in SOL(C, f_{\epsilon_k})$ for all k converges to the unique solution of the problem (4.3) as $k \rightarrow \infty$.

- (iii) In addition, if $g(x, y) + \epsilon_k f(x, y)$ is strictly pseudo- ∇ -monotone on C (in particular, $g(x, y)$ is ∇ -monotone), and $\bar{x}_k$ is any stationary point of the mathematical program $\min_{x \in C} \varphi_k(x)$ with

$$\varphi_k(x) := \min_{y \in C} \{g(x, y) + \epsilon_k f(x, y)\}, \quad (4.5)$$

then $\{\bar{x}_k\}$ converges to the unique solution of the problem (4.3) as $k \rightarrow \infty$.

5. Conclusion

We have considered a class of bilevel pseudomonotone equilibrium problems. The main difficulty of this problem is that its feasible domain is not given explicitly as in a standard mathematical programming problem. We have proposed a penalty function method to convert the bilevel problem into one-level ones. Then we have applied the regularized gap function method to solve the penalized equilibrium subproblems. We have generalized the pseudo- ∇ -monotonicity concept from ∇ -monotonicity. Under the pseudo- ∇ -monotonicity property, we have proved that any stationary point of the gap function is a solution to the original bilevel problem. As an application, we have shown how to apply the proposed method to the Tikhonov regularization method for pseudomonotone equilibrium problems.

Acknowledgment

This work is supported by the National Foundation for Science Technology Development of Vietnam (NAFOSTED).

References

- [1] J. Q. Luo, J. S. Pang, and D. Ralph, *Mathematical Programs with Equilibrium Constraints*, Cambridge University Press, Cambridge, UK, 1996.
- [2] M. A. Migdalas, P. Pardalos, and P. Varbrand, *Multilevel Optimization: Algorithms and Applications*, vol. 20 of *Nonconvex Optimization and its Applications*, Kluwer Academic Publishers, Dordrecht, The Netherlands, 1998.
- [3] L. D. Muu and N. V. Quy, "A global optimization method for solving convex quadratic bilevel programming problems," *Journal of Global Optimization*, vol. 26, no. 2, pp. 199–219, 2003.
- [4] P. N. Anh, J. Kim, and L. D. Muu, "An extragradient algorithm for solving bilevel variational inequalities," to appear in *Journal of Global Optimization*.
- [5] V. V. Kalashnikov and N. I. Kalashnikova, "Solving two-level variational inequality," *Journal of Global Optimization*, vol. 8, no. 3, pp. 289–294, 1996.
- [6] A. Moudafi, "Proximal methods for a class of bilevel monotone equilibrium problems," *Journal of Global Optimization*, vol. 47, no. 2, pp. 287–292, 2010.
- [7] X. P. Ding, "Auxiliary principle and algorithm for mixed equilibrium problems and bilevel mixed equilibrium problems in Banach spaces," *Journal of Optimization Theory and Applications*, vol. 146, no. 2, pp. 347–357, 2010.
- [8] G. Bigi, M. Castellani, and M. Pappalardo, "A new solution method for equilibrium problems," *Optimization Methods & Software*, vol. 24, no. 6, pp. 895–911, 2009.
- [9] M. Castellani and M. Pappalardo, "Gap functions for nonsmooth equilibrium problems," *Taiwanese Journal of Mathematics*, vol. 13, no. 6A, pp. 1837–1846, 2009.
- [10] A. N. Iusem and W. Sosa, "Iterative algorithms for equilibrium problems," *Optimization*, vol. 52, no. 3, pp. 301–316, 2003.

- [11] G. Mastroeni, "Gap functions for equilibrium problems," *Journal of Global Optimization*, vol. 27, no. 4, pp. 411–426, 2003.
- [12] L. D. Muu, V. H. Nguyen, and T. D. Quoc, "Extragradient algorithms extended to equilibrium problems," *Optimization*, vol. 57, no. 6, pp. 749–776, 2008.
- [13] L. D. Muu and T. D. Quoc, "Regularization algorithms for solving monotone Ky Fan inequalities with application to a Nash-Cournot equilibrium model," *Journal of Optimization Theory and Applications*, vol. 142, no. 1, pp. 185–204, 2009.
- [14] T. T. V. Nguyen, J. J. Strodiot, and V. H. Nguyen, "A bundle method for solving equilibrium problems," *Mathematical Programming*, vol. 116, no. 1-2, pp. 529–552, 2009.
- [15] L. D. Muu and W. Oettli, "Convergence of an adaptive penalty scheme for finding constrained equilibria," *Nonlinear Analysis*, vol. 18, no. 12, pp. 1159–1166, 1992.
- [16] M. Bianchi and S. Schaible, "Generalized monotone bifunctions and equilibrium problems," *Journal of Optimization Theory and Applications*, vol. 90, no. 1, pp. 31–43, 1996.
- [17] E. Blum and W. Oettli, "From optimization and variational inequalities to equilibrium problems," *The Mathematics Student*, vol. 63, no. 1–4, pp. 127–149, 1994.
- [18] I. Konnov, *Combined Relaxation Methods for Variational Inequalities*, vol. 495 of *Lecture Notes in Economics and Mathematical Systems*, Springer, Berlin, Germany, 2001.
- [19] A. Moudafi, "Proximal point algorithm extended to equilibrium problems," *Journal of Natural Geometry*, vol. 15, no. 1-2, pp. 91–100, 1999.
- [20] K. Taji and M. Fukushima, "A new merit function and a successive quadratic programming algorithm for variational inequality problems," *SIAM Journal on Optimization*, vol. 6, no. 3, pp. 704–713, 1996.
- [21] D. P. Bertsekas, *Nonlinear Programming*, Athena Scientific, Belmont, Mass, USA, 2nd edition, 1999.
- [22] A. B. Bakushinsky and A. Goncharsky, *Ill-Posed Problems: Theory and Applications*, vol. 301 of *Mathematics and its Applications*, Kluwer Academic Publishers Group, Dodrecht, The Netherlands, 1994.
- [23] L. G. Hung and L. D. Muu, "The Tikhonov regularization method extended to equilibrium problems involving pseudomonotone bifunctions," *Nonlinear Analysis*, vol. 74, no. 17, pp. 6121–6129, 2011.