Research Article

An Iteration Method Generating Analytical Solutions for Blasius Problem

Beong In Yun

Department of Informatics and Statistics, Kunsan National University, Kunsan 573-701, Republic of Korea

Correspondence should be addressed to Beong In Yun, paulllyun@gmail.com

Received 7 April 2011; Revised 31 May 2011; Accepted 29 June 2011

Academic Editor: Ch Tsitouras

Copyright © 2011 Beong In Yun. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We derive a new iteration method for finding solution of the generalized Blasius problem. This method results in the analytical series solutions which are consistent with the existing series solutions for some special cases.

1. Introduction

We consider the generalized Blasius' equation

$$y'''(x) + \alpha y(x)y''(x) = 0, \quad 0 \leq x < \infty, \quad (1.1)$$

where $\alpha = 1/2$ or $\alpha = 1$, with boundary conditions

$$y(0) = 0, \quad y'(0) = \mu, \quad y'(\infty) = 1. \quad (1.2)$$

This problem describes the boundary layer flow over a moving plate with constant velocity μ . For a special case of $\alpha = 1/2$ and $\mu = 0$, the series solution of the Blasius problem becomes

$$S(x) = \frac{\kappa}{2}x^2 - \frac{\kappa^2}{240}x^5 + \frac{11}{161280}\kappa^3x^8 - \frac{5}{4257792}\kappa^4x^{11} + \cdots, \quad (1.3)$$

where $\kappa = y''(0) \approx 0.3320573362$. The Blasius series, however, converges for $|x| < \rho = 5.6900380545$. In the literature [1–3], it was shown that the limitation can be overcome by Padé approximants or an Euler-accelerated series.

Lots of analytical methods such as Adomian decomposition methods [4–6], variational iteration methods [7–11], and homotopy analysis methods [12–14] have been proposed.

2. Derivation of an Iteration Formula

We develop a new iteration method to find the analytical series solution of the Blasius problem (1.1) subject to the boundary condition

$$y(0) = 0, \quad y'(0) = \mu, \quad y''(0) = \kappa, \quad (2.1)$$

where the curvature κ of the solution is assumed to be known. It should be noted that in order to make the problem easy to be solved, we consider the one point boundary conditions in (2.1) instead of the two-point boundary conditions in (1.2).

First, for $y = y(x)$ the Blasius equation (1.1) becomes

$$(y'' + \alpha y y')' = \alpha (y')^2. \quad (2.2)$$

From the boundary conditions in (2.1), it follows that

$$y'' + \alpha y y' = \alpha \int_0^x \{y'(t)\}^2 dt + \kappa := A(x, y). \quad (2.3)$$

This can be represented by

$$\left(y' + \frac{\alpha}{2} y^2\right)' = A(x, y), \quad (2.4)$$

which implies

$$y' = \int_0^x A(t, y) dt - \frac{\alpha}{2} y^2 + \mu := B(x, y). \quad (2.5)$$

In the result, we have

$$y(x) = \int_0^x B(t, y) dt. \quad (2.6)$$

If we denote by y_n the n th iterate solution and substitute it into the right hand side of (2.6), we have an iteration formula

$$y_{n+1}(x) = \int_0^x B(t, y_n(t)) dt, \quad n \geq 0, \quad (2.7)$$

where

$$B(x, y_n(x)) = \int_0^x A(t, y_n(t)) dt - \frac{\alpha}{2} y_n(x)^2 + \mu. \quad (2.8)$$

From (2.3) and (2.7), the function $A(x, y_n)$ can be represented by

$$A(x, y_n(x)) = \alpha \int_0^x \{B(t, y_{n-1}(t))\}^2 dt + \kappa, \quad (2.9)$$

for $n \geq 1$ with $A(t, y_0) \equiv \kappa$. Referring to the boundary conditions in (2.1), we may take the initial solution as

$$y_0(x) = \frac{\kappa}{2}x^2 + \mu x. \quad (2.10)$$

The proposed method can be summarized by the following algorithm.

Algorithm A. We have the following steps.

Step 1. Set initial guesses

$$\begin{aligned} y(x) &:= y_0(x), \\ B(x, y) &:= y'_0(x). \end{aligned} \quad (2.11)$$

Step 2. For a large integer $L > 0$, perform the iteration (2.7)–(2.9) using symbolic computations

$$\begin{aligned} l &:= 0, \quad \text{while } (l := l + 1) \leq L, \\ A(x, y) &:= \alpha \int_0^x \{B(t, y(t))\}^2 dt + \kappa, \\ B(x, y) &:= \int_0^x A(t, y(t)) dt - \frac{\alpha}{2} y(x)^2 + \mu, \\ y(x) &:= \int_0^x B(t, y(t)) dt. \end{aligned} \quad (2.12)$$

It should be noted that by performing this algorithm, we can also obtain the approximates $B(x, y_n)$ to the velocity $y'(x) = B(x, y)$.

3. Analytical Solutions

Performing the above algorithm by using the symbolic calculation software *Mathematica*, we have the successive approximate solutions below

$$\begin{aligned} y_1(x) &= \frac{\kappa x^2}{2} - \frac{\alpha k^2 x^5}{120} + \left(x - \frac{\alpha k x^4}{24}\right) \mu, \\ y_2(x) &= \frac{\kappa x^2}{2} - \frac{\alpha k^2 x^5}{120} + \frac{11\alpha^2 k^3 x^8}{40320} - \frac{\alpha^3 k^4 x^{11}}{712800} + \left(x - \frac{\alpha k x^4}{24} + \frac{11\alpha^2 k^2 x^7}{5040} - \frac{\alpha^3 k^3 x^{10}}{64800}\right) \mu \\ &\quad + \left(\frac{\alpha^2 k x^6}{240} - \frac{\alpha^3 k^2 x^9}{24192}\right) \mu^2, \end{aligned}$$

$$\begin{aligned}
y_3(x) = & \frac{\kappa x^2}{2} - \frac{\alpha k^2 x^5}{120} + \frac{11\alpha^2 k^3 x^8}{40320} - \frac{5\alpha^3 k^4 x^{11}}{532224} + \frac{10033\alpha^4 k^5 x^{14}}{87178291200} - \frac{5449\alpha^5 k^6 x^{17}}{3908653056000} \\
& + \frac{83\alpha^6 k^7 x^{20}}{8935557120000} - \frac{\alpha^7 k^8 x^{23}}{49080898944000} \\
& + \left(x - \frac{\alpha k x^4}{24} + \frac{11\alpha^2 k^2 x^7}{5040} - \frac{5\alpha^3 k^3 x^{10}}{48384} \right. \\
& \quad \left. + \frac{10033\alpha^4 k^4 x^{13}}{6227020800} - \frac{5449\alpha^5 k^5 x^{16}}{229920768000} + \frac{83\alpha^6 k^6 x^{19}}{446777856000} - \frac{\alpha^7 k^7 x^{22}}{2133952128000} \right) \mu \\
& + \left(\frac{\alpha^2 k x^6}{240} - \frac{43\alpha^3 k^2 x^9}{120960} + \frac{1157\alpha^4 k^3 x^{12}}{159667200} \right. \\
& \quad \left. - \frac{1147\alpha^5 k^4 x^{15}}{7925299200} + \frac{14057\alpha^6 k^5 x^{18}}{10260214272000} - \frac{197\alpha^7 k^6 x^{21}}{49145564160000} \right) \mu^2 \\
& - \left(\frac{\alpha^3 k x^8}{2688} - \frac{23\alpha^4 k^2 x^{11}}{2217600} + \frac{967\alpha^5 k^3 x^{14}}{2641766400} - \frac{1829\alpha^6 k^4 x^{17}}{414554112000} + \frac{\alpha^7 k^5 x^{20}}{66189312000} \right) \mu^3 \\
& - \left(\frac{\alpha^5 k^2 x^{13}}{3294720} - \frac{17\alpha^6 k^3 x^{16}}{3251404800} + \frac{\alpha^7 k^4 x^{19}}{47259168768} \right) \mu^4.
\end{aligned} \tag{3.1}$$

For the case of $\alpha = 1/2$, we have

$$\begin{aligned}
y_1(x) = & \frac{\kappa x^2}{2} - \frac{k^2 x^5}{240} + \left(x - \frac{k x^4}{48} \right) \mu, \\
y_2(x) = & \frac{\kappa x^2}{2} - \frac{k^2 x^5}{240} + \frac{11k^3 x^8}{161280} - \frac{k^4 x^{11}}{5702400} + \left(x - \frac{k x^4}{48} + \frac{11k^2 x^7}{20160} - \frac{k^3 x^{10}}{518400} \right) \mu \\
& + \left(\frac{k x^6}{960} - \frac{k^2 x^9}{193536} \right) \mu^2, \\
y_3(x) = & \frac{\kappa x^2}{2} - \frac{k^2 x^5}{240} + \frac{11k^3 x^8}{161280} - \frac{5k^4 x^{11}}{4257792} + \frac{10033k^5 x^{14}}{1394852659200} - \frac{5449k^6 x^{17}}{125076897792000} \\
& + \frac{83k^7 x^{20}}{571875655680000} - \frac{k^8 x^{23}}{6282355064832000} \\
& + \left(x - \frac{k x^4}{48} + \frac{11k^2 x^7}{20160} - \frac{5k^3 x^{10}}{387072} + \frac{10033k^4 x^{13}}{99632332800} \right. \\
& \quad \left. - \frac{5449k^5 x^{16}}{7357464576000} + \frac{83k^6 x^{19}}{28593782784000} - \frac{k^7 x^{22}}{273145872384000} \right) \mu
\end{aligned}$$

$$\begin{aligned}
& + \left(\frac{kx^6}{960} - \frac{43k^2x^9}{967680} + \frac{1157k^3x^{12}}{2554675200} - \frac{1147k^4x^{15}}{253609574400} \right. \\
& \quad \left. + \frac{14057k^5x^{18}}{656653713408000} - \frac{197k^6x^{21}}{6290632212480000} \right) \mu^2 \\
& - \left(\frac{kx^8}{21504} - \frac{23k^2x^{11}}{35481600} + \frac{967k^3x^{14}}{84536524800} - \frac{1829k^4x^{17}}{26531463168000} + \frac{k^5x^{20}}{8472231936000} \right) \mu^3 \\
& - \left(\frac{k^2x^{13}}{105431040} - \frac{17k^3x^{16}}{208089907200} + \frac{k^4x^{19}}{6049173602304} \right) \mu^4.
\end{aligned} \tag{3.2}$$

One can see that the result is consistent with the known series solution [11, 12].

In particular, when $\alpha = 1$ and $\mu = 0$, it follows that

$$\begin{aligned}
y_1(x) &= \frac{\kappa x^2}{2} - \frac{k^2 x^5}{120}, \\
y_2(x) &= \frac{\kappa x^2}{2} - \frac{k^2 x^5}{120} + \frac{11k^3 x^8}{40320} - \frac{k^4 x^{11}}{712800}, \\
y_3(x) &= \frac{\kappa x^2}{2} - \frac{k^2 x^5}{120} + \frac{11k^3 x^8}{40320} - \frac{5k^4 x^{11}}{532224} + \frac{10033k^5 x^{14}}{87178291200} - \frac{5449k^6 x^{17}}{3908653056000} \\
& \quad + \frac{83k^7 x^{20}}{8935557120000} - \frac{k^8 x^{23}}{49080898944000}, \\
y_4(x) &= \frac{\kappa x^2}{2} - \frac{k^2 x^5}{120} + \frac{11k^3 x^8}{40320} - \frac{5k^4 x^{11}}{532224} + \frac{9299k^5 x^{14}}{29059430400} - \frac{2173649k^6 x^{17}}{355687428096000} \\
& \quad + \frac{13722337k^7 x^{20}}{115852476579840000} - \frac{27184438601k^8 x^{23}}{12926008369442488320000} \\
& \quad + \frac{12320831753849k^9 x^{26}}{403291461126605635584000000} \\
& \quad + \dots - \frac{k^{16} x^{47}}{463172433275878342410240000000}.
\end{aligned} \tag{3.3}$$

In this case, $\kappa \approx 0.4695999883$. For comparison, we refer to another analytical solution obtained by the Adomian decomposition method as follows:

$$\begin{aligned}
y_1^A(x) &= \frac{\kappa x^2}{2} - \frac{k^2 x^5}{120}, \\
y_2^A(x) &= \frac{\kappa x^2}{2} - \frac{k^2 x^5}{120} + \frac{11k^3 x^8}{40320}, \\
y_3^A(x) &= \frac{\kappa x^2}{2} - \frac{k^2 x^5}{120} + \frac{11k^3 x^8}{40320} - \frac{5k^4 x^{11}}{532224}, \\
y_4^A(x) &= \frac{\kappa x^2}{2} - \frac{k^2 x^5}{120} + \frac{11k^3 x^8}{40320} - \frac{5k^4 x^{11}}{532224} + \frac{9299k^5 x^{14}}{29059430400}, \\
y_5^A(x) &= \frac{\kappa x^2}{2} - \frac{k^2 x^5}{120} + \frac{11k^3 x^8}{40320} - \frac{5k^4 x^{11}}{532224} + \frac{9299k^5 x^{14}}{29059430400} - \frac{1272379k^6 x^{17}}{118562476032000}.
\end{aligned} \tag{3.4}$$

This solution is based on

$$y_n^A(x) = y_0(x) + \sum_{k=1}^n u_k(x), \quad n = 1, 2, \dots, \quad (3.5)$$

with

$$u_{k+1}(x) = -L^{-1}(A_k(x)), \quad k = 0, 1, 2, \dots, \quad (3.6)$$

$u_0(x) = y_0(x)$, and Adomian polynomial $A_k(x)$ generated by the formula [6]

$$A_k(x) = \frac{1}{k!} \left[\frac{d^k}{d\lambda^k} \left(\sum_{j=0}^k \lambda^j u_j(x) \right) \left(\sum_{j=0}^k \lambda^j u_j''(x) \right) \right]_{\lambda=0}, \quad (3.7)$$

where L^{-1} is an inverse operator of $L = d^3/dx^3$. Comparing the formulas in (3.3) and (3.4), one can see that the presented analytical solution $y_n(x)$ has more terms than $y_n^A(x)$ in each n th iteration. In other words,

$$y_n(x) = y_n^A(x) + \sum_{j=1}^{2^n - n - 1} d_j x^{3(n+j)+2}, \quad (3.8)$$

for any integer $n \geq 2$. In practice, Figure 1 depicts that the presented solutions $y_n(x)$ and their derivatives $y_n'(x)$, $n = 1, 3, 5$ approximate exact ones better than $y_n^A(x)$ and $(y_n^A)'(x)$. Therein, we chose the initial solution $y_0(x) = (\kappa/2)x^2$ as given in (2.10) and took a numerical solution for the exact solution which is denoted by $y^*(x)$. Moreover, Table 1 includes numerical results of the errors and the CPU times spent in computations for the presented solution $y_n(x)$ compared with those of $y_n^A(x)$. The L_∞ error indicates the maximum error for the 50 nodes selected in the interval $(0, \rho)$, where ρ is a radius of convergence of the series solution given in the literature [2, 15]. In fact, $\rho \approx 4.02$ for $\alpha = 1$ and $\rho \approx 5.69$ for $\alpha = 1/2$. The L_2 error means $\|y_n - y^*\|_2$ over the same interval.

By the numerical performance, we can surmise the convergence of the algorithm proposed in this work, and the rate of convergence is better than that of the Adomian decomposition method though it spends more CPU time as shown above. In addition, for example, for the case of $\alpha = 1$ and $\mu = 0$, we may guess that the presented method has the same radius of convergence, $\rho = 4.0234644935$ as the well known Blasius' series [2] as follows:

$$y(x) = \sum_{k=0}^{\infty} (-1)^k p_k \frac{\kappa^{k+1}}{(3k+2)!} x^{3k+2}, \quad (3.9)$$

where $p_0 = 1$ and

$$p_k = \sum_{j=0}^{k-1} \binom{3k-1}{3j} p_j p_{k-j-1}, \quad k \geq 1. \quad (3.10)$$

Theoretical convergence analysis with extended application of the presented method to more general problems is left for further works.

Figure 1: Graphs of the presented solutions $y_n(x)$ and their derivatives $y'_n(x)$ in (a) and those of Adomian's solutions $y_n^A(x)$ and their derivatives $(y_n^A)'(x)$ in (b).

Table 1: Comparison of the numerical results obtained by the presented method and the Adomian's decomposition method for $\alpha = 1, 1/2$ with $\mu = 0$.

	n	Presented solution, $y_n(x)$			Adomian's solution, $y_n^A(x)$		
		L_∞ error	L_2 error	CPU time (sec)	L_∞ error	L_2 error	CPU time (sec)
$\alpha = 1$	2	0.690	0.318	0.14	0.995	0.445	0.09
	4	0.300	0.110	1.03	1.080	0.382	0.37
	6	0.078	0.025	6.37	1.110	0.335	1.48
	8	0.013	0.004	26.5	1.118	0.299	5.74
$\alpha = 1/2$	2	0.976	0.534	0.13	1.407	0.748	0.11
	4	0.424	0.185	1.01	1.527	0.643	0.41
	6	0.111	0.041	6.71	1.570	0.564	1.55
	8	0.019	0.006	27.4	1.581	0.504	5.93

Acknowledgments

The author would like to show his gratitude to the Department of Mathematics at the University of British Columbia, where the author worked as a visiting scholar for a year. Particularly, the author is heartily thankful to Professor Anthony Peirce for his favor and kind assistance. In addition, the author shows his sincere gratitude to reviewers for their helpful comments and valuable suggestions on the first draft of this paper. This research was supported by Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education, Science, and Technology (20110006106).

References

- [1] J. P. Boyd, "Padé approximant algorithm for solving nonlinear ordinary differential equation boundary value problems on an unbounded domain," *Computers in Physics*, vol. 11, pp. 299–303, 1997.

- [2] J. P. Boyd, "The Blasius function in the complex plane," *Experimental Mathematics*, vol. 8, no. 4, pp. 381–394, 1999.
- [3] J. P. Boyd, "The Blasius function: computations before computers, the value of tricks, undergraduate projects, and open research problems," *SIAM Review*, vol. 50, no. 4, pp. 791–804, 2008.
- [4] G. Adomian, "A review of the decomposition method in applied mathematics," *Journal of Mathematical Analysis and Applications*, vol. 135, no. 2, pp. 501–544, 1988.
- [5] G. Adomian, "Solution of the Thomas-Fermi equation," *Applied Mathematics Letters*, vol. 11, no. 3, pp. 131–133, 1998.
- [6] E. Alizadeh, M. Farhadi, K. Sedighi, H. R. Ebrahimi-Kebria, and A. Ghafourian, "Solution of the Falkner-Skan equation for wedge by Adomian Decomposition Method," *Communications in Nonlinear Science and Numerical Simulation*, vol. 14, no. 3, pp. 724–733, 2009.
- [7] J. He, "Approximate analytical solution of Blasius' equation," *Communications in Nonlinear Science and Numerical Simulation*, vol. 3, no. 4, pp. 260–263, 1998.
- [8] Ji-Huan He, "A review on some new recently developed nonlinear analytical techniques," *International Journal of Nonlinear Sciences and Numerical Simulation*, vol. 1, no. 1, pp. 51–70, 2000.
- [9] J. H. He, "A simple perturbation approach to Blasius equation," *Applied Mathematics and Computation*, vol. 140, no. 2-3, pp. 217–222, 2003.
- [10] J. Y. Parlange, R. D. Braddock, and G. Sander, "Analytical approximations to the solution of the Blasius equation," *Acta Mechanica*, vol. 38, no. 1-2, pp. 119–125, 1981.
- [11] A. M. Wazwaz, "The variational iteration method for solving two forms of Blasius equation on a half-infinite domain," *Applied Mathematics and Computation*, vol. 188, no. 1, pp. 485–491, 2007.
- [12] F. M. Allan and M. I. Syam, "On the analytic solutions of the nonhomogeneous Blasius problem," *Journal of Computational and Applied Mathematics*, vol. 182, no. 2, pp. 362–371, 2005.
- [13] S. J. Liao, "A uniformly valid analytic solution of two-dimensional viscous flow over a semi-infinite flat plate," *Journal of Fluid Mechanics*, vol. 385, pp. 101–128, 1999.
- [14] S. J. Liao, "An explicit, totally analytic approximate solution for Blasius' viscous flow problems," *International Journal of Non-Linear Mechanics*, vol. 34, no. 4, pp. 759–778, 1999.
- [15] S. Finch, "Prandtl-Blasius flow," 2008, <http://algo.inria.fr/csolve/bla.pdf>.