Research Article

The System of Mixed Equilibrium Problems for Quasi-Nonexpansive Mappings in Hilbert Spaces

Rabian Wangkeeree^{1,2} and Panatda Boonman¹

¹ Department of Mathematics, Faculty of Science, Naresuan University, Phitsanulok 65000, Thailand

² Centre of Excellence in Mathematics, CHE, Si Ayutthaya Road, Bangkok 10400, Thailand

Correspondence should be addressed to Rabian Wangkeeree, rabianw@nu.ac.th

Received 19 February 2012; Accepted 23 March 2012

Academic Editor: Rudong Chen

Copyright © 2012 R. Wangkeeree and P. Boonman. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We first introduce the iterative procedure to approximate a common element of the fixed-point set of two quasinonexpansive mappings and the solution set of the system of mixed equilibrium problem (SMEP) in a real Hilbert space. Next, we prove the weak convergence for the given iterative scheme under certain assumptions. Finally, we apply our results to approximate a common element of the set of common fixed points of asymptotic nonspreading mapping and asymptotic TJ mapping and the solution set of SMEP in a real Hilbert space.

1. Introduction

Let H be a real Hilbert space with inner product $\langle \cdot, \cdot \rangle$ and norm $\| \cdot \|$, let C be a nonempty closed convex subset of H , and let T be a mapping of C into H , then $T : C \rightarrow H$ is said to be nonexpansive if $\|Tx - Ty\| \leq \|x - y\|$ for all $x, y \in C$. A mapping $T : C \rightarrow H$ is said to be quasinonexpansive if $\|Tx - y\| \leq \|x - y\|$ for all $x \in C$ and $y \in F(T) := \{x \in C : Tx = x\}$. It is well known that the set $F(T)$ of fixed points of a quasi-nonexpansive mapping T is a closed and convex set [1]. A mapping $T : C \rightarrow C$ is said to be firmly nonexpansive [2] if

$$\|Tx - Ty\|^2 \leq \langle x - y, Tx - Ty \rangle, \quad (1.1)$$

for all $x, y \in C$, and it is an important example of nonexpansive mappings in a Hilbert space.

Let $\varphi : C \rightarrow \mathbb{R}$ be a real-valued function, and let $F : C \times C \rightarrow \mathbb{R}$ be an equilibrium bifunction, that is, $F(u, u) = 0$ for each $u \in C$. The mixed equilibrium problem is to find $x \in C$ such that

$$F(x, y) + \varphi(y) - \varphi(x) \geq 0 \quad \forall y \in C. \quad (1.2)$$

Denote the set of solution of (1.2) by $\text{MEP}(F, \varphi)$. In particular, if $\varphi = 0$, this problem reduces to the equilibrium problem, which is to find $x \in C$ such that

$$F(x, y) \geq 0 \quad \forall y \in C. \quad (1.3)$$

The set of solution of (1.3) is denoted by $\text{EP}(F)$.

The problem (1.2) is very general in the sense that it includes, as special cases, optimization problems, variational inequalities, Min-Max problems, the Nash equilibrium problems in noncooperative games, and others; see, for example, Blum and Oettli [3] and Moudafi [4]. Numerous problems in physics, optimization, and economics reduce to find a solution of (1.3).

Let $F_1, F_2 : C \times C \rightarrow \mathbb{R}$ be two monotone bifunctions and $\mu > 0$ is constant. In 2009, Moudafi [5] introduced an alternating algorithm for approximating a solution of the system of equilibrium problems, finding $(x', y') \in C \times C$ such that

$$(\text{SEP}) \begin{cases} F_1(x', z) + \frac{1}{\mu} \langle y' - x', x' - z \rangle \geq 0, & \forall z \in C, \\ F_2(y', z) + \frac{1}{\mu} \langle x' - y', y' - z \rangle \geq 0, & \forall z \in C. \end{cases} \quad (1.4)$$

For such mappings F_1 and F_2 and two given positive constants $\lambda, \mu > 0$, Plubtieng and Sombut [6] considered the following system of mixed equilibrium problem, finding $(x', y') \in C \times C$ such that

$$(\text{SMEP}) \begin{cases} F_1(x', z) + \varphi(z) - \varphi(x') + \frac{1}{\lambda} \langle y' - x', x' - z \rangle \geq 0, & \forall z \in C, \\ F_2(y', z) + \varphi(z) - \varphi(y') + \frac{1}{\mu} \langle x' - y', y' - z \rangle \geq 0, & \forall z \in C. \end{cases} \quad (1.5)$$

In particular, if $\lambda = \mu$ and $\varphi \equiv 0$, then problem (SMEP) reduces to (SEP). Furthermore, Plubtieng and Sombut [6] introduced the following iterative procedure to approximate a common element of the fixed-point set of a quasi-nonexpansive mapping T and the solution set of (SMEP) in a Hilbert space H . Let $\{x_n\}$, $\{y_n\}$, and $\{u_n\}$ be given by

$$\begin{aligned} & x_1 \in C \text{ chosen arbitrary,} \\ & u_n \in C, \quad F_2(u_n, z) + \varphi(z) - \varphi(u_n) + \frac{1}{\mu} \langle z - u_n, u_n - x_n \rangle \geq 0, \quad \forall z \in C, \\ & y_n \in C, \quad F_1(y_n, z) + \varphi(z) - \varphi(y_n) + \frac{1}{\lambda} \langle z - y_n, y_n - u_n \rangle \geq 0, \quad \forall z \in C, \\ & x_{n+1} = \alpha_n x_n + (1 - \alpha_n) T y_n, \quad \forall n \in \mathbb{N}, \end{aligned} \quad (1.6)$$

where $\{\alpha_n\} \subset [a, b]$ for some $a, b \in (0, 1)$ and satisfying appropriate conditions. The weak convergence theorems are obtained in a real Hilbert space.

On the other hand, in 1953, Mann [7] introduced the following iterative procedure to approximate a fixed point of a nonexpansive mapping T in a Hilbert space H :

$$x_{n+1} = \alpha_n x_n + (1 - \alpha_n) T x_n, \quad \forall n \in \mathbb{N}, \quad (1.7)$$

where the initial point x_0 is taken in C arbitrarily, and $\{\alpha_n\}$ is a sequence in $[0, 1]$.

For two nonexpansive mappings T_1, T_2 of C into itself, Moudafi [4] studied weak convergence theorems in the following iterative process:

$$\begin{aligned} x_0 &\in C \text{ chosen arbitrary,} \\ x_{n+1} &= \alpha_n x_n + (1 - \alpha_n)(\beta_n T_1 x_n + (1 - \beta_n) T_2 x_n), \end{aligned} \quad (1.8)$$

for all $n \in \mathbb{N}$, where $\{\alpha_n\}$ and $\{\beta_n\}$ are appropriate sequences in $[0, 1]$ and $F(T_1) \cap F(T_2) \neq \emptyset$. Recently, Iemoto and Takahashi [8] also considered this iterative procedure for T_1 is a nonexpansive mapping and $T_2 : C \rightarrow C$ is a nonspreading mapping. Very recently, Kim [9] studied the weak and strong convergence for the Moudafi's iterative scheme (1.8) of two quasi-nonexpansive mappings.

In this paper, inspired and motivated by Plubtieng and Sombut [6], Moudafi [4], Iemoto and Takahashi [8], and Kim [9], we first introduce the iterative procedure to approximate a common element of the common fixed point set of two quasi-nonexpansive mappings and the solution set of SMEP in a real Hilbert space. Next, we prove the weak convergence theorem for the given iterative scheme under certain assumptions. Finally, we apply our results to approximate a common element of the set of common fixed point of asymptotic nonspreading mapping and asymptotic TJ mapping and the solution set of SMEP in a real Hilbert space.

2. Preliminaries

Throughout this paper, let $\mathbb{N}$ be the set of positive integers, and let $\mathbb{R}$ be the set of real numbers. Let H be a real Hilbert space with inner product $\langle \cdot, \cdot \rangle$ and norm $\| \cdot \|$, respectively, and let C be a closed convex subset of H . We denote the strong convergence and the weak convergence of $\{x_n\}$ to $x \in H$ by $x_n \rightarrow x$ and $x_n \rightharpoonup x$, respectively.

From [10], for each $x, y \in H$ and $\lambda \in [0, 1]$, we have

$$\| \lambda x + (1 - \lambda)y \|^2 = \lambda \|x\|^2 + (1 - \lambda) \|y\|^2 - \lambda(1 - \lambda) \|x - y\|^2. \quad (2.1)$$

For every point $x \in H$, there exists a unique nearest point in C , denoted by $P_C x$, such that

$$\|x - P_C x\| \leq \|x - y\| \quad \forall y \in C. \quad (2.2)$$

P_C is called the metric projection of H onto C . It is well know that P_C is a nonexpansive mapping of H onto C and satisfies

$$\|P_C x - P_C y\|^2 \leq \langle P_C x - P_C y, x - y \rangle \quad \forall x, y \in H. \quad (2.3)$$

Moreover, $P_C x$ is characterized by the following properties: $P_C x \in C$,

$$\begin{aligned} \langle x - P_C x, P_C y - y \rangle &\geq 0, \\ \|x - y\|^2 &\geq \|x - P_C x\|^2 + \|y - P_C y\|^2 \quad \forall x \in H, y \in C. \end{aligned} \quad (2.4)$$

Further, for all $x \in H$ and $y \in C$, $y = P_C x$ if and only if $\langle x - y, y - z \rangle \geq 0$, for all $z \in C$.

Lemma 2.1 (see [11]). *Let C be a nonempty closed convex subset of a real Hilbert space H . Let P be the metric projection of H onto C , and let $\{x_n\}_{n \in \mathbb{N}}$ be in H . If*

$$\|x_{n+1} - u\| \leq \|x_n - u\|, \quad (2.5)$$

for all $u \in C$ and $n \in \mathbb{N}$. Then, $\{P_C x_n\}$ converges strongly to an element of C .

Theorem 2.2 (Opial's theorem, [10]). *Let H be a real Hilbert space, and suppose that $x_n \rightharpoonup x$, then*

$$\liminf_{n \rightarrow \infty} \|x_n - x\| < \liminf_{n \rightarrow \infty} \|x_n - y\|, \quad (2.6)$$

for all $y \in H$ with $x \neq y$.

All Hilbert space and l^p ($1 < p < \infty$) satisfy Opial's condition, while L^p with $1 < p \neq 2 < \infty$ do not.

For solving the mixed equilibrium problem for an equilibrium bifunction $F : C \times C \rightarrow \mathbb{R}$, let us assume that F satisfies the following conditions:

- (A1) $F(x, x) = 0$ for all $x \in C$,
- (A2) F is monotone, that is, $F(x, y) + F(y, x) \leq 0$ for all $x, y \in C$,
- (A3) for each $y \in C$, $x \mapsto F(x, y)$ is weakly upper semicontinuous,
- (A4) for each $x \in C$, $y \mapsto F(x, y)$ is convex and semicontinuous.

The following lemma appears implicitly in [3, 12].

Lemma 2.3 (see [3]). *Let C be a nonempty closed convex subset of H , and let $F : C \times C \rightarrow \mathbb{R}$ be a bifunction satisfying (A1)–(A4). Let $r > 0$ and $x \in H$, then there exists $z \in C$ such that*

$$F(z, y) + \frac{1}{r} \langle y - z, z - x \rangle \geq 0 \quad \forall y \in C. \quad (2.7)$$

The following lemma was also given in [12].

Lemma 2.4 (see [12]). *Let C be a nonempty closed convex subset of H and let $F : C \times C \rightarrow \mathbb{R}$ be a bifunction satisfying (A1)–(A4), then, for any $r > 0$ and $x \in H$, define a mapping $T_r x : H \rightarrow C$ as follows:*

$$T_r(x) = \left\{ z \in C : F(z, y) + \frac{1}{r} \langle y - z, z - x \rangle \geq 0, \quad \forall y \in C \right\}, \quad (2.8)$$

for all $z \in H, r \in \mathbb{R}$. Then the following hold:

- (i) T_r is single valued,
- (ii) T_r is firmly nonexpansive, that is,

$$\|T_r x - T_r y\|^2 \leq \langle T_r x - T_r y, x - y \rangle, \quad \forall x, y \in H; \quad (2.9)$$

- (iii) $F(T_r) = \text{EP}(F)$,
- (iv) $\text{EP}(F)$ is closed and convex.

We note that Lemma 2.4 is equivalent to the following lemma.

Lemma 2.5 (see [6]). *Let C be a nonempty closed convex subset of a real Hilbert space H . Let $F : C \times C \rightarrow \mathbb{R}$ be an equilibrium bifunction satisfying (A1)–(A4) and let $\varphi : C \rightarrow \mathbb{R}$ be a lower semicontinuous and convex functional. For each $r > 0$ and $x \in H$, define a mapping*

$$S_r(x) = \left\{ y \in C : F(y, z) + \varphi(z) - \varphi(y) + \frac{1}{r} \langle y - x, z - y \rangle \geq 0, \forall z \in C \right\}, \quad \forall x \in H. \quad (2.10)$$

Then, the following results hold:

- (i) for each $x \in C$, $S_r(x) \neq \emptyset$,
- (ii) S_r is single valued,
- (iii) S_r is firmly nonexpansive, that is, for any $x, y \in H$,

$$\|S_r(x) - S_r(y)\|^2 \leq \langle S_r(x) - S_r(y), x - y \rangle, \quad (2.11)$$

- (iv) $F(S_r) = \text{MEP}(F, \varphi)$,
- (v) $\text{MEP}(F, \varphi)$ is closed and convex.

3. Main Results

In this section, we prove the weak convergence for approximating a common element of the common fixed point set of two quasi-nonexpansive mappings and the solution set of the system of mixed equilibrium problems in a Hilbert space.

To begin with, let us state and prove the following characterizations of the solution set of GMEP.

Lemma 3.1. *Let C be a closed convex subset of a real Hilbert space H . Let F_1 and F_2 be two mappings from $C \times C \rightarrow \mathbb{R}$ satisfying (A1)–(A4), and let $S_{1,\lambda}$ and $S_{2,\mu}$ be defined as in Lemma 2.5 associated to F_1 and F_2 , respectively. For given $x', y' \in C$, (x', y') is a solution of problem (1.5) if and only if x' is a fixed point of the mapping $G : C \rightarrow C$ defined by*

$$G(x) = S_{1,\lambda}(S_{2,\mu}x), \quad \forall x \in C, \quad (3.1)$$

where $y' = S_{2,\mu}x'$.

$$\begin{aligned}
(x', y') \text{ is a solution of problem (1.5)} &\iff \begin{cases} F_1(x', z) + \varphi(z) - \varphi(x') + \frac{1}{\lambda} \langle y' - x', x' - z \rangle \geq 0 \\ \forall z \in C, \\ F_2(y', z) + \varphi(z) - \varphi(y') + \frac{1}{\mu} \langle x' - y', y' - z \rangle \geq 0 \\ \forall z \in C, \end{cases} \\
&\iff x' = S_{1,\lambda} y' \quad y' = S_{2,\mu} x', \\
&\iff x' = S_{1,\lambda}(S_{2,\mu} x'), \\
&\iff x' = Gx'.
\end{aligned} \tag{3.2}$$
☐

Theorem 3.2. *Let C be a nonempty closed convex subset of a real Hilbert space H . Let F_1 and F_2 be two bifunctions from $C \times C \rightarrow \mathbb{R}$ satisfying (A1)–(A4). Let $r, \lambda > 0$ and $S_{1,r}$ and $S_{2,\lambda}$ be defined as in Lemma 2.5 associated to F_1 and F_2 , respectively. Let $T_i : C \rightarrow C$, $i = 1, 2$, be two quasi-nonexpansive mappings such that $I - T_i$ are demiclosed at zero, that is, if $\{w_n\} \subset C$, $w_n \rightharpoonup w$, and $(I - T_i)w_n \rightarrow 0$, then $w \in F(T_i)$, with $F(T_1) \cap F(T_2) \cap \Omega \neq \emptyset$. Let the sequences $\{x_n\}$, $\{y_n\}$, and $\{u_n\}$, be given by*

where $\{a_n\}, \{b_n\} \subset [a, b]$ for some $a, b \in (0, 1)$, and satisfy

then $x_n \rightharpoonup \bar{x} := \lim_{n \rightarrow \infty} P_{F(T_1) \cap F(T_2) \cap \Omega} x_n$ and $(\bar{x}, \bar{y})$ is a solution of problem (1.5), where $\bar{y} = S_{2,\lambda} \bar{x}$.

Putting $y^* = S_{2,\lambda}x^*$, $y_n = S_{1,r}u_n$, and $u_n = S_{2,\lambda}x_n$, we have

$$\|y_n - x^*\| = \|S_{1,r}u_n - S_{1,r}y^*\| \leq \|u_n - y^*\| = \|S_{2,\lambda}x_n - S_{2,\lambda}x^*\| \leq \|x_n - x^*\|. \quad (3.5)$$

Next, we prove that

$$\lim_{n \rightarrow \infty} \|x_n - x^*\| \text{ exists.} \quad (3.6)$$

Since T_1 and T_2 are quasi-nonexpansive, we obtain that

$$\begin{aligned} \|b_n T_1 y_n + (1 - b_n) T_2 y_n - x^*\|^2 &= \|b_n (T_1 y_n - x^*) + (1 - b_n) (T_2 y_n - x^*)\|^2 \\ &= b_n \|T_1 y_n - x^*\|^2 + (1 - b_n) \|T_2 y_n - x^*\|^2 \\ &\quad - b_n (1 - b_n) \|T_1 y_n - T_2 y_n\|^2 \\ &\leq b_n \|T_1 y_n - x^*\|^2 + (1 - b_n) \|T_2 y_n - x^*\|^2 \\ &\leq b_n \|y_n - x^*\|^2 + (1 - b_n) \|y_n - x^*\|^2 \\ &= \|y_n - x^*\|^2 \\ &\leq \|x_n - x^*\|^2, \end{aligned} \quad (3.7)$$

which gives that

$$\begin{aligned} \|x_{n+1} - x^*\|^2 &= \|a_n x_n + (1 - a_n) (b_n T_1 y_n + (1 - b_n) T_2 y_n) - x^*\|^2 \\ &= \|a_n (x_n - x^*) + (1 - a_n) (b_n T_1 y_n + (1 - b_n) T_2 y_n - x^*)\|^2 \\ &= a_n \|x_n - x^*\|^2 + (1 - a_n) \|b_n T_1 y_n + (1 - b_n) T_2 y_n - x^*\|^2 \\ &\quad - a_n (1 - a_n) \|x_n - (b_n T_1 y_n + (1 - b_n) T_2 y_n)\|^2 \\ &\leq a_n \|x_n - x^*\|^2 + (1 - a_n) \|x_n - x^*\|^2 \\ &\quad - a_n (1 - a_n) \|x_n - (b_n T_1 y_n + (1 - b_n) T_2 y_n)\|^2 \\ &= \|x_n - x^*\|^2 - a_n (1 - a_n) \|x_n - (b_n T_1 y_n + (1 - b_n) T_2 y_n)\|^2 \\ &\leq \|x_n - x^*\|^2. \end{aligned} \quad (3.8)$$

Hence, $\{\|x_n - x^*\|\}$ is a nonincreasing sequence, and hence, $\lim_{n \rightarrow \infty} \|x_n - x^*\|$ exists. This implies that $\{x_n\}$, $\{y_n\}$, $\{u_n\}$, $\{T_1 y_n\}$, and $\{T_2 y_n\}$ are bounded. From (3.8), we have

$$a_n (1 - a_n) \|x_n - (b_n T_1 y_n + (1 - b_n) T_2 y_n)\|^2 \leq \|x_n - x^*\|^2 - \|x_{n+1} - x^*\|^2. \quad (3.9)$$

Since $\liminf_{n \rightarrow \infty} a_n (1 - a_n) > 0$, this implies that

$$\lim_{n \rightarrow \infty} \|x_n - (b_n T_1 y_n + (1 - b_n) T_2 y_n)\| = 0. \quad (3.10)$$

Furthermore, since $0 < a \leq a_n \leq b < 1$, we have

$$\begin{aligned}
 \|x_{n+1} - x_n\| &= \|a_n x_n + (1 - a_n)(b_n T_1 y_n + (1 - b_n) T_2 y_n) - x_n\| \\
 &= \|(1 - a_n)(b_n T_1 y_n + (1 - b_n) T_2 y_n) - (1 - a_n) x_n\| \\
 &= (1 - a_n) \|b_n T_1 y_n + (1 - b_n) T_2 y_n - x_n\| \\
 &\leq (1 - a) \|b_n T_1 y_n + (1 - b_n) T_2 y_n - x_n\|.
 \end{aligned} \tag{3.11}$$

From (3.10), we conclude that

$$\lim_{n \rightarrow \infty} \|x_{n+1} - x_n\| = 0. \tag{3.12}$$

From (3.7), we have

$$\begin{aligned}
 b_n(1 - b_n) \|T_1 y_n - T_2 y_n\|^2 &= b_n \|T_1 y_n - x^*\|^2 + (1 - b_n) \|T_2 y_n - x^*\|^2 \\
 &\quad - \|b_n T_1 y_n + (1 - b_n) T_2 y_n - x^*\|^2 \\
 &\leq b_n \|y_n - x^*\|^2 + (1 - b_n) \|y_n - x^*\|^2 \\
 &\quad - \|b_n T_1 y_n + (1 - b_n) T_2 y_n - x^*\|^2 \\
 &= \|y_n - x^*\|^2 - \|b_n T_1 y_n + (1 - b_n) T_2 y_n - x^*\|^2 \\
 &\leq \|x_n - x^*\|^2 - \|b_n T_1 y_n + (1 - b_n) T_2 y_n - x^*\|^2 \\
 &= (\|x_n - x^*\| - \|b_n T_1 y_n + (1 - b_n) T_2 y_n - x^*\|) \\
 &\quad \times (\|x_n - x^*\| + \|b_n T_1 y_n + (1 - b_n) T_2 y_n - x^*\|) \\
 &\leq M(\|x_n - x^*\| - \|b_n T_1 y_n + (1 - b_n) T_2 y_n - x^*\|) \\
 &\leq M(\|x_n - (b_n T_1 y_n + (1 - b_n) T_2 y_n)\|),
 \end{aligned} \tag{3.13}$$

where M is a constant satisfying $M \geq \sup_{n \geq 1} [\|x_n - x^*\| + \|b_n T_1 y_n + (1 - b_n) T_2 y_n - x^*\|]$. Again from (3.10), we conclude that

$$\lim_{n \rightarrow \infty} b_n(1 - b_n) \|T_1 y_n - T_2 y_n\| = 0. \tag{3.14}$$

Using $\liminf_{n \rightarrow \infty} b_n(1 - b_n) > 0$, we have

$$\lim_{n \rightarrow \infty} \|T_1 y_n - T_2 y_n\| = 0. \tag{3.15}$$

Now, we prove that

$$\lim_{n \rightarrow \infty} \|x_{n+1} - T_1 y_n\| = 0, \quad \lim_{n \rightarrow \infty} \|x_{n+1} - T_2 y_n\| = 0. \tag{3.16}$$

We observe that

$$\begin{aligned}
\|x_{n+1} - T_1 y_n\| &= \|a_n x_n + (1 - a_n)(b_n T_1 y_n + (1 - b_n)T_2 y_n) - T_1 y_n\| \\
&= \|a_n x_n + (1 - a_n)(b_n T_1 y_n + (1 - b_n)T_2 y_n) - (a_n + (1 - a_n))T_1 y_n\| \\
&= \|a_n(x_n - T_1 y_n) + (1 - a_n)(b_n T_1 y_n + (1 - b_n)T_2 y_n - T_1 y_n)\| \\
&= \|a_n(x_n - T_1 y_n) + (1 - a_n)(b_n T_1 y_n + T_2 y_n - b_n T_2 y_n - T_1 y_n)\| \\
&= \|a_n(x_n - T_1 y_n) + (1 - a_n)((1 - b_n)T_2 y_n - (1 - b_n)T_1 y_n)\| \\
&= \|a_n(x_n - T_1 y_n) + (1 - a_n)(1 - b_n)(T_2 y_n - T_1 y_n)\| \\
&\leq a_n \|x_n - T_1 y_n\| + (1 - a_n)(1 - b_n) \|T_2 y_n - T_1 y_n\| \\
&= a_n \|x_n - x_{n+1} + x_{n+1} - T_1 y_n\| + (1 - a_n)(1 - b_n) \|T_2 y_n - T_1 y_n\| \\
&\leq a_n \|x_n - x_{n+1}\| + a_n \|x_{n+1} - T_1 y_n\| + (1 - a_n)(1 - b_n) \|T_2 y_n - T_1 y_n\|,
\end{aligned} \tag{3.17}$$

which gives that

$$(1 - a_n) \|x_{n+1} - T_1 y_n\| \leq a_n \|x_n - x_{n+1}\| + (1 - a_n)(1 - b_n) \|T_2 y_n - T_1 y_n\|. \tag{3.18}$$

Since $0 < a \leq a_n \leq b < 1$, we have

$$\begin{aligned}
a_n(1 - a_n) \|x_{n+1} - T_1 y_n\| &\leq (1 - a_n) \|x_{n+1} - T_1 y_n\| \\
&\leq a_n \|x_n - x_{n+1}\| + (1 - a_n)(1 - b_n) \|T_2 y_n - T_1 y_n\|.
\end{aligned} \tag{3.19}$$

Using (3.12) and (3.15), we conclude that

$$\lim_{n \rightarrow \infty} a_n(1 - a_n) \|x_{n+1} - T_1 y_n\| = 0, \tag{3.20}$$

which gives that

$$\lim_{n \rightarrow \infty} \|x_{n+1} - T_1 y_n\| = 0, \tag{3.21}$$

since $\liminf_{n \rightarrow \infty} a_n(1 - a_n) > 0$. Similarly, we have

$$\begin{aligned}
\|x_{n+1} - T_2 y_n\| &= \|a_n x_n + (1 - a_n)(b_n T_1 y_n + (1 - b_n)T_2 y_n) - T_2 y_n\| \\
&= \|a_n(x_n - T_2 y_n) + (1 - a_n)(b_n T_1 y_n + (1 - b_n)T_2 y_n - T_2 y_n)\| \\
&= \|a_n(x_n - T_2 y_n) + (1 - a_n)b_n(T_1 y_n - T_2 y_n)\| \\
&\leq a_n \|x_n - T_2 y_n\| + (1 - a_n)b_n \|T_1 y_n - T_2 y_n\| \\
&= a_n \|x_n - x_{n+1} + x_{n+1} - T_2 y_n\| + (1 - a_n)b_n \|T_2 y_n - T_1 y_n\| \\
&\leq a_n \|x_n - x_{n+1}\| + a_n \|x_{n+1} - T_2 y_n\| + (1 - a_n)b_n \|T_2 y_n - T_1 y_n\|,
\end{aligned} \tag{3.22}$$

which implies that

$$(1 - a_n) \|x_{n+1} - T_2 y_n\| \leq a_n \|x_n - x_{n+1}\| + (1 - a_n) b_n \|T_2 y_n - T_1 y_n\|. \quad (3.23)$$

Thus, we have

$$\begin{aligned} a_n(1 - a_n) \|x_{n+1} - T_2 y_n\| &\leq (1 - a_n) \|x_{n+1} - T_2 y_n\| \\ &\leq a_n \|x_n - x_{n+1}\| + (1 - a_n) b_n \|T_2 y_n - T_1 y_n\| \\ &\leq b \|x_n - x_{n+1}\| + (1 - a_n) \|T_2 y_n - T_1 y_n\| \\ &\leq b \|x_n - x_{n+1}\| + (1 - a) \|T_2 y_n - T_1 y_n\|. \end{aligned} \quad (3.24)$$

Hence, $\lim_{n \rightarrow \infty} a_n(1 - a_n) \|x_{n+1} - T_2 y_n\| = 0$. Since $\liminf_{n \rightarrow \infty} a_n(1 - a_n) > 0$, we have

$$\lim_{n \rightarrow \infty} \|x_{n+1} - T_2 y_n\| = 0. \quad (3.25)$$

Next, we prove that

$$\lim_{n \rightarrow \infty} \|y_n - T_1 y_n\| = 0, \quad \lim_{n \rightarrow \infty} \|y_n - T_2 y_n\| = 0. \quad (3.26)$$

Since $S_{1,r}$ and $S_{2,\lambda}$ are firmly nonexpansive, it follows that

$$\|u_n - y^*\|^2 = \|S_{2,\lambda} x_n - S_{2,\lambda} x^*\|^2 \leq \langle S_{2,\lambda} x_n - S_{2,\lambda} x^*, x_n - x^* \rangle = \langle u_n - y^*, x_n - x^* \rangle, \quad (3.27)$$

which gives that

$$\begin{aligned} \|(u_n - y^*) - (x_n - x^*)\|^2 &= \|u_n - y^*\|^2 - 2\langle u_n - y^*, x_n - x^* \rangle + \|x_n - x^*\|^2 \\ &\leq \|u_n - y^*\|^2 - 2\|u_n - y^*\|^2 + \|x_n - x^*\|^2 \\ &= -\|u_n - y^*\|^2 + \|x_n - x^*\|^2. \end{aligned} \quad (3.28)$$

This implies that

$$\|u_n - y^*\|^2 \leq \|x_n - x^*\|^2 - \|(u_n - y^*) - (x_n - x^*)\|^2. \quad (3.29)$$

By the convexity of $\|\cdot\|^2$, we have

$$\begin{aligned} \|x_{n+1} - x^*\|^2 &= \|a_n x_n + (1 - a_n)(b_n T_1 y_n + (1 - b_n) T_2 y_n) - x^*\|^2 \\ &= \|a_n(x_n - x^*) + (1 - a_n)(b_n T_1 y_n + (1 - b_n) T_2 y_n - x^*)\|^2 \end{aligned}$$

$$\begin{aligned}
&\leq a_n \|x_n - x^*\|^2 + (1 - a_n) \|b_n T_1 y_n + (1 - b_n) T_2 y_n - x^*\|^2 \\
&\leq a_n \|x_n - x^*\|^2 + (1 - a_n) \|y_n - x^*\|^2 \\
&\leq a_n \|x_n - x^*\|^2 + (1 - a_n) \|u_n - y^*\|^2 \\
&\leq a_n \|x_n - x^*\|^2 + (1 - a_n) \left(\|x_n - x^*\|^2 - \|(u_n - y^*) - (x_n - x^*)\|^2 \right) \\
&= \|x_n - x^*\|^2 - (1 - a_n) \|(u_n - y^*) - (x_n - x^*)\|^2.
\end{aligned} \tag{3.30}$$

Thus,

$$(1 - a_n) \|(u_n - y^*) - (x_n - x^*)\|^2 \leq \|x_n - x^*\|^2 - \|x_{n+1} - x^*\|^2. \tag{3.31}$$

Since $0 < a \leq a_n \leq b < 1$, we have

$$(1 - b) \|(u_n - y^*) - (x_n - x^*)\|^2 \leq (1 - a_n) \|(u_n - y^*) - (x_n - x^*)\|^2 \leq \|x_n - x^*\|^2 - \|x_{n+1} - x^*\|^2. \tag{3.32}$$

Since $\lim_{n \rightarrow \infty} \|x_n - x^*\|$ exists, we have

$$\lim_{n \rightarrow \infty} \|u_n - x_n + x^* - y^*\| = 0. \tag{3.33}$$

Similarly, we have

$$\|y_n - x^*\|^2 = \|S_{1,r_n} u_n - S_{1,r_n} y^*\|^2 \leq \langle S_{1,r_n} u_n - S_{1,r_n} y^*, u_n - y^* \rangle = \langle y_n - x^*, u_n - x^* \rangle, \tag{3.34}$$

which gives that

$$\begin{aligned}
\|(y_n - x^*) - (u_n - y^*)\|^2 &= \|y_n - x^*\|^2 - 2\langle y_n - x^*, u_n - x^* \rangle + \|u_n - y^*\|^2 \\
&\leq \|y_n - x^*\|^2 - 2\|y_n - x^*\|^2 + \|u_n - y^*\|^2 \\
&= -\|y_n - x^*\|^2 + \|u_n - y^*\|^2.
\end{aligned} \tag{3.35}$$

This implies that

$$\|y_n - x^*\|^2 \leq \|u_n - y^*\|^2 - \|y_n - u_n - x^* + y^*\|^2. \tag{3.36}$$

By the convexity of $\|\cdot\|^2$, we have

$$\begin{aligned}
\|x_{n+1} - x^*\|^2 &= \|a_n x_n + (1 - a_n)(b_n T_1 y_n + (1 - b_n) T_2 y_n) - x^*\|^2 \\
&\leq a_n \|x_n - x^*\|^2 + (1 - a_n) \|y_n - x^*\|^2
\end{aligned}$$

$$\begin{aligned}
&\leq a_n \|x_n - x^*\|^2 + (1 - a_n) \left(\|u_n - y^*\|^2 - \|y_n - u_n - x^* + y^*\|^2 \right) \\
&\leq a_n \|x_n - x^*\|^2 + (1 - a_n) \|x_n - x^*\|^2 - (1 - a_n) \|y_n - u_n - x^* + y^*\|^2 \\
&= \|x_n - x^*\|^2 - (1 - a_n) \|(y_n - x^*) - (u_n - y^*)\|^2.
\end{aligned} \tag{3.37}$$

Thus,

$$(1 - a_n) \|y_n - u_n - x^* + y^*\|^2 \leq \|x_n - x^*\|^2 - \|x_{n+1} - x^*\|^2. \tag{3.38}$$

Since $0 < a \leq a_n \leq b < 1$, we have

$$(1 - b) \|y_n - u_n - x^* + y^*\|^2 \leq (1 - a_n) \|y_n - u_n - x^* + y^*\|^2 \leq \|x_n - x^*\|^2 - \|x_{n+1} - x^*\|^2. \tag{3.39}$$

Since $\lim_{n \rightarrow \infty} \|x_n - x^*\|$ exists, we have

$$\lim_{n \rightarrow \infty} \|y_n - u_n - x^* + y^*\| = 0. \tag{3.40}$$

Hence,

$$\begin{aligned}
&\|(b_n T_1 y_n + (1 - b_n) T_2 y_n) - y_n\| \\
&= \|(b_n T_1 y_n + (1 - b_n) T_2 y_n) - x_n + x_n - u_n + u_n - y^* + y^* - x^* + x^* - y_n\| \\
&\leq \|(b_n T_1 y_n + (1 - b_n) T_2 y_n) - x_n\| + \|x_n - u_n - x^* + y^*\| + \|u_n - y_n + x^* - y^*\|.
\end{aligned} \tag{3.41}$$

It follow from (3.10), (3.33), and (3.40) that

$$\lim_{n \rightarrow \infty} \|(b_n T_1 y_n + (1 - b_n) T_2 y_n) - y_n\| = 0, \tag{3.42}$$

from which it follows that

$$\begin{aligned}
\|y_n - x_n\| &= \|y_n - (b_n T_1 y_n + (1 - b_n) T_2 y_n) + (b_n T_1 y_n + (1 - b_n) T_2 y_n) - x_n\| \\
&\leq \|y_n - (b_n T_1 y_n + (1 - b_n) T_2 y_n)\| \\
&\quad + \|(b_n T_1 y_n + (1 - b_n) T_2 y_n) - x_n\| \longrightarrow 0 \quad \text{as } n \longrightarrow \infty,
\end{aligned} \tag{3.43}$$

that is,

$$\lim_{n \rightarrow \infty} \|y_n - x_n\| = 0. \tag{3.44}$$

Thus,

$$\|y_n - T_1 y_n\| \leq \|y_n - x_n\| + \|x_n - x_{n+1}\| + \|x_{n+1} - T_1 y_n\| \longrightarrow 0. \tag{3.45}$$

Similarly, we have $\|y_n - T_2 y_n\| \rightarrow 0$. Since $\{y_n\}$ is bounded sequence, there exists a subsequence $\{y_{n_i}\}$ of $\{y_n\}$ such that $y_{n_i} \rightharpoonup \bar{x}$ as $i \rightarrow \infty$. Since T_1 and T_2 are demiclosed at 0, we conclude that $\bar{x} \in F(T_1) \cap F(T_2)$. Let G be a mapping which is defined as in Lemma 3.1. Thus, we have

$$\|y_n - G(y_n)\| = \|S_{1,r}S_{2,\lambda}x_n - G(y_n)\| = \|G(x_n) - G(y_n)\| \leq \|x_n - y_n\|, \quad (3.46)$$

and hence,

$$\begin{aligned} \|x_n - G(x_n)\| &= \|x_n - y_n + y_n - G(y_n) + G(y_n) - G(x_n)\| \\ &\leq \|x_n - y_n\| + \|y_n - G(y_n)\| + \|G(y_n) - G(x_n)\| \\ &\leq \|x_n - y_n\| + \|x_n - y_n\| + \|y_n - x_n\| \\ &= 3\|x_n - y_n\| \rightarrow 0. \end{aligned} \quad (3.47)$$

This together with $x_{n_i} \rightharpoonup \bar{x}$ implies that $\bar{x} \in F(G) := \Omega$, if $\{y_{n_j}\}$ is another subsequence of $\{y_n\}$ such that $y_{n_j} \rightharpoonup \hat{x}$ as $j \rightarrow \infty$. Since T_1 and T_2 are demiclosed at 0, we conclude that $\hat{x} \in F(T_1) \cap F(T_2) \cap \Omega$. From $x_{n_i} \rightharpoonup \bar{x}$ and $x_{n_j} \rightharpoonup \hat{x}$, we will show that $\bar{x} = \hat{x}$. Assume that $\bar{x} \neq \hat{x}$. Since $\lim_{n \rightarrow \infty} \|x_n - x^*\|$ exists for all $x^* \in F(T_1) \cap F(T_2) \cap \Omega$, by Opial's Theorem 2.2, we have

$$\begin{aligned} \lim_{n \rightarrow \infty} \|x_n - \bar{x}\| &= \liminf_{i \rightarrow \infty} \|x_{n_i} - \bar{x}\| \\ &< \liminf_{i \rightarrow \infty} \|x_{n_i} - \hat{x}\| \\ &= \lim_{n \rightarrow \infty} \|x_n - \hat{x}\| \\ &= \liminf_{j \rightarrow \infty} \|x_{n_j} - \hat{x}\| \\ &< \liminf_{j \rightarrow \infty} \|x_{n_j} - \bar{x}\| \\ &= \lim_{n \rightarrow \infty} \|x_n - \bar{x}\|. \end{aligned} \quad (3.48)$$

This is a contradiction. Thus, we have $\bar{x} = \hat{x}$. This implies that $y_n \rightharpoonup \bar{x} \in F(T_1) \cap F(T_2) \cap \Omega$. Since $\|x_n - y_n\| \rightarrow 0$, we have $x_n \rightharpoonup \bar{x}$. Put $z_n = P_{F(T_1) \cap F(T_2) \cap \Omega} x_n$. Finally, we show that $\bar{x} = \lim_{n \rightarrow \infty} z_n$. Now from (2.4) and $\bar{x} \in F(T_1) \cap F(T_2) \cap \Omega$, we have

$$\langle \bar{x} - z_n, z_n - x_n \rangle \geq 0. \quad (3.49)$$

Since $\{\|x_n - x^*\|\}$ is nonnegative and nonincreasing for all $x^* \in F(T_1) \cap F(T_2) \cap \Omega$, it follows by Lemma 2.1 that $\{z_n\}$ converges strongly to some $\hat{x} \in F(T_1) \cap F(T_2) \cap \Omega$. By (3.49), we have

$$\langle \bar{x} - \hat{x}, \hat{x} - \bar{x} \rangle \geq 0. \quad (3.50)$$

Therefore, $\bar{x} = \hat{x}$. □

Setting $T := T_1 = T_2$ in Theorem 3.2, we have the following result.

Corollary 3.3 (see [6]). *Let C be a closed convex subset of a real Hilbert space H . Let F_1 and F_2 be two bifunctions from $C \times C \rightarrow \mathbb{R}$ satisfying (A1)–(A4). Let $\lambda, \mu > 0$, and let $S_{1,\lambda}$ and $S_{2,\mu}$ be defined as in Lemma 2.5 associated to F_1 and F_2 , respectively. Let $T : C \rightarrow C$ be a quasi-nonexpansive mapping such that $I - T$ is demiclosed at zero and $F(T) \cap \Omega \neq \emptyset$. Suppose that $x_0 = x \in C$ and $\{x_n\}$, $\{y_n\}$, and $\{z_n\}$ are given by*

$$\begin{aligned} z_n \in C, \quad F_2(z_n, z) + \varphi(z) - \varphi(z_n) + \frac{1}{\mu} \langle z - z_n, z_n - x_n \rangle &\geq 0, \quad \forall z \in C, \\ y_n \in C, \quad F_1(y_n, z) + \varphi(z) - \varphi(y_n) + \frac{1}{\lambda} \langle z - y_n, y_n - z_n \rangle &\geq 0, \quad \forall z \in C, \\ x_{n+1} &= a_n x_n + (1 - a_n) T y_n, \end{aligned} \quad (3.51)$$

for all $n \in \mathbb{N}$, where $\{a_n\} \subset [a, b]$ for some $a, b \in (0, 1)$, and satisfy $\liminf_{n \rightarrow \infty} a_n(1 - a_n) > 0$, then $\{x_n\}$ converges weakly to $\bar{x} = \lim_{n \rightarrow \infty} P_{F(T) \cap \Omega} x_n$ and $(\bar{x}, \bar{y})$ is a solution of problem (1.5), where $\bar{y} = S_{2,\mu} \bar{x}$.

Setting $F_1 = F_2 \equiv 0$, $\varphi \equiv 0$ in Theorem 3.2, we have the following result.

Corollary 3.4 (see [9]). *Let H be a Hilbert space, let C be a nonempty, closed, and convex subset of H , and let T_1, T_2 be two quasi-nonexpansive mappings of C into itself such that $I - T_1, I - T_2$ are demiclosed at zero with $F(T_1) \cap F(T_2) \neq \emptyset$. For any x_1 in C , let $\{x_n\}$ be defined by*

$$x_{n+1} = (1 - a_n)x_n + a_n(b_n T_1 x_n + (1 - b_n)T_2 x_n), \quad (3.52)$$

where $\{a_n\}$ and $\{b_n\}$ are chosen so that

$$\liminf_{n \rightarrow \infty} a_n(1 - a_n) > 0 \quad \liminf_{n \rightarrow \infty} b_n(1 - b_n) > 0, \quad (3.53)$$

then $x_n \rightharpoonup p \in F(T_1) \cap F(T_2)$.

4. Applications

In this section, we apply our results to approximate a common element of the set of common fixed points of an asymptotic nonspreading mapping and an asymptotic TJ mapping and the solution set of SMEP in a real Hilbert space. We recall the following definitions. A mapping $T : C \rightarrow C$ is called nonspreading [13] if

$$2\|Tx - Ty\|^2 \leq \|Tx - y\|^2 + \|Ty - x\|^2 \quad \forall x, y \in C. \quad (4.1)$$

Furthermore, Takahashi and Yao [14] also introduced two nonlinear mappings in Hilbert spaces. A mapping $T : C \rightarrow C$ is called a $TJ - 1$ mapping [14] if

$$2\|Tx - Ty\|^2 = \|x - y\|^2 + \|Tx - y\|^2, \quad (4.2)$$

for all $x, y \in C$. A mapping $T : C \rightarrow C$ is called a $TJ - 2$ [14] mapping if

$$3\|Tx - Ty\|^2 = 2\|Tx - y\|^2 + \|Ty - x\|^2, \quad (4.3)$$

for all $x, y \in C$. For these two nonlinear mappings, $TJ - 1$ and $TJ - 2$ mappings, Takahashi and Yao [14] studied the existence results of fixed points in Hilbert spaces. Very recently, Lin et al. [15] introduced the following definitions of new mappings.

Definition 4.1. Let C be a nonempty closed convex subset of a Hilbert space H . We say that $T : C \rightarrow C$ is an asymptotic nonspreading mapping if there exist two functions $\alpha : C \rightarrow [0, 2)$ and $\beta : C \rightarrow [0, k]$, $k < 2$, such that

$$(A1) \quad 2\|Tx - Ty\|^2 \leq \alpha(x)\|Tx - y\|^2 + \beta(x)\|Ty - x\|^2 \text{ for all } x, y \in C,$$

$$(A2) \quad 0 < \alpha(x) + \beta(x) \leq 2 \text{ for all } x \in C.$$

Remark 4.2. The class of asymptotic nonspreading mappings contains the class of nonspreading mappings and the class of $TJ - 2$ mappings in a Hilbert space. Indeed, in Definition 4.1, we know that

(i) if $\alpha(x) = \beta(x) = 1$ for all $x \in C$, then T is a nonspreading mapping,

(ii) if $\alpha(x) = 4/3$ and $\beta(x) = 2/3$ for all $x \in C$, then T is a $TJ - 2$ mapping.

Definition 4.3. Let C be a nonempty closed convex subset of a Hilbert space H . We say $T : C \rightarrow C$ is an asymptotic TJ mapping if there exists two functions $\alpha : C \rightarrow [0, 2]$ and $\beta : C \rightarrow [0, k]$, $k < 2$, such that

$$(B1) \quad 2\|Tx - Ty\|^2 \leq \alpha(x)\|x - y\|^2 + \beta(x)\|Tx - y\|^2 \text{ for all } x, y \in C,$$

$$(B2) \quad \alpha(x) + \beta(x) \leq 2 \text{ for all } x \in C.$$

Remark 4.4. The class of asymptotic TJ mappings contains the class of $TJ - 1$ mappings and the class of nonexpansive mappings in a Hilbert space. Indeed, in Definition 4.3, we know that

(i) if $\alpha(x) = 2$ and $\beta(x) = 0$ for each $x \in C$, then T is a nonexpansive mapping,

(ii) if $\alpha(x) = \beta(x) = 1$ for each $x \in C$, then T is a $TJ - 1$ mapping.

It is well known that the set $F(T)$ of fixed points of a quasi-nonexpansive mapping T is a closed and convex set [1]. Hence, if $T : C \rightarrow C$ is an asymptotic nonspreading mapping (resp., asymptotic TJ mapping) with $F(T) \neq \emptyset$, then T is a quasi-nonexpansive mapping, and this implies that $F(T)$ is a nonempty closed convex subset of C .

Theorem 4.5 (see [15]). Let C be a nonempty closed convex subset of a real Hilbert space H , and let $T : C \rightarrow C$ be an asymptotic nonspreading mapping, then $I - T$ is demiclosed at 0.

Theorem 4.6 (see [15]). Let C be a nonempty closed convex subset of a real Hilbert space H , and let $T : C \rightarrow C$ be an asymptotic TJ mapping, then $I - T$ is demiclosed at 0.

Applying the above results, we have the following theorem.

Theorem 4.7. Let C be a nonempty closed convex subset of a real Hilbert space H . Let F_1 and F_2 be two bifunctions from $C \times C \rightarrow \mathbb{R}$ satisfying (A1)–(A4). Let $r, \lambda > 0$ and $S_{1,r}$ and $S_{2,\lambda}$ be defined as in Lemma 2.5 associated to F_1 and F_2 , respectively. Let $T_i : C \rightarrow C$, $i = 1, 2$, be any one of asymptotic nonspreading mapping and asymptotic TJ mapping such that $F(T_1) \cap F(T_2) \cap \Omega \neq \emptyset$. Let $\{x_n\}$, $\{y_n\}$, and $\{u_n\}$ be given by

$$\begin{aligned} x_1 &\in C \text{ chosen arbitrary,} \\ u_n &\in C, \quad F_2(u_n, z) + \varphi(z) - \varphi(u_n) + \frac{1}{\lambda} \langle z - u_n, u_n - x_n \rangle \geq 0, \quad \forall z \in C, \\ y_n &\in C, \quad 2F_1(y_n, z) + \varphi(z) - \varphi(y_n) + \frac{1}{r} \langle z - y_n, y_n - u_n \rangle \geq 0, \quad \forall z \in C, \\ x_{n+1} &= a_n x_n + (1 - a_n)(b_n T_1 y_n + (1 - b_n) T_2 y_n), \quad \forall n \in \mathbb{N}, \end{aligned} \quad (4.4)$$

where $\{a_n\}, \{b_n\} \subset [a, b]$ for some $a, b \in (0, 1)$ and satisfying

$$\liminf_{n \rightarrow \infty} a_n(1 - a_n) > 0, \quad \liminf_{n \rightarrow \infty} b_n(1 - b_n) > 0, \quad (4.5)$$

then $x_n \rightharpoonup \bar{x} := \lim_{n \rightarrow \infty} P_{F(T_1) \cap F(T_2) \cap \Omega} x_n$, and $(\bar{x}, \bar{y})$ is a solution of problem (1.5), where $\bar{y} = S_{2,\lambda} \bar{x}$.

Setting $F_1 = F_2 := F$ and $\varphi \equiv 0$ in the above theorem, we have the following result.

Corollary 4.8. Let C be a nonempty closed convex subset of a real Hilbert space H . Let F be the bifunctions from $C \times C \rightarrow \mathbb{R}$ satisfying (A1)–(A4). Let $T_i : C \rightarrow C$, $i = 1, 2$, be any one of asymptotic nonspreading mapping and asymptotic TJ mapping such that $\mathcal{F} := F(T_1) \cap F(T_2) \cap \text{EP}(F) \neq \emptyset$. For given $u \in C$ and $r > 0$, let the sequences $\{x_n\}$ and $\{u_n\}$ be defined by

$$\begin{aligned} x_1 &\in C \text{ chosen arbitrary,} \\ u_n &\in C, \quad F(u_n, y) + \frac{1}{r} \langle y - u_n, u_n - x_n \rangle \geq 0 \quad \forall y \in C, \\ x_{n+1} &= a_n x_n + (1 - a_n)(b_n T_1 u_n + (1 - b_n) T_2 u_n) \quad \forall n \in \mathbb{N}, \end{aligned} \quad (4.6)$$

where $\{a_n\}, \{b_n\}$ are two sequences in $(0, 1)$ satisfying

$$\liminf_{n \rightarrow \infty} a_n(1 - a_n) > 0, \quad \liminf_{n \rightarrow \infty} b_n(1 - b_n) > 0, \quad (4.7)$$

then $x_n \rightharpoonup w$ for some $w \in \mathcal{F}$.

Setting $F_1 = F_2 \equiv 0$ and $\varphi \equiv 0$ in Theorem 4.7, we have the following result.

Corollary 4.9 (see [15]). Let C be a nonempty closed convex subset of a real Hilbert space H , and let $T_i : C \rightarrow C$, $i = 1, 2$, be any one of asymptotic nonspreading mapping and asymptotic TJ mapping. Let $\mathcal{F} := F(T_1) \cap F(T_2) \neq \emptyset$. Let $\{a_n\}$ and $\{b_n\}$ be two sequences in $(0, 1)$. Let $\{x_n\}$ be defined by

$$\begin{aligned} x_1 &\in C \text{ chosen arbitrary,} \\ x_{n+1} &= a_n x_n + (1 - a_n)(b_n T_1 x_n + (1 - b_n) T_2 x_n). \end{aligned} \tag{4.8}$$

Assume that $\liminf_{n \rightarrow \infty} a_n(1 - a_n) > 0$ and $\liminf_{n \rightarrow \infty} b_n(1 - b_n) > 0$, then $x_n \rightharpoonup w$ for some $w \in \mathcal{F}$.

Acknowledgment

This paper is supported by the Centre of Excellence in Mathematics under the Commission on Higher Education, Ministry of Education, Thailand.

References

- [1] S. Itoh and W. Takahashi, "The common fixed point theory of singlevalued mappings and multi-valued mappings," *Pacific Journal of Mathematics*, vol. 79, no. 2, pp. 493–508, 1978.
- [2] K. Goebel and W. A. Kirk, *Topics in Metric Fixed Point Theory*, vol. 28 of *Cambridge Studies in Advanced Mathematics*, Cambridge University Press, Cambridge, Mass, USA, 1990.
- [3] E. Blum and W. Oettli, "From optimization and variational inequalities to equilibrium problems," *The Mathematics Student*, vol. 63, no. 1–4, pp. 123–145, 1994.
- [4] A. Moudafi, "Krasnoselski-Mann iteration for hierarchical fixed-point problems," *Inverse Problems*, vol. 23, no. 4, pp. 1635–1640, 2007.
- [5] A. Moudafi, "From alternating minimization algorithms and systems of variational inequalities to equilibrium problems," *Communications on Applied Nonlinear Analysis*, vol. 16, no. 3, pp. 31–35, 2009.
- [6] S. Plubtieng and K. Sombut, "Weak convergence theorems for a system of mixed equilibrium problems and nonspreading mappings in a Hilbert space," *Journal of Inequalities and Applications*, vol. 2010, Article ID 246237, 12 pages, 2010.
- [7] W. R. Mann, "Mean value methods in iteration," *Proceedings of the American Mathematical Society*, vol. 4, pp. 506–510, 1953.
- [8] S. Iemoto and W. Takahashi, "Approximating common fixed points of nonexpansive mappings and nonspreading mappings in a Hilbert space," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 71, no. 12, pp. e2082–e2089, 2009.
- [9] G. E. Kim, "Weak and strong convergence theorems of quasi-nonexpansive mappings in a Hilbert spaces," *Journal of Optimization Theory and Applications*, vol. 152, no. 3, pp. 727–738, 2010.
- [10] W. Takahashi, *Introduction to Nonlinear and Convex Analysis*, Yokohama Publishers, Yokohama, Japan, 2009.
- [11] W. Takahashi and M. Toyoda, "Weak convergence theorems for nonexpansive mappings and monotone mappings," *Journal of Optimization Theory and Applications*, vol. 118, no. 2, pp. 417–428, 2003.
- [12] S. D. Flâm and A. S. Antipin, "Equilibrium programming using proximal-like algorithms," *Mathematical Programming*, vol. 78, no. 1, pp. 29–41, 1997.
- [13] F. Kohsaka and W. Takahashi, "Fixed point theorems for a class of nonlinear mappings related to maximal monotone operators in banach spaces," *Archiv der Mathematik*, vol. 91, no. 2, pp. 166–177, 2008.
- [14] W. Takahashi and J. C. Yao, "Fixed point theorems and ergodic theorems for nonlinear mappings in hilbert spaces," *Taiwanese Journal of Mathematics*, vol. 15, no. 2, pp. 457–472, 2011.
- [15] L. J. Lin, C. H. Chuang, and Z. T. Yu, "Fixed point theorems for some new nonlinear mappings in Hilbert spaces," *Fixed Point Theory and Applications*, vol. 2011, no. 1, article 51, 2011.