Research Article

Approximation Theorems for Generalized Complex Kantorovich-Type Operators

N. I. Mahmudov and M. Kara

Eastern Mediterranean University, Gazimagusa, T.R. North Cyprus, Mersin 10, Turkey

Correspondence should be addressed to N. I. Mahmudov, nazim.mahmudov@emu.edu.tr

Received 28 April 2012; Accepted 3 September 2012

Academic Editor: Jinyun Yuan

Copyright © 2012 N. I. Mahmudov and M. Kara. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The order of simultaneous approximation and Voronovskaja-type results with quantitative estimate for complex q -Kantorovich polynomials ($q > 0$) attached to analytic functions on compact disks are obtained. In particular, it is proved that for functions analytic in $\{z \in \mathbb{C} : |z| < R\}$, $R > q$, the rate of approximation by the q -Kantorovich operators ($q > 1$) is of order q^{-n} versus $1/n$ for the classical Kantorovich operators.

1. Introduction

For each integer $k \geq 0$, the q -integer $[k]_q$ and the q -factorial $[k]_q!$ are defined by

$$[k]_q := \begin{cases} \frac{1-q^k}{1-q}, & \text{if } q \in \mathbb{R}^+ \setminus \{1\}, \\ k, & \text{if } q = 1, \end{cases} \quad \text{for } k \in \mathbb{N}, [0]_q = 0, \quad (1.1)$$

$$[k]_q! := [1]_q [2]_q \cdots [k]_q \quad \text{for } k \in \mathbb{N}, [0]! = 1.$$

For integers $0 \leq k \leq n$, the q -binomial coefficient is defined by

$$\begin{bmatrix} n \\ k \end{bmatrix}_q := \frac{[n]_q!}{[k]_q! [n-k]_q!}. \quad (1.2)$$

For fixed $1 \neq q > 0$, we denote the q -derivative $D_q f(z)$ of f by

$$D_q f(z) = \begin{cases} \frac{f(qz) - f(z)}{(q-1)z}, & z \neq 0, \\ f'(0), & z = 0. \end{cases} \quad (1.3)$$

Let $\mathbb{D}_R$ be a disc $\mathbb{D}_R := \{z \in \mathbb{C} : |z| < R\}$ in the complex plane $\mathbb{C}$. Denote by $H(\mathbb{D}_R)$ the space of all analytic functions on $\mathbb{D}_R$. For $f \in H(\mathbb{D}_R)$ we assume that $f(z) = \sum_{m=0}^{\infty} a_m z^m$.

In several recent papers, convergence properties of complex q -Bernstein polynomials, proposed by Phillips [1], defined by

$$B_{n,q}(f; z) = \sum_{k=0}^n f\left(\frac{[k]_q}{[n]_q}\right) \begin{bmatrix} n \\ k \end{bmatrix}_q x^k \prod_{j=0}^{n-k-1} (1 - q^j x) = \sum_{k=0}^n f\left(\frac{[k]_q}{[n]_q}\right) p_{n,k}(q; z) \quad (1.4)$$

and attached to an analytic function f in closed disks, were intensively studied by many authors; see [2] and references therein. It is known that the cases $0 < q < 1$ and $q > 1$ are not similar to each other. This difference is caused by the fact that, for $0 < q < 1$, $B_{n,q}$ are positive linear operators on $C[0, 1]$ while for $q > 1$, the positivity fails. The lack of positivity makes the investigation of convergence in the case $q > 1$ essentially more difficult than that for $0 < q < 1$. There are few papers [3–6] studying systematically the convergence of the q -Bernstein polynomials in the case $q > 1$. If $q \geq 1$ then qualitative Voronovskaja-type and saturation results for complex q -Bernstein polynomials were obtained in Wang and Wu [5]. Wu [6] studied saturation of convergence on the interval $[0, 1]$ for the q -Bernstein polynomials of a continuous function f for arbitrary fixed $q > 1$. On the other hand, Gal [7, 8], Anastassiou and Gal [9, 10], Mahmudov [11–13], and Mahmudov and Gupta [14] obtained quantitative estimates of the convergence and of the Voronovskaja's theorem in compact disks, for different complex Bernstein-Durrmeyer type operators.

The goal of the present note is to extend these type of results to complex Kantorovich operators based on the q -integers, in the case $q > 0$, defined as follows:

$$K_{n,q}(f; z) = \sum_{k=0}^n p_{n,k}(q; z) \int_0^1 f\left(\frac{q[k]_q + t}{[n+1]_q}\right) dt. \quad (1.5)$$

Notice that in the case $q = 1$, these operators coincide with the classical Kantorovich operators. For $0 < q \leq 1$ the operator $K_{n,q} : C[0, 1] \rightarrow C[0, 1]$ is positive and for $q > 1$, it is not positive. The problems studied in this paper in the case $q = 1$ were investigated in [2, 9].

We start with the following quantitative estimates of the convergence for complex q -Kantorovich-type operators attached to an analytic function in a disk of radius $R > 1$ and center 0.

Theorem 1.1. Let $f \in H(\mathbb{D}_R)$.

(i) Let $0 < q \leq 1$ and $1 \leq r < R$. For all $z \in \mathbb{D}_r$ and $n \in \mathbb{N}$, one has

$$|K_{n,q}(f; z) - f(z)| \leq \frac{3 + q^{-1}}{2[n]_q} \sum_{m=1}^{\infty} |a_m| m(m+1) r^m. \quad (1.6)$$

(ii) Let $1 < q < R < \infty$ and $1 \leq r < R/q$. For all $z \in \mathbb{D}_r$ and $n \in \mathbb{N}$, one has

$$|K_{n,q}(f; z) - f(z)| \leq \frac{2}{[n]_q} \sum_{m=1}^{\infty} |a_m| m(m+1) q^m r^m. \quad (1.7)$$

Remark 1.2. (i) Since $[n]_q \rightarrow (1-q)^{-1}$ as $n \rightarrow \infty$ in the estimate in Theorem 1.1(i) we do not obtain convergence of $K_{n,q}(f; z)$ to $f(z)$. But this situation can be improved by choosing $0 < q = q_n < 1$ with $q_n \nearrow 1$ as $n \rightarrow \infty$. Since in this case $[n]_{q_n} \rightarrow \infty$ as $n \rightarrow \infty$, from Theorem 1.1(i) we get uniform convergence in $\mathbb{D}_r$.

(ii) Theorem 1.1(ii) says that for functions analytic in $\mathbb{D}_R$, $R > q$, the rate of approximation by the q -Kantorovich operators ($q > 1$) is of order q^{-n} versus $1/n$ for the classical Kantorovich operators.

Let $f \in H(\mathbb{D}_R)$. Let us define

$$L_q(f; z) := \begin{cases} \frac{1-2z}{2} f'(z) + \frac{(1-z)(D_q f(z) - f'(z))}{1-q^{-1}}, & \text{if } |z| < \frac{R}{q}, R > q > 1, \\ \frac{1-2z}{2} f'(z) + \frac{z(1-z)}{2} f''(z), & \text{if } |z| < R, 0 < q \leq 1. \end{cases} \quad (1.8)$$

It is not difficult to show that

$$\begin{aligned} L_q(f; z) &= q(1-z) \sum_{m=1}^{\infty} a_m \frac{[m]_q - m}{q-1} z^{m-1} + \frac{1-2z}{2} \sum_{m=1}^{\infty} a_m m z^{m-1} \\ &= q \sum_{m=1}^{\infty} a_m ([1]_q + \cdots + [m-1]_q) z^{m-1} (1-z) + \frac{1-2z}{2} \sum_{m=1}^{\infty} a_m m z^{m-1}, \quad q > 1. \end{aligned} \quad (1.9)$$

Here we used the identity

$$\frac{[m]_q - m}{q-1} = [1]_q + \cdots + [m-1]_q. \quad (1.10)$$

The next theorem gives Voronovskaja-type result in compact disks, for complex q -Kantorovich operators attached to an analytic function in $\mathbb{D}_R$, $R > 1$ and center 0.

Theorem 1.3. Let $f \in H(\mathbb{D}_R)$.

(i) Let $0 < q \leq 1$ and $1 \leq r < R$. For all $z \in \mathbb{D}_r$ and $n \in \mathbb{N}$ one has

$$\left| K_{n,q}(f; z) - f(z) - \frac{1-2z}{2[n+1]_q} f'(z) - \frac{z(1-z)}{2[n+1]_q} f''(z) \right| \leq \frac{28+q^{-1}}{2[n]_q^2} \sum_{m=2}^{\infty} |a_m| m^2 (m-1)^2 r^m. \quad (1.11)$$

(ii) Let $1 < q < R < \infty$ and $1 \leq r < R/q^2$. For all $z \in \mathbb{D}_r$ and $n \in \mathbb{N}$, one has

$$\left| K_{n,q}(f; z) - f(z) - \frac{1}{[n+1]_q} L_q(f; z) \right| \leq \frac{14}{[n]_q^2} \sum_{m=2}^{\infty} |a_m| m^2 (m-1)^2 q^{2m} r^m. \quad (1.12)$$

Remark 1.4. (i) In the hypothesis on f in Theorem 1.3(i) choosing $0 < q_n < 1$ with $q_n \nearrow 1$ as $n \rightarrow \infty$, it follows that

$$\lim_{n \rightarrow \infty} [n+1]_{q_n} [K_{n,q_n}(f; z) - f(z)] = \frac{1-2z}{2} f'(z) + \frac{z(1-z)}{2} f''(z) \quad (1.13)$$

uniformly in any compact disk included in the open disk $\mathbb{D}_R$.

(ii) Theorem 1.3(ii) gives explicit formulas of Voronovskaja-type for the q -Kantorovich polynomials for $q > 1$.

(iii) Obviously the best order of approximation that can be obtained from the estimate Theorem 1.3(i) is $O(1/[n]_{q_n}^2)$ and $O(1/n^2)$ for $q = 1$, while the order given by Theorem 1.3(ii) is $O(1/q^{2n})$, $q > 1$, which is essentially better.

Next theorem shows that $L_q(f; z)$, $q \geq 1$, is continuous about the parameter q for $f \in H(\mathbb{D}_R)$, $R > 1$.

Theorem 1.5. Let $R > 1$ and $f \in H(\mathbb{D}_R)$. Then for any r , $0 < r < R$,

$$\lim_{q \rightarrow 1^+} L_q(f; z) = L_1(f; z) \quad (1.14)$$

uniformly on $\mathbb{D}_r$.

As an application of Theorem 1.3, we present the order of approximation for complex q -Kantorovich operators.

Theorem 1.6. Let $1 < q < R$, $1 \leq r < R/q^2$ (or $0 < q \leq 1$, $1 \leq r < R$) and $f \in H(\mathbb{D}_R)$. If f is not a constant function then the estimate

$$\|K_{n,q}(f) - f\|_r \geq \frac{1}{[n+1]_q} C_{r,q}(f), \quad n \in \mathbb{N} \quad (1.15)$$

holds, where the constant $C_{r,q}(f)$ depends on f , q and r but is independent of n .

2. Auxiliary Results

Lemma 2.1. Let $q > 0$. For all $n \in \mathbb{N}$, $m \in \mathbb{N} \cup \{0\}$, $z \in \mathbb{C}$ one has

$$K_{n,q}(e_m; z) = \sum_{j=0}^m \binom{m}{j} \frac{q^j [n]_q^j}{[n+1]_q^m (m-j+1)} B_{n,q}(e_j; z), \quad (2.1)$$

where $e_m(z) = z^m$.

Proof. The recurrence formula can be derived by direct computation.

$$\begin{aligned} K_{n,q}(e_m; z) &= \sum_{k=0}^n p_{n,k}(z) \sum_{j=0}^m \int_0^1 \binom{m}{j} \frac{q^j [k]_q^j t^{m-j}}{[n+1]_q^m} dt = \sum_{k=0}^n p_{n,k}(z) \sum_{j=0}^m \binom{m}{j} \frac{q^j [k]_q^j}{[n+1]_q^m (m-j+1)} \\ &= \sum_{j=0}^m \binom{m}{j} \frac{q^j [n]_q^j}{[n+1]_q^m (m-j+1)} \sum_{k=0}^n \frac{[k]_q^j}{[n]_q^j} p_{n,k}(z) \\ &= \sum_{j=0}^m \binom{m}{j} \frac{q^j [n]_q^j}{[n+1]_q^m (m-j+1)} B_{n,q}(e_j; z). \end{aligned} \quad (2.2)$$

□

Lemma 2.2. For all $z \in \mathbb{D}_r$, $r \geq 1$ one has

$$|K_{n,q}(e_m; z)| \leq r^m, \quad n, m \in \mathbb{N}. \quad (2.3)$$

Proof. Indeed, using the inequality $|B_{n,q}(e_j; z)| \leq r^j$ (see [3]), we get

$$\begin{aligned} |K_{n,q}(e_m; z)| &\leq \sum_{j=0}^m \binom{m}{j} \frac{q^j [n]_q^j}{[n+1]_q^m (m-j+1)} |B_{n,q}(e_j; z)| \\ &\leq \frac{1}{[n+1]_q^m} \sum_{j=0}^m \binom{m}{j} q^j [n]_q^j r^m = \left(\frac{1 + q[n]_q}{[n+1]_q} \right)^m r^m = r^m. \end{aligned} \quad (2.4)$$

□

Lemma 2.3. For all $n, m \in \mathbb{N}$, $z \in \mathbb{C}$, $1 \neq q > 0$ one has

$$\begin{aligned} K_{n,q}(e_{m+1}; z) &= \frac{z(1-z)}{[n]_q} D_q K_{n,q}(e_m; z) + z K_{n,q}(e_m; z) \\ &\quad + \frac{1}{[n+1]_q^{m+1}} \sum_{j=0}^{m+1} \binom{m+1}{j} q^j [n]_q^j \frac{1}{(m-j+2)} \left(1 - \frac{j[n+1]_q}{q(m+1)[n]_q} \right) B_{n,q}(e_j; z). \end{aligned} \quad (2.5)$$

Proof. We know that (see [2])

$$\frac{z(1-z)}{[n]_q} D_q B_{n,q}(e_j; z) = B_{n,q}(e_{j+1}; z) - z B_{n,q}(e_j; z). \quad (2.6)$$

Taking the derivative of the formula (2.1) and using the above formula we have

$$\begin{aligned} \frac{z(1-z)}{[n]_q} D_q K_{n,q}(e_m; z) &= \sum_{j=0}^m \binom{m}{j} \frac{q^j [n]_q^j}{[n+1]_q^m (m-j+1)} \frac{z(1-z)}{[n]_q} D_q B_{n,q}(e_j; z) \\ &= \sum_{j=0}^m \binom{m}{j} \frac{q^j [n]_q^j}{[n+1]_q^m (m-j+1)} (B_{n,q}(e_{j+1}; z) - z B_{n,q}(e_j; z)) \\ &= \sum_{j=1}^{m+1} \binom{m}{j-1} \frac{q^{j-1} [n]_q^{j-1}}{[n+1]_q^m (m-j+2)} B_{n,q}(e_j; z) - z K_{n,q}(e_m; z). \end{aligned} \quad (2.7)$$

It follows that

$$\begin{aligned} K_{n,q}(e_{m+1}; z) &= \frac{z(1-z)}{[n]_q} D_q K_{n,q}(e_m; z) + z K_{n,q}(e_m; z) \\ &\quad + \sum_{j=0}^{m+1} \binom{m+1}{j} \frac{q^j [n]_q^j}{[n+1]_q^{m+1} (m-j+2)} B_{n,q}(e_j; z) \\ &\quad - \sum_{j=1}^{m+1} \binom{m}{j-1} \frac{q^{j-1} [n]_q^{j-1}}{[n+1]_q^m (m-j+2)} B_{n,q}(e_j; z) \\ &= \frac{z(1-z)}{[n]_q} D_q K_{n,q}(e_m; z) + z K_{n,q}(e_m; z) + \frac{1}{[n+1]_q^{m+1} (m+2)} \\ &\quad + \sum_{j=1}^{m+1} \binom{m+1}{j} \frac{q^{j-1} [n]_q^{j-1}}{[n+1]_q^m (m-j+2)} \frac{(m+1)q[n]_q - j[n+1]_q}{(m+1)[n+1]_q} B_{n,q}(e_j; z) \\ &= \frac{z(1-z)}{[n]_q} D_q K_{n,q}(e_m; z) + z K_{n,q}(e_m; z) \\ &\quad + \sum_{j=0}^{m+1} \binom{m+1}{j} \frac{q^{j-1} [n]_q^{j-1}}{[n+1]_q^m (m-j+2)} \frac{(m+1)q[n]_q - j[n+1]_q}{(m+1)[n+1]_q} B_{n,q}(e_j; z). \end{aligned} \quad (2.8)$$

Here we used the identity

$$\binom{m}{j-1} = \binom{m+1}{j} \frac{j}{(m+1)}. \quad (2.9)$$

□

For $m \in \mathbb{N} \cup \{0\}$ define

$$E_{n,m}(z) := \begin{cases} K_{n,q}(e_m; z) - e_m(z) - \frac{1-2z}{2[n+1]_q} m z^{m-1} - \frac{z(1-z)}{2[n+1]_q} m(m-1) z^{m-2}, & \text{if } 0 < q \leq 1, \\ K_{n,q}(e_m; z) - e_m(z) - \frac{1-2z}{2[n+1]_q} m z^{m-1} - \sum_{j=1}^{m-1} [j]_q \frac{q z^{m-1}(1-z)}{[n+1]_q}, & \text{if } q > 1. \end{cases} \quad (2.10)$$

Here it is assumed that $\sum_{j=1}^0 [j]_q = 0$.

Lemma 2.4. Let $n, m \in \mathbb{N}$.

(a) If $0 < q < 1$, one has the following recurrence formula:

$$\begin{aligned} E_{n,m}(z) &= \frac{z(1-z)}{[n]_q} D_q(K_{n,q}(e_{m-1}; z) - e_{m-1}(z)) + z E_{n,m-1}(z) \\ &\quad + \left(\frac{m-1}{[n+1]_q} - \frac{[m-1]_q}{[n]_q} \right) z^{m-1}(1-z) - \frac{1-2z}{2[n+1]_q} z^{m-1} \\ &\quad + \frac{1}{[n+1]_q^m} \sum_{j=0}^m \binom{m}{j} \frac{q^j [n]_q^j}{(m-j+1)} \frac{mq[n]_q - j[n+1]_q}{mq[n]_q} B_{n,q}(e_j; z). \end{aligned} \quad (2.11)$$

(b) If $q > 1$, one has

$$\begin{aligned} E_{n,m}(z) &= \frac{z(1-z)}{[n]_q} D_q(K_{n,q}(e_{m-1}; z) - e_{m-1}(z)) + z E_{n,m-1}(z) + \frac{[m-1]_q}{[n]_q [n+1]_q} z^{m-1}(1-z) \\ &\quad - \frac{1-2z}{2[n+1]_q} z^{m-1} + \frac{1}{[n+1]_q^m} \sum_{j=0}^m \binom{m}{j} \frac{q^j [n]_q^j}{(m-j+1)} \frac{mq[n]_q - j[n+1]_q}{mq[n]_q} B_{n,q}(e_j; z). \end{aligned} \quad (2.12)$$

Proof. We give the proof for the case $q > 1$. The case $0 < q < 1$ is similar to that of $q > 1$.

(b) It is immediate that $E_{n,m}(z)$ is a polynomial of degree less than or equal to m and that $E_{n,0}(z) = E_{n,1}(z) = 0$.

Using the formula (2.5), we get

$$\begin{aligned}
 E_{n,m}(z) &= \frac{z(1-z)}{[n]_q} D_q(K_{n,q}(e_{m-1}; z) - e_{m-1}(z)) + \frac{[m-1]_q}{[n]_q} z^{m-1}(1-z) \\
 &\quad + z \left(E_{n,m-1}(z) + \frac{1-2z}{2[n+1]_q} (m-1)z^{m-2} + \sum_{j=1}^{m-2} [j]_q \frac{qz^{m-2}(1-z)}{[n+1]_q} \right) \\
 &\quad - \frac{1-2z}{2[n+1]_q} mz^{m-1} - \sum_{j=1}^{m-1} [j]_q \frac{qz^{m-1}(1-z)}{[n+1]_q} \\
 &\quad + \frac{1}{[n+1]_q^m} \sum_{j=0}^m \binom{m}{j} \frac{q^j [n]_q^j}{(m-j+1)} \frac{mq[n]_q - j[n+1]_q}{mq[n]_q} B_{n,q}(e_j; z).
 \end{aligned} \tag{2.13}$$

A simple calculation leads us to the following relationship:

$$\begin{aligned}
 E_{n,m}(z) &= \frac{z(1-z)}{[n]_q} D_q(K_{n,q}(e_{m-1}; z) - e_{m-1}(z)) + zE_{n,m-1}(z) + \frac{[m-1]_q}{[n]_q[n+1]_q} z^{m-1}(1-z) \\
 &\quad - \frac{1-2z}{2[n+1]_q} z^{m-1} + \frac{1}{[n+1]_q^m} \sum_{j=0}^m \binom{m}{j} \frac{q^j [n]_q^j}{(m-j+1)} \frac{mq[n]_q - j[n+1]_q}{mq[n]_q} B_{n,q}(e_j; z),
 \end{aligned} \tag{2.14}$$

which is the desired recurrence formula. $\square$

Remark 2.5. Lemmas 2.3 and 2.4 are true in the case $q = 1$. In the formulae, we have to replace q -derivative by the ordinary derivative.

3. Proofs of the Main Results

We give proofs for the case $q > 1$. The case $0 < q < 1$ and $q = 1$ are similar to that of $q > 1$.

Proof of Theorem 1.1. The use of the above recurrence we obtain the following relationship:

$$\begin{aligned}
 K_{n,q}(e_m; z) - e_m(z) &= \frac{z(1-z)}{[n]_q} D_q K_{n,q}(e_{m-1}; z) + z(K_{n,q}(e_{m-1}; z) - e_{m-1}(z)) \\
 &\quad + \frac{1}{[n+1]_q^m} \sum_{j=0}^m \binom{m}{j} \frac{q^j [n]_q^j}{(m-j+1)} \frac{mq[n]_q - j[n+1]_q}{mq[n]_q} B_{n,q}(e_j; z).
 \end{aligned} \tag{3.1}$$

We can easily estimate the sum in the above formula as follows:

$$\begin{aligned}
& \left| \frac{1}{[n+1]_q^m} \sum_{j=0}^m \binom{m}{j} q^j [n]_q^j \frac{1}{(m-j+1)} \left(1 - \frac{j}{m} - \frac{j}{mq[n]_q} \right) B_{n,q}(e_j; z) \right| \\
& \leq \frac{1}{[n+1]_q^m} \left(\sum_{j=0}^{m-1} \binom{m-1}{j} \frac{m}{m-j} \frac{q^j [n]_q^j}{m-j+1} \left| 1 - \frac{j}{m} - \frac{j}{mq[n]_q} \right| \right) |B_{n,q}(e_j; z)| + \frac{q^{m-1} [n]_q^{m-1}}{[n+1]_q^m} r^m \\
& \leq \frac{2m \left(q[n]_q + 1 \right)^{m-1} + q^{m-1} [n]_q^{m-1}}{[n+1]_q^m} r^m \leq \frac{2m+1}{[n+1]_q} r^m.
\end{aligned} \tag{3.2}$$

It is known that by a linear transformation, the Bernstein inequality in the closed unit disk becomes

$$|P'_m(z)| \leq \frac{m}{qr} \|P_m\|_{qr}, \quad \forall |z| \leq qr, \quad r \geq 1, \tag{3.3}$$

(where $\|P_m\|_{qr} = \max\{|P_m(z)| : |z| \leq qr\}$) which combined with the mean value theorem in complex analysis implies

$$|D_q(P_m; z)| = \left| \frac{P_m(qz) - P_m(z)}{qz - z} \right| \leq \|P'_m\|_{qr} \leq \frac{m}{qr} \|P_m\|_{qr}, \tag{3.4}$$

for all $|z| \leq r$, where $P_m(z)$ is a complex polynomial of degree $\leq m$. From the above recurrence formula (3.1), we get

$$\begin{aligned}
|K_{n,q}(e_m; z) - e_m(z)| & \leq \frac{|z||1-z|}{[n]_q} |D_q K_{n,q}(e_{m-1}; z)| + |z| |K_{n,q}(e_{m-1}; z) - e_{m-1}(z)| + \frac{2m+1}{[n+1]_q} r^m \\
& \leq \frac{r(1+r)}{[n]_q} \frac{m-1}{qr} \|K_{n,q}(e_{m-1})\|_{qr} + r |K_{n,q}(e_{m-1}; z) - e_{m-1}(z)| + \frac{2m+1}{[n+1]_q} r^m \\
& \leq r |K_{n,q}(e_{m-1}; z) - e_{m-1}(z)| + \frac{2(m-1)}{[n]_q} q^{m-1} r^m + \frac{2m+1}{[n+1]_q} r^m \\
& \leq r |K_{n,q}(e_{m-1}; z) - e_{m-1}(z)| + \frac{4m}{[n]_q} q^m r^m.
\end{aligned} \tag{3.5}$$

By writing the last inequality for $m = 1, 2, \dots$, we easily obtain, step by step, the following:

$$\begin{aligned} |K_n(e_m; z) - e_m(z)| &\leq \frac{4m}{[n]_q} q^m r^m + r \frac{4(m-1)}{[n]_q} q^{m-1} r^{m-1} + r^2 \frac{4(m-2)}{[n]_q} q^{m-2} r^{m-2} + \dots + r^{m-1} \frac{4}{[n]_q} q r \\ &= \frac{4}{[n]_q} q^m r^m (m + m - 1 + \dots + 1) \leq \frac{2m(m+1)}{[n]_q} q^m r^m. \end{aligned} \quad (3.6)$$

Since $K_{n,q}(f; z)$ is analytic in $\mathbb{D}_R$, we can write

$$K_{n,q}(f; z) = \sum_{m=0}^{\infty} a_m K_{n,q}(e_m; z), \quad z \in \mathbb{D}_R, \quad (3.7)$$

which together with (3.6) immediately implies for all $|z| \leq r$

$$|K_{n,q}(f; z) - f(z)| \leq \sum_{m=0}^{\infty} |a_m| |K_{n,q}(e_m; z) - e_m(z)| \leq \frac{2}{[n]_q} \sum_{m=1}^{\infty} |c_m| m(m+1) (qr)^m. \quad (3.8) \quad \square$$

Proof of Theorem 1.3. A simple calculation and the use of the recurrence formula (2.5) lead us to the following relationship:

$$\begin{aligned} E_{n,m}(z) &= \frac{z(1-z)}{[n]_q} D_q(K_{n,q}(e_{m-1}; z) - e_{m-1}(z)) + zE_{n,m-1}(z) + \frac{[m-1]_q}{[n]_q[n+1]_q} z^{m-1}(1-z) \\ &\quad + \frac{1}{[n+1]_q} (z^m - B_{n,q}(e_m; z)) + \frac{1}{[n+1]_q} \left(1 - \frac{q^{m-1}[n]_q^{m-1}}{[n+1]_q^{m-1}} \right) B_{n,q}(e_m; z) \\ &\quad + \frac{1}{2[n+1]_q} \left(\frac{q^{m-1}[n]_q^{m-1}}{[n+1]_q^{m-1}} - 1 \right) B_{n,q}(e_{m-1}; z) + \frac{1}{2[n+1]_q} (B_{n,q}(e_{m-1}; z) - z^{m-1}) \\ &\quad - \frac{(m-1)q^{m-2}[n]_q^{m-2}}{2[n+1]_q^m} B_{n,q}(e_{m-1}; z) \\ &\quad + \frac{1}{[n+1]_q^m} \sum_{j=0}^{m-2} \binom{m}{j} \frac{q^j [n]_q^j}{(m-j+1)} \frac{mq[n]_q - j[n+1]_q}{mq[n]_q} B_{n,q}(e_j; z) \\ &:= \sum_{k=1}^9 I_k. \end{aligned} \quad (3.9)$$

Firstly, we estimate I_3, I_8 . It is clear that

$$\begin{aligned} |I_3| &\leq \frac{[m-1]_q}{[n]_q[n+1]_q} r^{m-1}(1+r), \\ |I_8| &\leq \frac{(m-1)}{2[n+1]_q^2} |B_{n,q}(e_{m-1}; z)| \leq \frac{(m-1)}{2[n+1]_q^2} r^{m-1}. \end{aligned} \quad (3.10)$$

Secondly, using the known inequality

$$1 - \prod_{k=1}^m x_k \leq \sum_{k=1}^m (1 - x_k), \quad 0 \leq x_k \leq 1, \quad (3.11)$$

to estimate I_5, I_6, I_9 .

$$\begin{aligned} |I_5| &\leq \frac{1}{[n+1]_q} \left(1 - \frac{q^{m-1}[n]_q^{m-1}}{[n+1]_q^{m-1}} \right) |B_{n,q}(e_m; z)| \leq \frac{m-1}{[n+1]_q^2} r^m, \\ |I_6| &\leq \frac{1}{2[n+1]_q} \left(1 - \frac{q^{m-1}[n]_q^{m-1}}{[n+1]_q^{m-1}} \right) |B_{n,q}(e_{m-1}; z)| \leq \frac{m-1}{2[n+1]_q^2} r^{m-1}, \\ |I_9| &\leq \frac{1}{[n+1]_q^m} \sum_{j=0}^{m-2} \binom{m-2}{j} \frac{m(m-1)}{(m-j)(m-j-1)} \frac{q^j [n]_q^j}{(m-j+1)} \left(1 - \frac{j}{m} - \frac{j}{mq[n]_q} \right) r^j \\ &\leq \frac{2m(m-1)[n+1]_q^{m-2}}{[n+1]_q^m} r^m = \frac{2m(m-1)}{[n+1]_q^2} r^m. \end{aligned} \quad (3.12)$$

Finally, we estimate I_4, I_7 . We use [2, Theorem 1.1.2]

$$\begin{aligned} |I_4| + |I_7| &\leq \frac{1}{[n+1]_q} |z^m - B_{n,q}(e_m; z)| + \frac{1}{2[n+1]_q} |B_n(e_{m-1}; z) - z^{m-1}| \\ &\leq \frac{2[m-1]_q(m-1)}{[n]_q[n+1]_q} r^m + \frac{[m-2]_q(m-2)}{[n]_q[n+1]_q} r^{m-1}. \end{aligned} \quad (3.13)$$

Using (3.6), (3.10), (3.12), and (3.13) in (3.9) finally we have ($m \geq 3$)

$$\begin{aligned}
|E_{n,m}(z)| &\leq \frac{r(1+r)}{[n]_q} |D_q(K_{n,q}(e_{m-1}; z) - e_{m-1}(z))| + r|E_{n,m-1}(z)| + \frac{[m-1]_q}{[n]_q[n+1]_q} r^{m-1}(1+r) \\
&\quad + \frac{2[m-1]_q(m-1)}{[n]_q[n+1]_q} r^m + \frac{m-1}{[n+1]_q^2} r^m + \frac{m-1}{2[n+1]_q^2} r^{m-1} + \frac{[m-2]_q(m-2)}{[n]_q[n+1]_q} r^{m-1} \\
&\quad + \frac{(m-1)}{2[n+1]_q^2} r^{m-1} + \frac{2m(m-1)}{[n+1]_q^2} r^m \\
&\leq \frac{r(1+r)}{[n]_q} \frac{m-1}{qr} \|K_{n,q}(e_{m-1}) - e_{m-1}\|_{qr} + r|E_{n,m-1}(z)| + \frac{10m[m-1]_q}{[n]_q^2} r^m \\
&\leq \frac{(m-1)(1+r)}{[n]_q} \frac{2(m-1)m}{[n]_q} q^{2(m-1)} r^{m-1} + r|E_{n,m-1}(z)| + \frac{10m[m-1]_q}{[n]_q^2} r^m \\
&\leq r|E_{n,m-1}(z)| + \frac{4m(m-1)^2}{[n]_q^2} q^{2m} r^m + \frac{10m(m-1)}{[n]_q^2} q^m r^m \\
&\leq r|E_{n,m-1}(z)| + \frac{14m^2(m-1)^2}{[n]_q^2} q^{2m} r^m.
\end{aligned} \tag{3.14}$$

As a consequence, we get

$$|E_{n,m}(z)| \leq \frac{14m^2(m-1)^2}{[n]_q^2} q^{2m} r^m. \tag{3.15}$$

This inequality combined with

$$\left| K_{n,q}(f; z) - f(z) - \frac{1}{[n+1]_q} L_q(f, z) \right| \leq \sum_{m=1}^{\infty} |a_m| |E_{n,m}(z)| \tag{3.16}$$

immediately implies the required estimate in statement.

Note that since $f^{(4)} = \sum_{m=4}^{\infty} a_m m(m-1)(m-2)(m-3)z^{m-4}$ and the series is absolutely convergent for all $|z| < R$, it easily follows the finiteness of the involved constants in the statement. $\square$

Proof of Theorem 1.6. For all $z \in \mathbb{D}_R$ and $n \in \mathbb{N}$, we get

$$K_{n,q}(f; z) - f(z) = \frac{1}{[n+1]_q} \left\{ L_q(f; z) + [n+1]_q \left(K_{n,q}(f; z) - f(z) - \frac{1}{[n+1]_q} L_q(f; z) \right) \right\}. \tag{3.17}$$

We apply

$$\|F + G\|_r \geq \| \|F\|_r - \|G\|_r \| \geq \|F\|_r - \|G\|_r \quad (3.18)$$

to get

$$\|K_{n,q}(f) - f\|_r \geq \frac{1}{[n+1]_q} \left\{ \|L_q(f; z)\|_r - [n+1]_q \left\| K_{n,q}(f; z) - f(z) - \frac{1}{[n+1]_q} L_q(f; z) \right\|_r \right\}. \quad (3.19)$$

Because by hypothesis f is not a constant in $\mathbb{D}_R$, it follows $\|L_q(f; z)\|_r > 0$. Indeed, assuming the contrary, it follows that $L_q(f; z) = 0$ for all $z \in \overline{\mathbb{D}}_R$ that is

$$\begin{aligned} \sum_{m=1}^{\infty} a_m \left(\frac{1}{2} - z \right) m z^{m-1} + \sum_{m=1}^{\infty} a_m \sum_{j=1}^{m-1} [j]_q z^{m-1} (1-z) &= 0, \\ \frac{1}{2} a_1 + a_1 + \sum_{m=1}^{\infty} \left(\frac{1}{2} (m+1) a_{m+1} - a_m + a_{m+1} \sum_{j=1}^m [j]_q - a_m \sum_{j=1}^{m-1} [j]_q \right) z^m &= 0 \end{aligned} \quad (3.20)$$

for all $z \in \overline{\mathbb{D}}_R \setminus \{0\}$. Thus $a_m = 0$, $m = 1, 2, 3, \dots$. Thus, f is constant, which is contradiction with the hypothesis.

Now, by Theorem 1.3, we have

$$\begin{aligned} [n+1]_q \left\| K_{n,q}(f; z) - f(z) - \frac{1}{[n+1]_q} L_q(f; z) \right\| \\ \leq \frac{[n+1]_q}{[n]_q} \frac{14}{[n]_q} \sum_{m=2}^{\infty} |a_m| m^2 (m-1)^2 q^{2m} r^m \rightarrow 0 \quad \text{as } n \rightarrow \infty. \end{aligned} \quad (3.21)$$

Consequently, there exists n_1 (depending only on f and r) such that for all $n \geq n_1$ we have

$$\|L_q(f; z)\|_r - [n+1]_q \left\| K_{n,q}(f; z) - f(z) - \frac{1}{[n+1]_q} L_q(f; z) \right\|_r \geq \frac{1}{2} \|L_q(f; z)\|_r, \quad (3.22)$$

which implies

$$\|K_{n,q}(f) - f\|_r \geq \frac{1}{[n+1]_q} \frac{1}{2} \|L_q(f; z)\|_r, \quad \forall n \geq n_1. \quad (3.23)$$

For $1 \leq n \leq n_1 - 1$, we have

$$\|K_{n,q}(f) - f\|_r \geq \frac{1}{[n+1]_q} \left([n+1]_q \|K_{n,q}(f) - f\|_r \right) = \frac{1}{[n+1]_q} M_{r,n}(f) > 0, \quad (3.24)$$

which finally implies that

$$\|K_{n,q}(f) - f\|_r \geq \frac{1}{[n+1]_q} C_{r,q}(f), \quad (3.25)$$

for all n , with $C_{r,q}(f) = \min\{M_{r,1}(f), \dots, M_{r,n_1-1}(f), (1/2)\|L_q(f; z)\|_r\}$. $\square$

Proof of Theorem 1.5. Proof is similar to that of Theorem 1.3 [5]. $\square$

References

- [1] G. M. Phillips, "Bernstein polynomials based on the q -integers," *Annals of Numerical Mathematics*, vol. 4, no. 1–4, pp. 511–518, 1997.
- [2] S. G. Gal, *Approximation by complex Bernstein and convolution type operators*, vol. 8, World Scientific, 2009.
- [3] S. Ostrovska, " q -Bernstein polynomials and their iterates," *Journal of Approximation Theory*, vol. 123, no. 2, pp. 232–255, 2003.
- [4] S. Ostrovska, "The sharpness of convergence results for q -Bernstein polynomials in the case $q > 1$," *Czechoslovak Mathematical Journal*, vol. 58, no. 4, pp. 1195–1206, 2008.
- [5] H. Wang and X. Wu, "Saturation of convergence for q -Bernstein polynomials in the case $q > 1$," *Journal of Mathematical Analysis and Applications*, vol. 337, no. 1, pp. 744–750, 2008.
- [6] Z. Wu, "The saturation of convergence on the interval $[0, 1]$ for the q -Bernstein polynomials in the case $q > 1$," *Journal of Mathematical Analysis and Applications*, vol. 357, no. 1, pp. 137–141, 2009.
- [7] S. G. Gal, "Voronovskaja's theorem and iterations for complex Bernstein polynomials in compact disks," *Mediterranean Journal of Mathematics*, vol. 5, no. 3, pp. 253–272, 2008.
- [8] S. G. Gal, "Approximation by complex genuine Durrmeyer type polynomials in compact disks," *Applied Mathematics and Computation*, vol. 217, no. 5, pp. 1913–1920, 2010.
- [9] G. A. Anastassiou and S. G. Gal, "Approximation by complex Bernstein-Schurer and Kantorovich-Schurer polynomials in compact disks," *Computers & Mathematics with Applications*, vol. 58, no. 4, pp. 734–743, 2009.
- [10] G. A. Anastassiou and S. G. Gal, "Approximation by complex bernstein-durrmeyer polynomials in compact disks," *Mediterranean Journal of Mathematics*, vol. 7, no. 4, pp. 471–482, 2010.
- [11] N. I. Mahmudov, "Approximation by genuine q -Bernstein-Durrmeyer polynomials in compact disks," *Hacetatepe Journal of Mathematics and Statistics*, vol. 40, no. 1, pp. 77–89, 2011.
- [12] N. I. Mahmudov, "Convergence properties and iterations for q -Stancu polynomials in compact disks," *Computers & Mathematics with Applications. An International Journal*, vol. 59, no. 12, pp. 3763–3769, 2010.
- [13] N. I. Mahmudov, "Approximation by Bernstein-Durrmeyer-type operators in compact disks," *Applied Mathematics Letters*, vol. 24, no. 7, pp. 1231–1238, 2011.
- [14] N. I. Mahmudov and V. Gupta, "Approximation by genuine Durrmeyer-Stancu polynomials in compact disks," *Mathematical and Computer Modelling*, vol. 55, no. 3–4, pp. 278–285, 2012.