Research Article

Asymptotic Stability of a Class of Impulsive Delay Differential Equations

G. L. Zhang, M. H. Song, and M. Z. Liu

Department of Mathematics, Harbin Institute of Technology, Harbin 150001, China

Correspondence should be addressed to M. H. Song, songmh@lsec.cc.ac.cn

Received 8 June 2012; Revised 22 August 2012; Accepted 9 September 2012

Academic Editor: Jaime Munoz Rivera

Copyright © 2012 G. L. Zhang et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

This paper is concerned with a class of linear impulsive delay differential equations. Asymptotic stability of analytic solutions of this kind of equations is studied by the property of delay differential equations without impulsive perturbations. New numerical methods for this kind of equations are constructed. The convergence and asymptotic stability of the methods for this kind of equations are studied.

1. Introduction

Impulsive differential equations arise widely in the study of medicine, biology, economics, and engineering, and so forth. In recent years, theory of impulsive differential delay equations (IDDEs) has been an object of active research (see [1–18] and the reference therein). The results about the existence and uniqueness of IDDEs have been studied in [2, 3, 10]. And the stability of IDDEs have attracted increasing interest in both theoretical research and practical applications (see [1, 3, 4, 6–18] and the reference therein). In this paper, we use the property of delay differential equations without impulsive perturbations to study asymptotic stability of analytic solutions of a class of linear impulsive delay differential equations.

There are a few papers on numerical methods of impulsive differential equations. In [5], Covachev et al. obtained a convergent difference approximation for a nonlinear impulsive ordinary system in a Banach space. In [9, 19], the authors studied the stability of Runge-Kutta methods for linear impulsive ordinary differential equations. In [20], Ding et al. studied the convergence property of an Euler method for IDDEs. In this paper, we construct new numerical methods for a class of linear impulsive delay differential equations. The convergence and asymptotic stability of the methods for this kind of equations are studied.

2. Stability of Analytic Solutions of a Class of Linear IDDEs

In this paper, we consider the following equation:

$$\begin{aligned} x'(t) &= px(t) + qx(t - \tau), \quad t \geq 0, \quad t \neq k\tau, \quad k = 0, 1, 2, \dots, \\ x(t) &= rx(t^-), \quad t = k\tau, \quad k = 0, 1, 2, \dots, \\ x(t) &= \Phi(t), \quad t \in [-\tau, 0), \end{aligned} \quad (2.1)$$

where $r \neq 0$, $\tau > 0$, p and q are real constants, Φ is a continuous function on $[-\tau, 0)$, and $x'(t)$ denotes the right-hand derivative of $x(t)$.

Definition 2.1. The zero solution of (2.1) is said to be asymptotically stable, if

$$\lim_{t \rightarrow \infty} x(t) = 0, \quad (2.2)$$

where $x(t)$ is the solution of (2.1) for any initial function $\Phi \in C([-\tau, 0), R)$.

We define $\Phi(0) = \lim_{t \rightarrow 0^-} \Phi(t)$ and the delay differential equations without impulsive perturbations as follows:

$$\begin{aligned} y'(t) &= \left(\frac{\ln r}{\tau} + p \right) y(t) + qy(t - \tau), \quad t \geq 0, \\ y(t) &= \Psi(t) = r^{t/\tau+1} \Phi(t), \quad t \in [-\tau, 0]. \end{aligned} \quad (2.3)$$

In the rest of the paper, $\{t\}$ is defined as $\{t\} = t - [t]$ where $[\cdot]$ denotes the greatest integer function.

Theorem 2.2. Assume that $x(t)$ is the solution of (2.1) and $y(t) = r^{\{t/\tau\}} x(t)$, $t \in [-\tau, \infty)$, then $y(t)$ is the solution of (2.3). On the other hand, assume that $y(t)$ is the solution of (2.3) and $x(t) = r^{-\{t/\tau\}} y(t)$, $t \in [-\tau, \infty)$, then $x(t)$ is the solution of (2.1).

Proof. (i) We prove that $y(t)$ is continuous if $x(t)$ is the solution of (2.1). Obviously,

$$y(t) = r^{\{t/\tau\}} x(t) = r^{t/\tau-k} x(t) \quad (2.4)$$

is continuous on $[k\tau, (k+1)\tau)$, $k = -1, 0, 1, \dots$. We know that $y(t) = \Psi(t) = r^{t/\tau+1} \Phi(t)$, $t \in [-\tau, 0]$ is continuous. Therefore,

$$\begin{aligned} y(0^-) &= \lim_{t \rightarrow 0^-} r^{t/\tau+1} x(t) = rx(0^-) = r\Phi(0^-), \\ y(0) &= x(0) = rx(0^-) = r\Phi(0^-). \end{aligned} \quad (2.5)$$

Hence, $y(0) = y(0^-)$. So $y(t)$ is continuous at $t = 0$. It follows from

$$\begin{aligned} y((k+1)\tau) &= r^{\lceil((k+1)\tau)/\tau\rceil} x((k+1)\tau) = r^{\lceil k+1 \rceil} x((k+1)\tau) = x((k+1)\tau) = rx((k+1)\tau^-), \\ y((k+1)\tau^-) &= \lim_{t \rightarrow (k+1)\tau^-} r^{\lceil t/\tau \rceil} x(t) = rx((k+1)\tau^-), \end{aligned} \quad (2.6)$$

that $y(t)$ is continuous at $t = (k+1)\tau$, $k = 0, 1, 2, \dots$. Hence $y(t)$ is continuous on $t \in [-\tau, \infty)$.

(ii) We prove that $y(t)$ is the solution of (2.3) if $x(t)$ is the solution of (2.1). For $t \in [k\tau, (k+1)\tau)$, $k = 0, 1, \dots$, we have

$$\begin{aligned} y'(t) &= r^{t/\tau-k} x(t) \frac{\ln r}{\tau} + r^{t/\tau-k} x'(t) \\ &= r^{t/\tau-k} x(t) \frac{\ln r}{\tau} + r^{t/\tau-k} (px(t) + qx(t-\tau)) \\ &= \left(p + \frac{\ln r}{\tau} \right) \left(r^{t/\tau-k} x(t) \right) + q \left(r^{((t-\tau)/\tau)-(k-1)} x(t-\tau) \right) \\ &= \left(p + \frac{\ln r}{\tau} \right) y(t) + qy(t-\tau). \end{aligned} \quad (2.7)$$

Because $y'(t)$ denotes the right-hand derivative of $y(t)$, $y'(t) = (p + \ln r/\tau)y(t) + qy(t-\tau)$, hence $y(t)$ is the solution of (2.3).

(iii) We prove that $x(t)$ is the solution of (2.1) if $y(t)$ is the solution of (2.3). For $t \in [k\tau, (k+1)\tau)$, $k = 0, 1, \dots$,

$$\begin{aligned} x'(t) &= -r^{-(t/\tau-k)} y(t) \frac{\ln r}{\tau} + r^{-(t/\tau-k)} y'(t) \\ &= -r^{-(t/\tau-k)} y(t) \frac{\ln r}{\tau} + r^{-(t/\tau-k)} \left(\left(p + \frac{\ln r}{\tau} \right) y(t) + qy(t-\tau) \right) \\ &= p \left(r^{-(t/\tau-k)} y(t) \right) + q \left(r^{-(t/\tau-k)} y(t-\tau) \right) \\ &= p \left(r^{-(t/\tau-k)} y(t) \right) + q \left(r^{-((t-\tau)/\tau-(k-1))} y(t-\tau) \right) = px(t) + qx(t-\tau). \end{aligned} \quad (2.8)$$

Obviously,

$$\begin{aligned} x((k+1)\tau) &= r^{-\lceil k+1 \rceil} y((k+1)\tau) = y((k+1)\tau), \\ x((k+1)\tau^-) &= \lim_{t \rightarrow (k+1)\tau^-} r^{-\lceil t/\tau \rceil} y(t) = r^{-1} y((k+1)\tau). \end{aligned} \quad (2.9)$$

So $x((k+1)\tau) = rx((k+1)\tau^-)$, $k = 0, 1, \dots$. Consequently, $x(t)$ is the solution of (2.1). $\square$

By Theorem 2.2, we obtain the following corollary.

Corollary 2.3. Assume that $\lim_{t \rightarrow \infty} x(t) = 0$, where $x(t)$ is the solution of (2.1). Then $\lim_{t \rightarrow \infty} y(t) = 0$, where $r^{(t/\tau)} x(t)$, $t \in [-\tau, \infty)$, and $y(t)$ also is the solution of (2.3). On the other hand, assume that $\lim_{t \rightarrow \infty} y(t) = 0$, where $y(t)$ is the solution of (2.3). Then $\lim_{t \rightarrow \infty} x(t) = 0$, where $x(t) = r^{-(t/\tau)} x(t)$, $t \in [-\tau, \infty)$, and $x(t)$ also is the solution of (2.1).

Lemma 2.4 (see [21]). Suppose $\alpha_0 = \max\{\operatorname{Re} \lambda : \lambda - (\ln r/\tau + p) - qe^{-\lambda\tau} = 0\}$ and $y(t)$ is the solution of (2.3). Then, for any $\alpha > \alpha_0$, there is a constant $K = K(\alpha)$ such that

$$|y(t)| \leq Ke^{\alpha t} \|\Psi\|_\tau, \quad t \geq 0, \quad \|\Psi\|_\tau = \sup_{-\tau \leq \theta \leq 0} \left| r^{\theta/\tau+1} \Phi(\theta) \right|. \quad (2.10)$$

Theorem 2.5. Assume that $|r| \exp((p + |q|)\tau) < 1$. Then the zero solution of (2.1) is asymptotically stable.

Proof. First we prove that $\alpha_0 < 0$, where

$$\alpha_0 = \max \left\{ \Re \lambda : \lambda - \left(\frac{\ln r}{\tau} + p \right) - qe^{-\lambda\tau} = 0 \right\}. \quad (2.11)$$

Suppose that $\alpha_0 \geq 0$. Then there exist a λ_0 such that $\Re \lambda_0 \geq 0$ and $\lambda_0 - (\ln r/\tau + p) - qe^{-\lambda_0\tau} = 0$. So

$$\Re \lambda_0 \leq \left(\frac{\ln |r|}{\tau} + p \right) + |q|e^{-\Re \lambda_0 \tau} \leq \frac{\ln |r|}{\tau} + p + |q|. \quad (2.12)$$

And $|r| \exp((p + |q|)\tau) < 1$ implies that

$$\frac{\ln |r|}{\tau} + p + |q| < 0. \quad (2.13)$$

Hence (2.12) and (2.13) imply that $\Re \lambda_0 < 0$, which is a contradiction.

Consequently, $\alpha_0 < 0$. By Lemma 2.4, we know that $\lim_{t \rightarrow \infty} y(t) = 0$, where $y(t)$ the solution of (2.3). By Corollary 2.3, we know that $\lim_{t \rightarrow \infty} x(t) = 0$, where $x(t)$ is the solution of (2.1). $\square$

Corollary 2.6. Assume that there is a constant $\lambda > 0$, such that $|r| < e^{\lambda\tau}$ and $p + |q|e^{\lambda\tau} \leq -\lambda$. Then the zero solution of (2.1) is asymptotically stable.

Proof. Obviously, $p + |q| \leq p + |q|e^{\lambda\tau} \leq -\lambda < 0$. And because $|r| < e^{\lambda\tau}$, so

$$|r|e^{(p+|q|)\tau} \leq |r|e^{-\lambda\tau} < 1. \quad (2.14)$$

By Theorem 2.5, we have that the zero solution of (2.1) is asymptotically stable. $\square$

Similarly, we can get the following corollary.

Corollary 2.7. Assume that there is a constant $\lambda > 0$, such that $|r| < e^{-\lambda\tau}$ and $p + |q||r|^{-1} \leq \lambda$. Then the zero solution of (2.1) is asymptotically stable.

3. Numerical Solutions of IDDEs

We consider the Runge-Kutta methods for (2.3) as follows:

$$\begin{aligned} y_{n+1} &= y_n + h \sum_{i=1}^v b_i \left(\left(\frac{\ln r}{\tau} + p \right) Y_n^i + q Y_{n-m}^i \right), \\ Y_n^i &= y_n + h \sum_{j=1}^v a_{ij} \left(\left(\frac{\ln r}{\tau} + p \right) Y_n^j + q Y_{n-m}^j \right), \end{aligned} \quad (3.1)$$

where $h = 1/m$, is a given stepsize with integer $m > 1$. v is referred to as the number of stages. The weights b_i , the abscissas $c_i = \sum_{j=1}^v a_{ij}$, and the matrix $A = [a_{ij}]_{i,j=1}^v$ will be denoted by (A, b, c) . $y_n = r^{nh/\tau+1} \Phi(nh)$, $n = -m, -m+1, \dots, 0$, and y_n is an approximation of $y(nh)$, $n = 0, 1, \dots$

Define

$$x_n = \Re \left(r^{-\{n/m\}} y_n \right). \quad (3.2)$$

Then x_n is the numerical solution of (2.1).

Theorem 3.1. Assume that the Runge-Kutta method (A, b, c) is of order p_1 . Then the order of (3.1) and (3.2) for (2.1) is also p_1 , when $\Psi \in C^{p_1}([- \tau, 0], R)$.

Proof. In [22], it has been proved that the order of (3.1) is p_1 , if the Runge-Kutta method (A, b, c) is of order p_1 . And for any fixed $T > 0$, and any $t_n = nh \leq T$, we have $|x(t_n) - x_n| \leq \sqrt{(x(t_n) - x_n)^2 + (\text{Im}(r^{-\{n/m\}} y_n))^2} = |x(t_n) - r^{-\{n/m\}} y_n| = |r^{-\{t_n/\tau\}} y(t_n) - r^{-\{n/m\}} y_n| = |r^{-\{n/m\}}| |y(t_n) - y_n| \leq C |y(t_n) - y_n| = O(h^{p_1})$, where $C = \max\{1, |r|^{-1}\}$. Hence the order of (3.1) and (3.2) is p_1 . $\square$

By (3.2), we know that $x_n \rightarrow 0$ and $\text{Im}(r^{-\{n/m\}} y_n) \rightarrow 0$, $n \rightarrow \infty$ if and only if $y_n \rightarrow 0$, $n \rightarrow \infty$. The definition of P -stability region and P -stable has been given in many papers, for example, [23]. The P -stability region of (3.1) is the set S_p of pairs of numbers (β, γ) , $\beta = h(\ln r/\tau + p)$, $\gamma = hq$, such that the numerical solution $\{y_n\}_{n \geq 0}$ satisfies

$$\lim_{n \rightarrow \infty} y_n = 0 \quad (3.3)$$

for all constant delays τ and all initial function Ψ .

The Runge-Kutta method (3.1) is P -stable, if

$$S_p \supseteq \left\{ (\beta, \gamma) \in \mathbb{C}^2 : \text{Re } \beta + |\gamma| < 0 \right\}. \quad (3.4)$$

Therefore we obtain the following theorem.

Theorem 3.2. If $|r| \exp((p+|q|)\tau) < 1$, then the solutions of (2.3) and (2.1) are asymptotically stable. In addition, if the Runge-Kutta method is P -stable, then the process (3.1) preserves asymptotically stable of (2.3); consequently, the process (3.1)-(3.2) preserves asymptotically stable of (2.1).

Figure 1: The analytic solution of (4.1) and the real part analytic solution of (4.3).

4. Numerical Experiments

Consider the following IDDE:

$$\begin{aligned} x'(t) &= -4x(t) + x(t-1), \quad t \geq 0, \quad t \neq k, \quad k = 0, 1, 2, \dots, \\ x(t) &= -2x(t^-), \quad t = k, \quad k = 0, 1, 2, \dots, \\ x(t) &= \Phi(t), \quad t \in [-1, 0). \end{aligned} \quad (4.1)$$

The analytic solution of (4.1) (see Figure 1) as $\Phi(t) \equiv 1, \quad t \in [-1, 0)$ is

$$x(t) = \begin{cases} 1, & t \in [-1, 0), \\ -\frac{9}{4}e^{-4t} + \frac{1}{4}, & t \in [0, 1), \\ \left(\frac{9}{2}e^{-4} - \frac{9}{4}t + \frac{27}{16}\right)e^{-4(t-1)} + \frac{1}{16}, & t \in [1, 2), \\ -2\left(\left(\frac{9}{2}e^{-4} - \frac{9}{4}t + \frac{27}{16}\right)e^{-4} + \frac{1}{16}\right), & t = 2. \end{cases} \quad (4.2)$$

Note that $e_{0.5}$, e_1 , $e_{1.5}$, and e_2 are the errors of the analytic solution and the numerical solution (3.1) (3.2) for (4.1) at $t = 0.5, 1, 1.5$, and 2 , respectively.

In Tables 1 and 2, we have listed the errors at $t = 0.5, 1.1.5$ and 2 of the explicit Euler method and the Trapezoid method.

Figure 2: The numerical solution of (4.1), when (3.1) is the Trapezoid method as $h = 1/100$.

Figure 3: The numerical solution of (4.1), when (3.1) is the implicit Euler method as $h = 1/100$.

Obviously, $|r|e^{(p+|q|)\tau} = 2e^{-4+1} = 2e^{-3} < 1$, so the zero solution of (4.1) is asymptotically stable by Theorem 2.5. Assume that $x(t)$ is the solution of (4.1) and $y(t) = (-2)^{[t]}x(t)$, for $t \in [-1, \infty)$. Then $y(t)$ is the solution of the following equation:

$$\begin{aligned} y'(t) &= (\ln(-2) - 4)y(t) + y(t-1), \quad t \geq 0, \quad t \neq k, \quad k = 0, 1, 2, \dots, \\ y(t) &= (-2)^{t+1}\Phi(t), \quad t \in [-1, 0). \end{aligned} \tag{4.3}$$

Because $\Re(\ln(-2) - 4) + 1 < 0$, we know that $y(t) \rightarrow 0$ as $t \rightarrow \infty$ (see Figure 1).

The numerical process (3.1) and (3.2) for (4.1) is asymptotically stable as (3.2) is the Trapezoid method (see Figure 2) and the implicit Euler method (see Figure 3).

Table 1

Stepsize	$\epsilon_{0.5}$	ϵ_1	$\epsilon_{1.5}$	ϵ_2
$h = 1/50$	0.007237306	0.013957231	0.009751641	0.000998634
$h = 1/100$	0.003480570	0.006575988	0.004535880	0.000421508
$h = 1/200$	0.001706846	0.003192398	0.002188540	0.000193107
$h = 1/400$	0.000845200	0.0015728959	0.001075069	0.000092351
Ratio	2.045995126	2.070658147	2.086056611	2.214324691

Table 2

Stepsize	$\epsilon_{0.5}$	ϵ_1	$\epsilon_{1.5}$	ϵ_2
$h = 1/50$	$3.50207e - 4$	$2.52492e - 4$	$2.21015e - 4$	$2.45482e - 4$
$h = 1/100$	$8.74678e - 5$	$6.30749e - 5$	$5.52490e - 5$	$6.13071e - 5$
$h = 1/200$	$2.18617e - 5$	$1.57657e - 5$	$1.38119e - 5$	$1.53228e - 5$
$h = 1/400$	$5.46510e - 6$	$3.94124e - 6$	$3.45297e - 6$	$3.83045e - 6$
Ratio	4.00167	4.00133	4.00015	4.00181

Acknowledgment

The authors wish to thank referees for valuable comments. The research was supported by the NSF of China no. 11071050.

References

- [1] A. Anokhin, L. Berezansky, and E. Braverman, "Exponential stability of linear delay impulsive differential equations," *Journal of Mathematical Analysis and Applications*, vol. 193, no. 3, pp. 923–941, 1995.
- [2] G. Ballinger and X. Liu, "Existence and uniqueness results for impulsive delay differential equations," *Dynamics of Continuous, Discrete and Impulsive Systems*, vol. 5, no. 1–4, pp. 579–591, 1999.
- [3] H. Bereketoglu, G. Seyhan, and A. Ogun, "Advanced impulsive differential equations with piecewise constant arguments," *Mathematical Modelling and Analysis*, vol. 15, no. 2, pp. 175–187, 2010.
- [4] L. Berezansky and L. Idels, "Exponential stability of some scalar impulsive delay differential equations," *Communications in Applied Analysis*, vol. 2, no. 3, pp. 301–308, 1998.
- [5] V. Covachev, H. Akça, and F. Yeniçerioglu, "Difference approximations for impulsive differential equations," *Applied Mathematics and Computation*, vol. 121, no. 2–3, pp. 383–390, 2001.
- [6] M. De la Sen, "Stability of impulsive time-varying systems and compactness of the operators mapping the input space into the state and output spaces," *Journal of Mathematical Analysis and Applications*, vol. 321, no. 2, pp. 621–650, 2006.
- [7] M. De la Sen and N. Luo, "On the uniform exponential stability of a wide class of linear time-delay systems," *Journal of Mathematical Analysis and Applications*, vol. 289, no. 2, pp. 456–476, 2004.
- [8] C. Z. Li, Y. K. Li, and Y. Ye, "Exponential stability of fuzzy Cohen-Grossberg neural networks with time delays and impulsive effects," *Communications in Nonlinear Science and Numerical Simulation*, vol. 15, no. 11, pp. 3599–3606, 2010.
- [9] M. Z. Liu, H. Liang, and Z. W. Yang, "Stability of Runge-Kutta methods in the numerical solution of linear impulsive differential equations," *Applied Mathematics and Computation*, vol. 192, no. 2, pp. 346–357, 2007.
- [10] X. Liu, "Stability results for impulsive differential systems with applications to population growth models," *Dynamics and Stability of Systems*, vol. 9, no. 2, pp. 163–174, 1994.
- [11] X. Liu, X. Shen, and Y. Zhang, "Exponential stability of singularly perturbed systems with time delay," *Applicable Analysis*, vol. 82, no. 2, pp. 117–130, 2003.
- [12] X. Z. Liu, X. M. Shen, Y. Zhang, and Q. Wang, "Stability criteria for impulsive systems with time delay and unstable system matrices," *IEEE Transactions on Circuits and Systems I*, vol. 54, no. 10, pp. 2288–2298, 2007.

- [13] I. M. Stamova and G. T. Stamov, "Lyapunov-Razumikhin method for impulsive functional differential equations and applications to the population dynamics," *Journal of Computational and Applied Mathematics*, vol. 130, no. 1-2, pp. 163–171, 2001.
- [14] J. H. Shen, "Razumikhin techniques in impulsive functional-differential equations," *Nonlinear Analysis. Theory, Methods & Applications A*, vol. 36, pp. 119–130, 1999.
- [15] J. Shen and J. Yan, "Razumikhin type stability theorems for impulsive functional-differential equations," *Nonlinear Analysis. Theory, Methods & Applications*, vol. 33, no. 5, pp. 519–531, 1998.
- [16] Q. Wang and X. Liu, "Exponential stability for impulsive delay differential equations by Razumikhin method," *Journal of Mathematical Analysis and Applications*, vol. 309, no. 2, pp. 462–473, 2005.
- [17] Q. Wang and X. Liu, "Impulsive stabilization of delay differential systems via the Lyapunov-Razumikhin method," *Applied Mathematics Letters*, vol. 20, no. 8, pp. 839–845, 2007.
- [18] Y. Zhang and J. Sun, "Stability of impulsive functional differential equations," *Nonlinear Analysis. Theory, Methods & Applications A*, vol. 68, no. 12, pp. 3665–3678, 2008.
- [19] X. J. Ran, M. Z. Liu, and Q. Y. Zhu, "Numerical methods for impulsive differential equation," *Mathematical and Computer Modelling*, vol. 48, no. 1-2, pp. 46–55, 2008.
- [20] X. Ding, K. Wu, and M. Liu, "The Euler scheme and its convergence for impulsive delay differential equations," *Applied Mathematics and Computation*, vol. 216, no. 5, pp. 1566–1570, 2010.
- [21] J. K. Hale, *Introduction to Functional Equations*, New York, NY, USA, 1993.
- [22] E. Hairer, S. P. Nørsett, and G. Wanner, *Solving Ordinary Differential Equations. I*, Berlin, Germany, 1991.
- [23] A. Bellen and M. Zennaro, *Numerical Methods for Delay Differential Equations*, Oxford University Press, Oxford, UK, 2003.