Research Article

Strong Convergence for Hybrid S-Iteration Scheme

Shin Min Kang,¹ Arif Rafiq,² and Young Chel Kwun³

¹ Department of Mathematics and RINS, Gyeongsang National University, Jinju 660-701, Republic of Korea

² School of CS and Mathematics, Hajvery University, 43-52 Industrial Area, Gulberg-III, Lahore 54660, Pakistan

³ Department of Mathematics, Dong-A University, Pusan 614-714, Republic of Korea

Correspondence should be addressed to Young Chel Kwun; yckwun@dau.ac.kr

Received 19 November 2012; Accepted 4 February 2013

Academic Editor: D. R. Sahu

Copyright © 2013 Shin Min Kang et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We establish a strong convergence for the hybrid S-iterative scheme associated with nonexpansive and Lipschitz strongly pseudocontractive mappings in real Banach spaces.

1. Introduction and Preliminaries

Let E be a real Banach space and let K be a nonempty convex subset of E . Let J denote the normalized duality mapping from E to 2^{E^*} defined by

$$J(x) = \{f^* \in E^* : \langle x, f^* \rangle = \|x\|^2, \|f^*\| = \|x\|\}, \quad \forall x, y \in E, \quad (1)$$

where E^* denotes the dual space of E and $\langle \cdot, \cdot \rangle$ denotes the generalized duality pairing. We will denote the single-valued duality map by j .

Let $T : K \rightarrow K$ be a mapping.

Definition 1. The mapping T is said to be *Lipschitzian* if there exists a constant $L > 1$ such that

$$\|Tx - Ty\| \leq L \|x - y\|, \quad \forall x, y \in K. \quad (2)$$

Definition 2. The mapping T is said to be *nonexpansive* if

$$\|Tx - Ty\| \leq \|x - y\|, \quad \forall x, y \in K. \quad (3)$$

Definition 3. The mapping T is said to be *pseudocontractive* if for all $x, y \in K$, there exists $j(x - y) \in J(x - y)$ such that

$$\langle Tx - Ty, j(x - y) \rangle \leq \|x - y\|^2. \quad (4)$$

Definition 4. The mapping T is said to be *strongly pseudocontractive* if for all $x, y \in K$, there exists $k \in (0, 1)$ such that

$$\langle Tx - Ty, j(x - y) \rangle \leq k \|x - y\|^2. \quad (5)$$

Let K be a nonempty convex subset C of a normed space E .

(a) The sequence $\{x_n\}$ defined by, for arbitrary $x_1 \in K$,

$$\begin{aligned} x_{n+1} &= (1 - \alpha_n)x_n + \alpha_n Ty_n, \\ y_n &= (1 - \beta_n)x_n + \beta_n Tx_n, \quad n \geq 1, \end{aligned} \quad (6)$$

where $\{\alpha_n\}$ and $\{\beta_n\}$ are sequences in $[0, 1]$, is known as the Ishikawa iteration process [1].

If $\beta_n = 0$ for $n \geq 1$, then the Ishikawa iteration process becomes the Mann iteration process [2].

(b) The sequence $\{x_n\}$ defined by, for arbitrary $x_1 \in K$,

$$\begin{aligned} x_{n+1} &= Ty_n, \\ y_n &= (1 - \beta_n)x_n + \beta_n Tx_n, \quad n \geq 1, \end{aligned} \quad (7)$$

where $\{\beta_n\}$ is a sequence in $[0, 1]$, is known as the S-iteration process [3, 4].

In the last few years or so, numerous papers have been published on the iterative approximation of fixed points of Lipschitz *strongly pseudocontractive* mappings using the *Ishikawa iteration scheme* (see, e.g., [1]). Results which had

been known only in Hilbert spaces and only for Lipschitz mappings have been extended to more general Banach spaces (see, e.g., [5–10] and the references cited therein).

In 1974, Ishikawa [1] proved the following result.

Theorem 5. Let K be a compact convex subset of a Hilbert space H and let $T : K \rightarrow K$ be a Lipschitzian pseudocontractive mapping. For arbitrary $x_1 \in K$, let $\{x_n\}$ be a sequence defined iteratively by

$$\begin{aligned} x_{n+1} &= (1 - \alpha_n)x_n + \alpha_n T y_n, \\ y_n &= (1 - \beta_n)x_n + \beta_n T x_n, \quad n \geq 1, \end{aligned} \quad (8)$$

where $\{\alpha_n\}$ and $\{\beta_n\}$ are sequences satisfying

- (i) $0 \leq \alpha_n \leq \beta_n \leq 1$,
- (ii) $\lim_{n \rightarrow \infty} \beta_n = 0$,
- (iii) $\sum_{n \geq 1} \alpha_n \beta_n = \infty$.

Then the sequence $\{x_n\}$ converges strongly at a fixed point of T .

In [6], Chidume extended the results of Schu [9] from Hilbert spaces to the much more general class of real Banach spaces and approximated the fixed points of (strongly) pseudocontractive mappings.

In [11], Zhou and Jia gave the more general answer of the question raised by Chidume [5] and proved the following.

If X is a real Banach space with a uniformly convex dual X^* , K is a nonempty bounded closed convex subset of X , and $T : K \rightarrow K$ is a continuous strongly pseudocontractive mapping, then the Ishikawa iteration scheme converges strongly at the unique fixed point of T .

In this paper, we establish the strong convergence for the hybrid S -iterative scheme associated with nonexpansive and Lipschitz strongly pseudocontractive mappings in real Banach spaces. We also improve the result of Zhou and Jia [11].

2. Main Results

We will need the following lemmas.

Lemma 6 (see [12]). Let $J : E \rightarrow 2^E$ be the normalized duality mapping. Then for any $x, y \in E$, one has

$$\begin{aligned} \|x + y\|^2 &\leq \|x\|^2 + 2 \langle y, j(x + y) \rangle, \\ \forall j(x + y) &\in J(x + y). \end{aligned} \quad (9)$$

Lemma 7 (see [10]). Let $\{\rho_n\}$ be nonnegative sequence satisfying

$$\rho_{n+1} \leq (1 - \theta_n) \rho_n + \omega_n, \quad (10)$$

where $\theta_n \in [0, 1]$, $\sum_{n \geq 1} \theta_n = \infty$, and $\omega_n = o(\theta_n)$. Then

$$\lim_{n \rightarrow \infty} \rho_n = 0. \quad (11)$$

The following is our main result.

Theorem 8. Let K be a nonempty closed convex subset of a real Banach space E , let $S : K \rightarrow K$ be nonexpansive, and let $T : K \rightarrow K$ be Lipschitz strongly pseudocontractive mappings such that $p \in F(S) \cap F(T) = \{x \in K : Sx = Tx = x\}$ and

$$\begin{aligned} \|x - Sy\| &\leq \|Sx - Sy\|, \quad \forall x, y \in K, \\ \|x - Ty\| &\leq \|Tx - Ty\|, \quad \forall x, y \in K. \end{aligned} \quad (C)$$

Let $\{\beta_n\}$ be a sequence in $[0, 1]$ satisfying

- (iv) $\sum_{n \geq 1} \beta_n = \infty$,
- (v) $\lim_{n \rightarrow \infty} \beta_n = 0$.

For arbitrary $x_1 \in K$, let $\{x_n\}$ be a sequence iteratively defined by

$$x_{n+1} = Sy_n, \quad (12)$$

$$y_n = (1 - \beta_n)x_n + \beta_n T x_n, \quad n \geq 1.$$

Then the sequence $\{x_n\}$ converges strongly at the common fixed point p of S and T .

Proof. For strongly pseudocontractive mappings, the existence of a fixed point follows from Delmling [13]. It is shown in [11] that the set of fixed points for strongly pseudocontractions is a singleton.

By (v), since $\lim_{n \rightarrow \infty} \beta_n = 0$, there exists $n_0 \in \mathbb{N}$ such that for all $n \geq n_0$,

$$\beta_n \leq \min \left\{ \frac{1}{4k}, \frac{1 - k}{(1 + L)(1 + 3L)} \right\}, \quad (13)$$

where $k < 1/2$. Consider

$$\begin{aligned} \|x_{n+1} - p\|^2 &= \langle x_{n+1} - p, j(x_{n+1} - p) \rangle \\ &= \langle Sy_n - p, j(x_{n+1} - p) \rangle \\ &= \langle Tx_{n+1} - p, j(x_{n+1} - p) \rangle \\ &\quad + \langle Sy_n - Tx_{n+1}, j(x_{n+1} - p) \rangle \\ &\leq k \|x_{n+1} - p\|^2 + \|Sy_n - Tx_{n+1}\| \|x_{n+1} - p\|, \end{aligned} \quad (14)$$

which implies that

$$\|x_{n+1} - p\| \leq \frac{1}{1 - k} \|Sy_n - Tx_{n+1}\|, \quad (15)$$

where

$$\begin{aligned} \|Sy_n - Tx_{n+1}\| &\leq \|Sy_n - Ty_n\| + \|Ty_n - Tx_{n+1}\| \\ &\leq \|x_n - Sy_n\| + \|x_n - Ty_n\| + \|Ty_n - Tx_{n+1}\| \\ &\leq \|Sx_n - Sy_n\| + \|Tx_n - Ty_n\| + \|Ty_n - Tx_{n+1}\| \\ &\leq \|Sx_n - Sy_n\| + L(\|x_n - y_n\| + \|y_n - x_{n+1}\|), \end{aligned} \quad (16)$$

$$\begin{aligned} \|y_n - x_{n+1}\| &\leq \|y_n - x_n\| + \|x_n - x_{n+1}\| \\ &= \|y_n - x_n\| + \|x_n - Sy_n\| \\ &\leq \|y_n - x_n\| + \|Sx_n - Sy_n\|, \end{aligned} \quad (17)$$

and consequently from (16), we obtain

$$\begin{aligned}\|S y_n - T x_{n+1}\| &\leq (1+L)\|S x_n - S y_n\| + 2L\|x_n - y_n\| \\ &\leq (1+3L)\|x_n - y_n\| \\ &= (1+3L)\beta_n\|x_n - T x_n\| \\ &\leq (1+L)(1+3L)\beta_n\|x_n - p\|.\end{aligned}\quad (18)$$

Substituting (18) in (15) and using (13), we get

$$\begin{aligned}\|x_{n+1} - p\| &\leq \frac{(1+L)(1+3L)}{1-k}\beta_n\|x_n - p\| \\ &\leq \|x_n - p\|.\end{aligned}\quad (19)$$

So, from the above discussion, we can conclude that the sequence $\{x_n - p\}$ is bounded. Since T is Lipschitzian, so $\{T x_n - p\}$ is also bounded. Let $M_1 = \sup_{n \geq 1} \|x_n - p\| + \sup_{n \geq 1} \|T x_n - p\|$. Also by (ii), we have

$$\begin{aligned}\|x_n - y_n\| &= \beta_n\|x_n - T x_n\| \\ &\leq M_1\beta_n \\ &\longrightarrow 0\end{aligned}\quad (20)$$

as $n \rightarrow \infty$, implying that $\{x_n - y_n\}$ is bounded, so let $M_2 = \sup_{n \geq 1} \|x_n - y_n\| + M_1$. Further,

$$\begin{aligned}\|y_n - p\| &\leq \|y_n - x_n\| + \|x_n - p\| \\ &\leq M_2,\end{aligned}\quad (21)$$

which implies that $\{y_n - p\}$ is bounded. Therefore, $\{T y_n - p\}$ is also bounded.

Set

$$M_3 = \sup_{n \geq 1} \|y_n - p\| + \sup_{n \geq 1} \|T y_n - p\|.\quad (22)$$

Denote $M = M_1 + M_2 + M_3$. Obviously, $M < \infty$. Now from (12) for all $n \geq 1$, we obtain

$$\|x_{n+1} - p\|^2 = \|S y_n - p\|^2 \leq \|y_n - p\|^2,\quad (23)$$

and by Lemma 6, we get

$$\begin{aligned}\|y_n - p\|^2 &= \|(1-\beta_n)x_n + \beta_n T x_n - p\|^2 \\ &= \|(1-\beta_n)(x_n - p) + \beta_n(T x_n - p)\|^2 \\ &\leq (1-\beta_n)^2\|x_n - p\|^2 + 2\beta_n\langle T x_n - p, j(y_n - p) \rangle \\ &= (1-\beta_n)^2\|x_n - p\|^2 + 2\beta_n\langle T y_n - p, j(y_n - p) \rangle \\ &\quad + 2\beta_n\langle T x_n - T y_n, j(y_n - p) \rangle \\ &\leq (1-\beta_n)^2\|x_n - p\|^2 + 2k\beta_n\|y_n - p\|^2 \\ &\quad + 2\beta_n\|T x_n - T y_n\|\|y_n - p\| \\ &\leq (1-\beta_n)^2\|x_n - p\|^2 + 2k\beta_n\|y_n - p\|^2 \\ &\quad + 2ML\beta_n\|x_n - y_n\|,\end{aligned}\quad (24)$$

which implies that

$$\begin{aligned}\|y_n - p\|^2 &\leq \frac{(1-\beta_n)^2}{1-2k\beta_n}\|x_n - p\|^2 + \frac{2ML\beta_n}{1-2k\beta_n}\|x_n - y_n\| \\ &\leq (1-\beta_n)\|x_n - p\|^2 + 4ML\beta_n\|x_n - y_n\|\end{aligned}\quad (25)$$

because by (13), we have $((1-\beta_n)/(1-2k\beta_n)) \leq 1$ and $(1/(1-2k\beta_n)) \leq 2$. Hence, (23) gives us

$$\|x_{n+1} - p\|^2 \leq (1-\beta_n)\|x_n - p\|^2 + 4ML\beta_n\|x_n - y_n\|.\quad (26)$$

For all $n \geq 1$, put

$$\begin{aligned}\rho_n &= \|x_n - p\|, \\ \theta_n &= \beta_n, \\ \omega_n &= 4ML\beta_n\|x_n - y_n\|,\end{aligned}\quad (27)$$

then according to Lemma 7, we obtain from (26) that

$$\lim_{n \rightarrow \infty} \|x_n - p\| = 0.\quad (28)$$

This completes the proof. $\square$

Corollary 9. Let K be a nonempty closed convex subset of a real Hilbert space H , let $S : K \rightarrow K$ be nonexpansive, and let $T : K \rightarrow K$ be Lipschitz strongly pseudocontractive mappings such that $p \in F(S) \cap F(T)$ and the condition (C). Let $\{\beta_n\}$ be a sequence in $[0, 1]$ satisfying the conditions (iv) and (v).

For arbitrary $x_1 \in K$, let $\{x_n\}$ be a sequence iteratively defined by (12). Then the sequence $\{x_n\}$ converges strongly at the common fixed point p of S and T .

Example 10. As a particular case, we may choose, for instance, $\beta_n = 1/n$.

Remark 11. (1) The condition (C) is not new and it is due to Liu et al. [14].

(2) We prove our results for a hybrid iteration scheme, which is simple in comparison to the previously known iteration schemes.

Acknowledgment

This study was supported by research funds from Dong-A University.

References

- [1] S. Ishikawa, "Fixed points by a new iteration method," *Proceedings of the American Mathematical Society*, vol. 44, pp. 147-150, 1974.
- [2] W. R. Mann, "Mean value methods in iteration," *Proceedings of the American Mathematical Society*, vol. 4, pp. 506-510, 1953.
- [3] D. R. Sahu, "Applications of the S-iteration process to constrained minimization problems and split feasibility problems," *Fixed Point Theory*, vol. 12, no. 1, pp. 187-204, 2011.

- [4] D. R. Sahu and A. Petruşel, "Strong convergence of iterative methods by strictly pseudocontractive mappings in Banach spaces," *Nonlinear Analysis. Theory, Methods & Applications*, vol. 74, no. 17, pp. 6012–6023, 2011.
- [5] C. E. Chidume, "Approximation of fixed points of strongly pseudocontractive mappings," *Proceedings of the American Mathematical Society*, vol. 120, no. 2, pp. 545–551, 1994.
- [6] C. E. Chidume, "Iterative approximation of fixed points of Lipschitz pseudocontractive maps," *Proceedings of the American Mathematical Society*, vol. 129, no. 8, pp. 2245–2251, 2001.
- [7] C. E. Chidume and C. Moore, "Fixed point iteration for pseudocontractive maps," *Proceedings of the American Mathematical Society*, vol. 127, no. 4, pp. 1163–1170, 1999.
- [8] C. E. Chidume and H. Zegeye, "Approximate fixed point sequences and convergence theorems for Lipschitz pseudocontractive maps," *Proceedings of the American Mathematical Society*, vol. 132, no. 3, pp. 831–840, 2004.
- [9] J. Chu, "Approximating fixed points of Lipschitzian pseudocontractive mappings," *Houston Journal of Mathematics*, vol. 19, no. 1, pp. 107–115, 1993.
- [10] X. Weng, "Fixed point iteration for local strictly pseudocontractive mapping," *Proceedings of the American Mathematical Society*, vol. 113, no. 3, pp. 727–731, 1991.
- [11] H. Zhou and Y. Jia, "Approximation of fixed points of strongly pseudocontractive maps without Lipschitz assumption," *Proceedings of the American Mathematical Society*, vol. 125, no. 6, pp. 1705–1709, 1997.
- [12] S. S. Chang, "Some problems and results in the study of nonlinear analysis," *Nonlinear Analysis*, vol. 30, no. 7, pp. 4197–4208, 1997.
- [13] K. Delmling, "Zeros of accretive operators," *Manuscripta Mathematica*, vol. 13, pp. 283–288, 1974.
- [14] Z. Liu, C. Feng, J. S. Ume, and S. M. Kang, "Weak and strong convergence for common fixed points of a pair of nonexpansive and asymptotically nonexpansive mappings," *Taiwanese Journal of Mathematics*, vol. 11, no. 1, pp. 27–42, 2007.