Erratum

Erratum to “Ranks of a Constrained Hermitian Matrix Expression with Applications”

Shao-Wen Yu

Department of Mathematics, East China University of Science and Technology, Shanghai 200237, China

Correspondence should be addressed to Shao-Wen Yu; yushaowen@ecust.edu.cn

Received 14 June 2013; Accepted 8 July 2013

Copyright © 2013 Shao-Wen Yu. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The equalities (15) and (16) should be deleted from Lemma 3 which will be reformatted as follows.

Lemma 3. Let $A = A^* \in \mathbb{H}^{m \times m}$, $B \in \mathbb{H}^{m \times n}$, and $C \in \mathbb{H}^{p \times m}$ be given. Assume that $\mathcal{R}(B) \subseteq \mathcal{R}(C^*)$. Then,

$$\begin{aligned} & \max_X r[A - BXC - (BXC)^*] \\ &= \min \left\{ r[A \ C^*], r \begin{bmatrix} A & B \\ B^* & 0 \end{bmatrix} \right\}, \\ & \min_X r[A - BXC - (BXC)^*] \\ &= 2r[A \ C^*] + r \begin{bmatrix} A & B \\ B^* & 0 \end{bmatrix} - 2r \begin{bmatrix} A & B \\ C & 0 \end{bmatrix}. \end{aligned} \tag{17}$$