Research Article

Nonlinear Super Integrable Couplings of Super Classical-Boussinesq Hierarchy

Xiuzhi Xing,¹ Jingzhu Wu,¹ and Xianguo Geng²

¹ Department of Mathematics, Zhoukou Normal University, Zhoukou 466000, China

² Department of Mathematics, Zhengzhou University, Zhengzhou 450052, China

Correspondence should be addressed to Xiuzhi Xing; xingxiuzhi@zknz.edu.cn

Received 15 December 2013; Accepted 26 February 2014; Published 7 April 2014

Academic Editor: Chein-Shan Liu

Copyright © 2014 Xiuzhi Xing et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Nonlinear integrable couplings of super classical-Boussinesq hierarchy based upon an enlarged matrix Lie super algebra were constructed. Then, its super Hamiltonian structures were established by using super trace identity. As its reduction, nonlinear integrable couplings of the classical integrable hierarchy were obtained.

1. Introduction

With the development of soliton theory, super integrable systems associated with Lie super algebra have aroused growing attentions by many mathematicians and physicists. It was known that super integrable systems contained the odd variables, which would provide more prolific fields for mathematical researchers and physical ones. Several super integrable systems including super AKNS hierarchy, super KdV hierarchy, super C-KdV hierarchy, and super classical-Boussinesq hierarchy have been studied [1–4]. There are some interesting results on the super integrable systems, such as Darboux transformation [5], super Hamiltonian structures [6, 7], binary nonlinearization [8], and reciprocal transformation [9].

The research of integrable couplings of the well-known integrable hierarchy has received considerable attentions [10–15]. A few approaches to construct linear integrable couplings of the classical soliton equation are presented by permutation, enlarging spectral problem, using matrix Lie algebra constructing new loop Lie algebra [16], and creating semidirect sums of Lie algebra. Recently, Ma and Zhu [17, 18] presented a scheme for constructing nonlinear continuous and discrete integrable couplings using the block type matrix algebra. However, there is one interesting question for us which is how to generate nonlinear super integrable couplings for the super integrable hierarchy.

In this paper, we would like to construct nonlinear super integrable couplings of the super soliton equations through enlarging matrix Lie super algebra. We take the Lie algebra $sl(2, 1)$ as an example to illustrate the approach for extending Lie super algebra. Based on the enlarged Lie super algebra $sl(4, 1)$, we work out nonlinear super integrable Hamiltonian couplings of the super classical-Boussinesq hierarchy. Finally, we will reduce the nonlinear super classical-Boussinesq integrable Hamiltonian couplings to some special cases.

2. Enlargement of Lie Superalgebra

Consider the Lie superalgebra $sl(2, 1)$. Its basis is

$$\begin{aligned} E_1 &= \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 0 \end{pmatrix}, & E_2 &= \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \\ E_3 &= \begin{pmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, & E_4 &= \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & -1 & 0 \end{pmatrix}, \\ E_5 &= \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}, \end{aligned} \quad (1)$$

where E_1, E_2, E_3 are even elements and E_4, E_5 are odd ones. Their nonzero (anti)commutation relations are

$$\begin{aligned} [E_1, E_2] &= 2E_2, & [E_1, E_3] &= -2E_3, \\ [E_2, E_3] &= E_1, & [E_1, E_4] &= [E_2, E_5] = E_4, \\ [E_1, E_5] &= [E_4, E_3] = -E_5, & [E_4, E_5] &= E_1, \\ [E_4, E_4] &= -2E_2, & [E_5, E_5] &= 2E_3. \end{aligned} \quad (2)$$

Let us enlarge the Lie superalgebra $\mathfrak{sl}(2, 1)$ to the Lie superalgebra $\mathfrak{sl}(4, 1)$ with a basis

$$\begin{aligned} e_1 &= \begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}, & e_2 &= \begin{pmatrix} 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}, \\ e_3 &= \begin{pmatrix} 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}, & e_4 &= \begin{pmatrix} 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & -1 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}, \\ e_5 &= \begin{pmatrix} 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}, & e_6 &= \begin{pmatrix} 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}, \\ e_7 &= \begin{pmatrix} 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & -1 & 0 & 1 & 0 \end{pmatrix}, & e_8 &= \begin{pmatrix} 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & -1 & 0 & 0 \end{pmatrix}, \end{aligned} \quad (3)$$

where $e_1, e_2, e_3, e_4, e_5, e_6$ are even and e_7, e_8 are odd. The generators of the Lie superalgebra $\mathfrak{sl}(4, 1)$, $e_i, 0 \leq i \leq 8$, satisfy the following (anti)commutation relations:

$$\begin{aligned} [e_1, e_2] &= 2e_2, & [e_1, e_3] &= -2e_3, \\ [e_1, e_5] &= 2e_5, & [e_1, e_6] &= -2e_6, & [e_1, e_7] &= e_7, \\ [e_1, e_8] &= -e_8, & [e_2, e_3] &= e_1, \\ [e_2, e_4] &= -2e_5, & [e_2, e_6] &= e_4, & [e_2, e_8] &= e_7, \\ [e_3, e_4] &= 2e_6, & [e_3, e_5] &= -e_4, \\ [e_3, e_7] &= e_8, & [e_4, e_5] &= 2e_5, \\ [e_4, e_6] &= -2e_6, & [e_5, e_6] &= e_4, \\ [e_7, e_8] &= e_1 - e_4, & [e_7, e_7] &= 2e_5 - 2e_2, \end{aligned}$$

$$[e_8, e_8] = 2e_3 - 2e_6,$$

$$[e_1, e_4] = [e_2, e_5] = [e_2, e_7] = [e_3, e_6] = [e_3, e_8] = 0,$$

$$[e_4, e_7] = [e_4, e_8] = [e_5, e_7] = [e_5, e_8]$$

$$= [e_6, e_7] = [e_6, e_8] = 0. \quad (4)$$

Define a loop superalgebra corresponding to the Lie superalgebra $\mathfrak{sl}(4, 1)$, denoted by

$$\begin{aligned} \widetilde{\mathfrak{sl}}(4, 1) &= \mathfrak{sl}(4, 1) \otimes [\lambda, \lambda^{-1}] \\ &= \{e_i \lambda^m, e_i \in \mathfrak{sl}(4, 1), i = 1, 2, \dots, 8; m = 0, \pm 1, \pm 2, \dots\}. \end{aligned} \quad (5)$$

The corresponding (anti)commutative relations are given as

$$[e_i \lambda^m, e_j \lambda^n] = [e_i, e_j] \lambda^{m+n}, \quad \forall e_i, e_j \in \mathfrak{sl}(4, 1). \quad (6)$$

3. Nonlinear Super Integrable Couplings of the Super Classical-Boussinesq Hierarchy

Let us start from an enlarged spectral problem associated with $\mathfrak{sl}(4, 1)$

$$\varphi_x = U\varphi, \quad (7)$$

where

U

$$\begin{aligned} &= -e_1(1) - \frac{1}{q}e_1(0) + re_2(0) - e_3(0) - u_1e_4(0) \\ &\quad + u_2e_5(0) + \alpha e_7(0) + \beta e_8(0) \end{aligned}$$

$$= \begin{pmatrix} -\lambda - \frac{1}{4}q & r & -u_1 & u_2 & \alpha \\ -1 & \lambda + \frac{1}{4}q & 0 & u_1 & \beta \\ 0 & 0 & -\lambda - \frac{1}{4}q - u_1 & r + u_2 & 0 \\ 0 & 0 & -1 & \lambda + \frac{1}{4}q + u_1 & 0 \\ \beta & -\alpha & -\beta & \alpha & 0 \end{pmatrix}, \quad (8)$$

where r, s, u_1 , and u_2 are even potentials but α and β are odd ones. In order to obtain super integrable couplings of super integrable hierarchy, we first solve the adjoint representation (8)

$$V_x = [U, V] \quad (9)$$

with

$$V = Ae_1(0) + Be_2(0) + Ce_3(0) + Ee_4(0) + Fe_5(0) + Ge_6(0) + \rho e_7(0) + \delta e_8(0)$$

$$= \begin{pmatrix} A & B & E & F & \rho \\ C & -A & G & -E & \delta \\ 0 & 0 & A+E & B+F & 0 \\ 0 & 0 & C+G & -A-E & 0 \\ \delta & -\rho & -\delta & \rho & 0 \end{pmatrix}, \quad (10)$$

where A, B, C, E, F , and G are commuting fields, ρ and δ are anticommuting fields, and α, β are anticommuting fields.

Substituting

$$A = \sum_{m \geq 0} A_m \lambda^{-m}, \quad B = \sum_{m \geq 0} B_m \lambda^{-m},$$

$$C = \sum_{m \geq 0} C_m \lambda^{-m}, \quad F = \sum_{m \geq 0} F_m \lambda^{-m},$$

$$G = \sum_{m \geq 0} G_m \lambda^{-m}, \quad \rho = \sum_{m \geq 0} \rho_m \lambda^{-m},$$

$$\sigma = \sum_{m \geq 0} \sigma_m \lambda^{-m} \quad (11)$$

into the above equation gives the following recursive formulas:

$$A_{m,x} = B_m + rC_m - \beta\rho_m + \alpha\delta_m,$$

$$B_{m,x} = -2rA_m - 2B_{m+1} - \frac{1}{2}qB_m - 2\alpha\rho_m,$$

$$C_{m,x} = -2A_m + 2C_{m+1} + \frac{1}{2}qC_m + 2\beta\delta_m,$$

$$E_{m,x} = u_2C_m + F_m + rG_m + u_2G_m - \alpha\delta_m - \beta\rho_m,$$

$$F_{m,x} = -2u_2A_m - 2u_1B_m - 2rE_m - 2u_2E_m - 2F_{m+1} - \frac{1}{2}qF_m - 2u_1F_m + 2\alpha\rho_m,$$

$$G_{m,x} = 2u_1C_m - 2E_m + 2G_{m+1} + \frac{1}{2}qG_m + 2u_1G_m - 2\beta\delta_m,$$

$$\rho_{m,x} = -\alpha A_m - \beta B_m - \rho_{m+1} - \frac{1}{4}q\rho_m + r\delta_m,$$

$$\delta_{m,x} = \beta A_m - \alpha C_m - \rho_m + \delta_{m+1} + \frac{1}{4}q\delta_m. \quad (12)$$

From these equations, we can successively deduce

$$A_0 = -1, \quad B_0 = C_0 = F_0 = G_0 = \rho_0 = \delta_0 = 0,$$

$$E_0 = \varepsilon = \text{const}, \quad A_1 = 0,$$

$$B_1 = r, \quad C_1 = -1,$$

$$F_1 = u_2 - r\varepsilon - u_2\varepsilon, \quad E_1 = 0,$$

$$G_1 = \varepsilon, \quad \rho_1 = \alpha, \quad \delta_1 = \beta,$$

$$A_2 = -\frac{1}{2}r + \alpha\beta, \quad B_2 = -\frac{1}{2}r_x - \frac{1}{4}qr,$$

$$C_2 = \frac{1}{4}q,$$

$$F_2 = -\frac{1}{2}u_{2x} + \frac{1}{2}\varepsilon r_x + \frac{1}{2}\varepsilon u_{2x} - \frac{1}{4}qu_2 + \frac{1}{4}q\varepsilon u_2 + \frac{1}{4}qr\varepsilon$$

$$- u_1r - u_1u_2 + \varepsilon u_1u_2 + \varepsilon r u_1,$$

$$E_2 = -\frac{1}{2}u_2 + \frac{1}{2}\varepsilon r + \frac{1}{2}\varepsilon u_2 - \alpha\beta,$$

$$G_2 = u_1 - \frac{1}{4}q\varepsilon - u_1\varepsilon, \quad \rho_2 = -\alpha_x - \frac{1}{4}q\alpha,$$

$$\delta_2 = \beta_x - \frac{1}{4}q\beta,$$

$$A_3 = \frac{1}{2}r_x + \frac{1}{4}qr - \frac{1}{2}q\alpha\beta - \beta\alpha_x + \alpha\beta_x,$$

$$B_3 = \frac{1}{4}r_{xx} + \frac{1}{8}q_xr + \frac{1}{4}qr_x + \frac{1}{2}r^2 - r\alpha\beta + \frac{1}{16}q^2r + \alpha\alpha_x,$$

$$C_3 = \frac{1}{8}q_x - \frac{1}{2}r + \alpha\beta - \frac{1}{16}q^2 - \beta\beta_x,$$

$$E_3 = (1 - \varepsilon) \left[u_1u_2 + u_1r + \frac{1}{4}qu_2 + \frac{1}{4}u_{2x} \right]$$

$$- \alpha\beta_x + \beta\alpha_x + \frac{1}{2}q\alpha\beta - \frac{1}{4}qr - \frac{1}{4}\varepsilon r_x,$$

$$F_3 = (1 - \varepsilon) \left[\frac{1}{4}u_{2xx} + \frac{1}{8}q_xu_2 + \frac{1}{4}qu_{2x} + \frac{1}{2}u_{1x}r + u_1r_x \right.$$

$$+ \frac{1}{2}u_{1x}u_2 + u_1u_{2x} + \frac{1}{16}q^2u_2 + \frac{1}{2}qu_1r$$

$$+ \left. \frac{1}{2}qu_1u_2 + \frac{1}{2}u_2^2 + ru_2 + ru_1^2 + u_1^2u_2 \right]$$

$$- \frac{1}{4}\varepsilon r_{xx} - \frac{1}{8}\varepsilon q_xr - \frac{1}{4}\varepsilon qr_x + r\alpha\beta - \frac{1}{2}\varepsilon r^2$$

$$- \frac{1}{16}\varepsilon q^2r - \alpha\alpha_x,$$

$$G_3 = \frac{1}{2}u_{1x} - \frac{1}{8}q_x\varepsilon - \frac{1}{2}u_{1x}\varepsilon - \frac{1}{2}qu_1 - \frac{1}{2}u_2 + \frac{1}{2}r\varepsilon + \frac{1}{2}u_2\varepsilon$$

$$+ \frac{1}{16}q^2\varepsilon + \frac{1}{2}qu_1\varepsilon - \alpha\beta - u_1^2 + u_1^2\varepsilon + \beta\beta_x,$$

$$\rho_3 = \alpha_{xx} + \frac{1}{4}q_x\alpha + \frac{1}{2}q\alpha_x + \frac{1}{2}\alpha r + \frac{1}{2}\beta r_x$$

$$\begin{aligned}
& + \frac{1}{2}\beta qr - r\beta_x + \frac{1}{16}q^2\alpha, \\
\delta_3 &= \beta_{xx} - \frac{1}{4}q_x\beta - \frac{1}{2}q\beta_x + \frac{1}{2}\beta r + \frac{1}{16}q^2\beta - \alpha_x.
\end{aligned} \tag{13}$$

$$\begin{pmatrix} 4A_{m+1} + 2E_{m+1} \\ 2C_{m+1} + G_{m+1} \\ 2A_{m+1} + 2E_{m+1} \\ C_{m+1} + G_{m+1} \\ 2\delta_{m+1} \\ -2\rho_{m+1} \end{pmatrix} = L \begin{pmatrix} 4A_m + 2E_m \\ 2C_m + G_m \\ 2A_m + 2E_m \\ C_m + G_m \\ 2\delta_m \\ -2\rho_m \end{pmatrix}, \tag{14}$$

Equations (12) can be written as

where

$$L = \begin{pmatrix} -\frac{1}{2}\partial - \frac{1}{4}\partial^{-1}q\partial & r + \partial^{-1}r\partial & -\partial^{-1}u_1\partial & u_2 + \partial^{-1}u_2\partial & \frac{1}{2}\partial + \frac{1}{2}\partial^{-1}\alpha\partial & \frac{1}{2}\beta + \partial^{-1}\beta\partial \\ \frac{1}{2} & \frac{1}{2}\partial - \frac{1}{4} & 0 & -u_1 & -\frac{1}{2}\beta & 0 \\ 0 & 0 & L_{33} & L_{34} & 0 & 0 \\ 0 & 0 & \frac{1}{2} & \frac{1}{2}\partial - u_1 - \frac{1}{4}q & 0 & 0 \\ -\beta & 2\alpha & \beta & -2\alpha & \partial - \frac{1}{4}q & -1 \\ \alpha + \beta\partial & -2\beta r & -\alpha - \beta\partial & 2\beta r & -\alpha\beta - r & -\partial - \frac{1}{4}q \end{pmatrix}, \tag{15}$$

$$L_{33} = -\frac{1}{2}\partial - \frac{1}{4}\partial^{-1}q\partial - \partial^{-1}u_1\partial, \quad L_{34} = r + u_2 + \partial^{-1}r\partial + \partial^{-1}u_2\partial.$$

Then, let us consider the spectral problem (8) with the following auxiliary problem:

which gives a nonlinear Lax super integrable hierarchy

$$\varphi_{t_n} = V^{(n)}\varphi \tag{16}$$

with

$$V^{(n)} = \sum_{j=0}^n \begin{pmatrix} A_j & B_j & E_j & F_j & \rho_j \\ C_j & -A_j & G_j & -E_j & \delta_j \\ 0 & 0 & A_j + E_j & B_j + F_j & 0 \\ 0 & 0 & C_j + G_j & -A_j - E_j & 0 \\ \delta_j & -\rho_j & -\delta_j & \rho_j & 0 \end{pmatrix} \lambda^{n-j} + C_{n+1}e_1(0) + G_{n+1}e_4(0). \tag{17}$$

$$u_{t_n} = \begin{pmatrix} q \\ r \\ u_1 \\ u_2 \\ \alpha \\ \beta \end{pmatrix}_{t_n} = \begin{pmatrix} -4C_{n+1,x} \\ 2B_{n+1} + 2rC_{n+1} \\ -G_{n+1,x} \\ 2F_{n+1} + 2u_2C_{n+1} + 2rG_{n+1} + 2u_2G_{n+1} \\ -\rho_{n+1} - \alpha C_{n+1} \\ \delta_{n+1} + \beta C_{n+1} \end{pmatrix}. \tag{19}$$

The super integrable hierarchy (19) is a nonlinear super integrable coupling for the super classical-Boussinesq hierarchy

From the compatible condition $\varphi_{x,t_n} = \varphi_{t_n,x}$ according to (8) and (17), we get the zero equation:

$$U_{t_n} - V_x^{(n)} + [U, V^{(n)}] = 0, \tag{18}$$

$$\bar{u}_{t_n} = \begin{pmatrix} q \\ r \\ \alpha \\ \beta \end{pmatrix}_{t_n} = \begin{pmatrix} -4C_{n+1,x} \\ 2B_{n+1} + 2rC_{n+1} \\ -\rho_{n+1} - \alpha C_{n+1} \\ \delta_{n+1} + \beta C_{n+1} \end{pmatrix}. \tag{20}$$

4. Super Hamiltonian Structures

A direct calculation reads

$$\begin{aligned}\text{Str}(U_\lambda, V) &= -4A - 2E, & \text{Str}(U_q, V) &= -A - \frac{1}{2}E, \\ \text{Str}(U_r, V) &= 2C + G, & \text{Str}(U_{u_1}, V) &= -2A - 2E, \\ \text{Str}(U_{u_2}, V) &= C + G, & \text{Str}(U_\alpha, V) &= 2\delta, \\ \text{Str}(U_\beta, V) &= -2\rho.\end{aligned}\quad (21)$$

Substituting the above results into the super trace identity in [6, 7]

$$\frac{\delta}{\delta u} \int \text{Str} \left(\frac{\delta U}{\delta \lambda} V \right) dx = \lambda^{-\gamma} \frac{\partial}{\partial \lambda} \lambda^\gamma \text{Str} \left(\frac{\delta U}{\delta u} V \right) \quad (22)$$

yields that

$$\frac{\delta}{\delta u} \int (-4A - 2E) dx = \lambda^{-\gamma} \frac{\partial}{\partial \lambda} \lambda^\gamma \begin{pmatrix} -A - \frac{1}{2}E \\ 2C + G \\ -2A - 2E \\ C + G \\ 2\delta \\ -2\rho \end{pmatrix}. \quad (23)$$

Comparing the coefficients of λ^{-n-1} on both sides of (23),

$$\frac{\delta}{\delta u} \int (-4A_{n+1} - 2E_{n+1}) dx = \lambda^{-\gamma} \frac{\partial}{\partial \lambda} \lambda^\gamma \begin{pmatrix} -A_n - \frac{1}{2}E_n \\ 2C_n + G_n \\ -2A_n - 2E_n \\ C_n + G_n \\ 2\delta_n \\ -2\rho_n \end{pmatrix}, \quad n \geq 0. \quad (24)$$

From the initial values in (12), we obtain $\gamma = 0$. Thus, we have

$$\begin{aligned}\frac{\delta H_n}{\delta u} &= \begin{pmatrix} -A_{n+1} - \frac{1}{2}E_{n+1} \\ 2C_{n+1} + G_{n+1} \\ -2A_{n+1} - 2E_{n+1} \\ C_{n+1} + G_{n+1} \\ 2\delta_{n+1} \\ -2\rho_{n+1} \end{pmatrix}, \\ H_n &= \int \frac{4A_{n+2} + 2E_{n+2}}{n+1} dx, \\ n &\geq 0.\end{aligned}\quad (25)$$

It then follows that the nonlinear super integrable couplings (22) possess the following super Hamiltonian form:

$$u_{t_n} = K_n(u) = J \frac{\delta H_n}{\delta u}, \quad (26)$$

where

$$J = \begin{pmatrix} 0 & -4\partial & 0 & 4\partial & 0 & 0 \\ -4\partial & 0 & \partial & 0 & -\alpha & \beta \\ 0 & \partial & 0 & -2\partial & 0 & 0 \\ 4\partial & 0 & -2\partial & 0 & -\alpha & \beta \\ 0 & \alpha & 0 & -\alpha & 0 & \frac{1}{2} \\ 0 & -\beta & 0 & \beta & \frac{1}{2} & 0 \end{pmatrix} \quad (27)$$

is a super Hamiltonian operator and H_n ($n \geq 0$) are Hamiltonian functions.

5. Reductions

Taking $\alpha = \beta = 0$, the hierarchy (26) reduces to a nonlinear integrable coupling of the super classical-Boussinesq hierarchy.

When $n = 2$ in (26), we obtain the nonlinear super integrable couplings of the second-order super classical-Boussinesq equations

$$\begin{aligned}q_{t_2} &= -\frac{1}{2}q_{xx} + 2r_x - 4\alpha_x\beta - 4\alpha\beta_x + \frac{1}{2}qq_x + 4\beta\beta_{xx} + 4\beta_x^2, \\ r_{t_2} &= \frac{1}{2}r_{xx} + \frac{1}{2}q_xr + \frac{1}{2}qr_x + 2\alpha\alpha_x - 2r\beta\beta_x, \\ u_{1,t_2} &= \frac{1}{2}(1-\varepsilon) \left[u_{2x} + q_xu_1 + qu_{1x} + 2u_1u_{1x} - u_{1xx} \right. \\ &\quad \left. + r_x - \frac{1}{4}q_{xx} + \frac{1}{4}qq_x \right] - \frac{1}{2}r_x + \frac{1}{8}q_{xx} \\ &\quad - \frac{1}{8}qq_x + \alpha_x\beta + \alpha\beta_x - \beta\beta_{xx}, \\ u_{t_2} &= (1-\varepsilon) \left[2u_{1x}r + 2u_{1x}u_2 + 2u_1u_{2x} + \frac{1}{2}u_{2xx} \right. \\ &\quad \left. + \frac{1}{2}q_xu_2 + \frac{1}{2}qu_{2x} + \frac{1}{2}qr_x \right. \\ &\quad \left. + \frac{1}{2}q_xr + \frac{1}{2}r_{xx} + 2r_xu_1 \right] \\ &\quad + 2r\beta\beta_x - \frac{1}{2}q_xr - \frac{1}{2}qr_x - \frac{1}{2}r_{xx} - 2\alpha\alpha_x\alpha_{t_2} \\ &= -\alpha_{xx} - \frac{1}{2}q\alpha_x - \frac{3}{8}q_x\alpha - \frac{1}{2}\beta r_x + \alpha\beta\beta_x, \\ \beta_{t_2} &= \beta_{xx} - \frac{1}{8}q_x\beta - \frac{1}{2}q\beta_x - \alpha_x.\end{aligned}\quad (28)$$

Taking $\alpha = \beta = 0$ in (28), we obtain the nonlinear super integrable couplings of the second-order super classical-Boussinesq equation

$$\begin{aligned}
 q_{t_2} &= -\frac{1}{2}q_{xx} + 2r_x + \frac{1}{2}qq_x, & r_{t_2} &= \frac{1}{2}r_{xx} + \frac{1}{2}q_x r + \frac{1}{2}qr_x, \\
 u_{1,t_2} &= \frac{1}{2}(1-\varepsilon) \left[u_{2x} + q_x u_1 + qu_{1x} + 2u_1 u_{1x} \right. \\
 &\quad \left. - u_{1xx} + r_x - \frac{1}{4}q_{xx} + \frac{1}{4}qq_x \right] \\
 &\quad - \frac{1}{2}r_x + \frac{1}{8}q_{xx} - \frac{1}{8}qq_x, \\
 u_{t_2} &= (1-\varepsilon) \left[2u_{1x}r + 2u_{1x}u_2 + 2u_1 u_{2x} + \frac{1}{2}u_{2xx} + \frac{1}{2}q_x u_2 \right. \\
 &\quad \left. + \frac{1}{2}qu_{2x} + \frac{1}{2}qr_x + \frac{1}{2}q_x r + \frac{1}{2}r_{xx} + 2r_x u_1 \right] \\
 &\quad - \frac{1}{2}q_x r - \frac{1}{2}qr_x - \frac{1}{2}r_{xx}.
 \end{aligned} \tag{29}$$

Letting $\varepsilon = 0$ in (28), we have

$$\begin{aligned}
 q_{t_2} &= -\frac{1}{2}q_{xx} + 2r_x - 4\alpha_x \beta - 4\alpha \beta_x + \frac{1}{2}qq_x + 4\beta \beta_{xx} + 4\beta_x^2, \\
 r_{t_2} &= \frac{1}{2}r_{xx} + \frac{1}{2}q_x r + \frac{1}{2}qr_x + 2\alpha \alpha_x - 2r\beta \beta_x, \\
 u_{1,t_2} &= \frac{1}{2} [u_{2x} + q_x u_1 + qu_{1x} + 2u_1 u_{1x} - u_{1xx}] \\
 &\quad + \alpha_x \beta + \alpha \beta_x - \beta \beta_{xx}, \\
 u_{t_2} &= \left[2u_{1x}r + 2u_{1x}u_2 + 2u_1 u_{2x} + \frac{1}{2}u_{2xx} \right. \\
 &\quad \left. + \frac{1}{2}q_x u_2 + \frac{1}{2}qu_{2x} + 2r_x u_1 \right] + 2r\beta \beta_x - 2\alpha \alpha_x, \\
 \alpha_{t_2} &= -\alpha_{xx} - \frac{1}{2}q\alpha_x - \frac{3}{8}q_x \alpha - \frac{1}{2}\beta r_x + \alpha \beta \beta_x, \\
 \beta_{t_2} &= \beta_{xx} - \frac{1}{8}q_x \beta - \frac{1}{2}q\beta_x - \alpha_x.
 \end{aligned} \tag{30}$$

When setting $\varepsilon = 1$, $u_1 = -(1/4)q$, $u_2 = r$ in (28), we obtain the second-order super classical-Boussinesq equations

$$\begin{aligned}
 q_{t_2} &= -\frac{1}{2}q_{xx} + 2r_x - 4\alpha_x \beta - 4\alpha \beta_x + \frac{1}{2}qq_x + 4\beta \beta_{xx} + 4\beta_x^2, \\
 r_{t_2} &= \frac{1}{2}r_{xx} + \frac{1}{2}q_x r + \frac{1}{2}qr_x + 2\alpha \alpha_x - 2r\beta \beta_x, \\
 \alpha_{t_2} &= -\alpha_{xx} - \frac{1}{2}q\alpha_x - \frac{3}{8}q_x \alpha - \frac{1}{2}\beta r_x + \alpha \beta \beta_x, \\
 \beta_{t_2} &= \beta_{xx} - \frac{1}{8}q_x \beta - \frac{1}{2}q\beta_x - \alpha_x.
 \end{aligned} \tag{31}$$

Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

References

- [1] Y. S. Li and L. N. Zhang, "A note on the super AKNS equations," *Journal of Physics A: Mathematical and General*, vol. 21, no. 7, pp. 1549–1552, 1988.
- [2] M. H. Tu and J. C. Shaw, "Hamiltonian structures of generalized Manin-Radul super-KdV and constrained super KP hierarchies," *Journal of Mathematical Physics*, vol. 40, no. 6, pp. 3021–3034, 1999.
- [3] S. X. Tao and T. C. Xia, "Lie algebra and Lie super algebra for integrable couplings of C-KdV hierarchy," *Chinese Physics Letters*, vol. 27, no. 4, Article ID 040202, 2010.
- [4] F. J. Yu, "Conservation laws and self-consistent sources for a super-classical-Boussinesq hierarchy," *Chinese Physics Letters*, vol. 28, no. 12, Article ID 120201, 2011.
- [5] H. Aratyn, E. Nissimov, and S. Pacheva, "Supersymmetric Kadomtsev-Petviashvili hierarchy: 'ghost' symmetry structure, reductions, and Darboux-Bäcklund solutions," *Journal of Mathematical Physics*, vol. 40, no. 6, pp. 2922–2932, 1999.
- [6] X. B. Hu, "An approach to generate superextensions of integrable systems," *Journal of Physics A: Mathematical and General*, vol. 30, no. 2, pp. 619–632, 1997.
- [7] W. X. Ma, J. S. He, and Z. Y. Qin, "A supertrace identity and its applications to superintegrable systems," *Journal of Mathematical Physics*, vol. 49, no. 3, Article ID 033511, 2008.
- [8] J. S. He, J. Yu, R. G. Zhou, and Y. Cheng, "Binary nonlinearization of the super AKNS system," *Modern Physics Letters B*, vol. 22, no. 4, pp. 275–288, 2008.
- [9] Q. P. Liu, Z. Popowicz, and K. Tian, "Supersymmetric reciprocal transformation and its applications," *Journal of Mathematical Physics*, vol. 51, no. 9, Article ID 093511, 2010.
- [10] W.-X. Ma, "Enlarging spectral problems to construct integrable couplings of soliton equations," *Physics Letters A*, vol. 316, no. 1-2, pp. 72–76, 2003.
- [11] W. X. Ma, X. X. Xu, and Y. F. Zhang, "Semidirect sums of Lie algebras and discrete integrable couplings," *Journal of Mathematical Physics*, vol. 47, no. 5, Article ID 053501, 2006.
- [12] F. K. Guo and Y. Zhang, "The quadratic-form identity for constructing the Hamiltonian structure of integrable systems," *Journal of Physics A: Mathematical and General*, vol. 38, no. 40, pp. 8537–8548, 2005.
- [13] W. X. Ma and M. Chen, "Hamiltonian and quasi-Hamiltonian structures associated with semi-direct sums of Lie algebras," *Journal of Physics A: Mathematical and General*, vol. 39, no. 34, pp. 10787–10801, 2006.
- [14] T. C. Xia and F. C. You, "Multi-component Dirac equations hierarchy and its multi-component integrable couplings system," *Chinese Physics*, vol. 16, no. 3, article 605, 2007.
- [15] Y. F. Zhang and E. G. Fan, "Coupling integrable couplings and bi-Hamiltonian structure associated with the Boiti-Pempinelli-Tu hierarchy," *Journal of Mathematical Physics*, vol. 51, no. 8, Article ID 083506, 2010.
- [16] Y. F. Zhang and H. Q. Zhang, "A direct method for integrable couplings of TD hierarchy," *Journal of Mathematical Physics*, vol. 43, no. 1, pp. 466–472, 2002.

- [17] W. X. Ma, “Nonlinear continuous integrable Hamiltonian couplings,” *Applied Mathematics and Computation*, vol. 217, no. 17, pp. 7238–7244, 2011.
- [18] W. X. Ma and Z. N. Zhu, “Constructing nonlinear discrete integrable Hamiltonian couplings,” *Computers & Mathematics with Applications*, vol. 60, no. 9, pp. 2601–2608, 2010.