Research Article

On $(a, 1)$ -Vertex-Antimagic Edge Labeling of Regular Graphs

Martin Bača,¹ Andrea Semaničová-Feňovčíková,¹ Tao-Ming Wang,² and Guang-Hui Zhang³

¹Department of Applied Mathematics and Informatics, Technical University, Letná 9, 04200 Košice, Slovakia

²Department of Applied Mathematics, Tunghai University, Taichung 40704, Taiwan

³Department of Applied Mathematics, National Chung Hsing University, Taichung 402, Taiwan

Correspondence should be addressed to Tao-Ming Wang; wang@go.thu.edu.tw

Received 1 April 2015; Accepted 26 May 2015

Academic Editor: Heping Zhang

Copyright © 2015 Martin Bača et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

An (a, s) -vertex-antimagic edge labeling (or an (a, s) -VAE labeling, for short) of G is a bijective mapping from the edge set $E(G)$ of a graph G to the set of integers $1, 2, \dots, |E(G)|$ with the property that the vertex-weights form an arithmetic sequence starting from a and having common difference s , where a and s are two positive integers, and the vertex-weight is the sum of the labels of all edges incident to the vertex. A graph is called (a, s) -antimagic if it admits an (a, s) -VAE labeling. In this paper, we investigate the existence of $(a, 1)$ -VAE labeling for disconnected 3-regular graphs. Also, we define and study a new concept (a, s) -vertex-antimagic edge deficiency, as an extension of (a, s) -VAE labeling, for measuring how close a graph is away from being an (a, s) -antimagic graph. Furthermore, the $(a, 1)$ -VAE deficiency of Hamiltonian regular graphs of even degree is completely determined. More open problems are mentioned in the concluding remarks.

1. Background and Introduction

All graphs in this paper are finite simple, undirected, and possibly disconnected, but without any isolated vertex or any isolated edge. For graph theoretic notations, we follow [1]. Over the past few decades, many kinds of graph labelings have been studied intensively, and an excellent survey of graph labeling can be found in Gallian's paper [2].

Hartsfield and Ringel in [3] introduced the concept of an *antimagic labeling*. In their terminology, a (p, q) -graph G with p vertices and q edges is called *antimagic* if its edges are labeled with labels $1, 2, \dots, q$ in such a way that all *vertex-weights* are pairwise distinct, where a vertex-weight of vertex v is the sum of labels of all edges incident with v . Hartsfield and Ringel [3] pointed out that antimagic graphs include paths P_n , $n \geq 3$, cycles, wheels, and complete graphs K_n , $n \geq 3$. They conjecture that every connected graph, except K_2 , is antimagic. Alon et al. [4] used several probabilistic tools and some techniques from analytic number theory to show that this conjecture is true for all graphs having minimum degree $\Omega(\log|V(G)|)$.

In 1993, Bodendiek and Walther [5] investigated antimagic labelings with certain restriction placed on the

vertex-weights. They defined the concept of an (a, s) -vertex-antimagic edge labeling as follows.

Definition 1. An (a, s) -vertex-antimagic edge labeling (or an (a, s) -VAE labeling for short) of a (p, q) -graph G is a bijective mapping f from the edge set $E(G)$ of a graph G to the set of integers $1, 2, \dots, q$ with the property that the vertex-weights form an arithmetic sequence starting from a and having common difference s , where a and s are two positive integers. The vertex-weight $wt_f(u)$ of the vertex u is the sum of the labels of all edges incident with the vertex u under the mapping f . A graph is called (a, s) -antimagic if it admits an (a, s) -VAE labeling.

Bodendiek and Walther in [6, 7] proved that the Herschel graph is not (a, s) -VAE and obtained both positive and negative results about (a, s) -VAE labelings for various cases of graphs called parachutes $P_{\alpha, \beta}$. They investigated (a, s) -VAE labelings for paths, cycles, and complete graphs in [8]. Characterization of all (a, s) -antimagic graphs of the prism $C_n \square P_2$ when n is even is given in [9]. In [10, 11] the (a, s) -VAE labelings for antiprisms have been investigated. It is proved in [12]

that the generalized Petersen graph $P(n, m)$ has an $(a, 1)$ -VAE labeling if and only if n is even, $n \geq 4$, $1 \leq m \leq n/2 - 1$, and $a = (7n + 4)/2$. Nicholas et al. [13] obtained several results about (a, d) -VAE labelings for caterpillars, unicyclic graphs, and complete bipartite graphs.

On the other hand, in 2002, MacDougall et al. [14] introduced the concept of *vertex magic total labeling* as follows. If G is a finite simple undirected graph with p vertices and q edges, then a vertex magic total labeling is a bijection f from $V(G) \cup E(G)$ to the integers $1, 2, \dots, p + q$ with the property that, for every $u \in V(G)$, the sum $f(u) + \sum_{uv \in E(G)} f(uv)$ is a constant. Note that, for regular graphs, the vertex magic total labeling is equivalent to the $(a, 1)$ -VAE labeling, while the vertices (edges) are assigned the smallest labels. More recently, Wang and Zhang [15] verified the existence of $(a, 1)$ -VAE labeling for particular classes of 3-regular graph H , where H contains a 1-factor and a 2-factor which consists of two 2-regular subgraphs with equal size. These results generalize and contain previous known examples such as Generalized Petersen Graphs.

The following theorem was proved in [16] by Ivančo and Semaničová, which guarantees the existence of the $(a, 1)$ -VAE-ness by adding an arbitrary even factor to an arbitrary $(a, 1)$ -antimagic graph.

Theorem 2 (see [16]). *Let the graph G admit an $(a, 1)$ -VAE labeling, and let H be any 2-factor over $V(G)$. Then, $G \cup H$ still admits an $(b, 1)$ -VAE labeling for some b .*

Therefore in order to study the $(a, 1)$ -antimagicness of a general regular graph, based upon Theorem 2 and the fact, pointed out by Petersen [17], that any regular graph of even degree has a 2-factorization, it is natural to explore the $(a, 1)$ -antimagicness of 2-regular graphs and 3-regular graphs, respectively. In this paper, we in particular investigate the existence of $(a, 1)$ -VAE labeling for disconnected 3-regular graphs and also define a new concept $(a, 1)$ -VAE deficiency, as an extension of $(a, 1)$ -VAE labeling, for studying those (regular) graphs not admitting an $(a, 1)$ -VAE labeling. The $(a, 1)$ -VAE deficiency is a parameter to measure how close a graph is from being $(a, 1)$ -antimagic. We notice that the method employed here is also valid for those graphs with multiple edges and loops. More examples and open problems will be provided in Section 5.

2. Preliminary Results

Suppose G is a (p, q) -graph with p vertices and q edges. The following are necessary conditions for graphs to admit an (a, s) -VAE labeling and $(a, 1)$ -VAE labeling in particular.

Lemma 3. *If a (p, q) -graph is (a, s) -antimagic, then $q(q+1) = pa + s((p-1)p/2)$.*

Proof. Consider that the total sum of all vertex-weights in (p, q) -graph is

$$2(1 + 2 + \dots + q) = a + (a + s) + \dots + (a + s(p-1)) \quad (1)$$

by two-way counting, and it implies that $q(q+1) = pa + s((p-1)p/2)$. $\square$

By Lemma 3 and for $s = 1$, in case $q = p$, then $a = (p + 3)/2$. Note that a is a positive integer, which implies that p is odd. Furthermore, in the case $q = p-1$, one has $a = (p-1)/2$, which implies that p is odd. Therefore, we have the following properties for $(a, 1)$ -VAE-ness.

Corollary 4. *Let G be a $(p, p-1)$ -graph. If G is $(a, 1)$ -antimagic, then p is odd.*

Corollary 5. *Trees of even order are not $(a, 1)$ -antimagic. In particular, a path of even order is not $(a, 1)$ -antimagic.*

Corollary 6. *Let G be a (p, p) -graph. If G is $(a, 1)$ -antimagic, then p is odd.*

Corollary 7. *The even cycle C_{2n} is not $(a, 1)$ -antimagic.*

Proposition 8. *Let G be an r -regular graph of order p and let G admit an $(a, 1)$ -VAE labeling. Then, we have the following:*

- (1) *If r is odd, then $p \equiv 0 \pmod{4}$.*
- (2) *If r is even, then $p \equiv 1 \pmod{2}$.*

Proof. Let G have q edges. Then, by hand-shaking lemma, we have that $pr = 2q$. Since G admits an $(a, 1)$ -VAE labeling, by Lemma 3, we have $q(q+1) = pa + (p-1)p/2$. Combining these equations, one has that $a = (1+rp/2)(r/2) - (p-1)/2 = (2r + r^2p - 2p + 2)/4$, which must be an integer. Therefore, it follows that if r is odd, then $p \equiv 0 \pmod{4}$, and if r is even, then $p \equiv 1 \pmod{2}$. $\square$

3. $(a, 1)$ -VAE for 3-Regular Graphs

In this section, we study the $(a, 1)$ -VAE labeling of disjoint union of 3-regular graphs. The main result is the following.

Theorem 9. *Let G be an $(a, 1)$ -antimagic 3-regular graph having a perfect matching. Then, the disjoint union of arbitrary number of copies of G , that is, mG , $m \geq 1$, is also a $(b, 1)$ -antimagic graph.*

Proof. Let G be a 3-regular graph with p vertices having a perfect matching.

Suppose that G admits an $(a, 1)$ -VAE labeling f such that

$$\begin{aligned} f : E(G) &\longrightarrow \left\{1, 2, \dots, \frac{3p}{2}\right\}, \\ \left\{wt_f(v) = \sum_{uv \in E(G)} f(uv) : v \in V(G)\right\} &= \{a, a+1, \dots, a+p-1\}. \end{aligned} \quad (2)$$

As G contains a perfect matching, we can divide the edges of G into two subsets $E_1(G)$ and $E_2(G)$, such that

$$\begin{aligned} E_1(G) \cup E_2(G) &= E(G), \\ E_1(G) \cap E_2(G) &= \emptyset, \end{aligned} \quad (3)$$

where the subset $E_1(G)$ consists of all edges belonging to the perfect matching.

For every vertex v in G , we denote by symbol v_i the corresponding vertex in the i th copy of G in mG . Analogously, let $e_i = u_i v_i$ denote the corresponding edge in the i th copy of G in mG .

Define the labeling f for the edges of mG in the following way:

$$g(e_i) = \begin{cases} m(f(e) - 1) + i, & \text{if } e \in E_2(G), i = 1, 2, \dots, m, \\ mf(e) + 1 - i, & \text{if } e \in E_1(G), i = 1, 2, \dots, m. \end{cases} \quad (4)$$

Let $t \in \{1, 2, \dots, 3p/2\}$. We consider two cases.

Case 1. If the number t is assigned by the labeling f to an edge from $E_2(G)$, then the corresponding edges in the copies in mG will receive labels under the labeling g :

$$\begin{array}{ccccccc} mt(t-1)+1, & m(t-1)+2, & \dots & m(t-1)+i, & \dots & mt. \\ \text{in } G_1 & \text{in } G_2 & \dots & \text{in } G_i & \dots & \text{in } G_m. \end{array} \quad (5)$$

Case 2. If the number t is assigned by the labeling f to an edge from $E_1(G)$, then the corresponding edges in the copies in mG will have the following labels under the labeling g :

$$\begin{array}{ccccccc} mt, & mt-1, & \dots & mt+1-i, & \dots & m(t-1)+1. \\ \text{in } G_1 & \text{in } G_2 & \dots & \text{in } G_i & \dots & \text{in } G_m. \end{array} \quad (6)$$

It is easy to see that the edge labels in mG are not overlapping; thus, the labeling g is a bijective function which

assigns the integers $\{1, 2, \dots, 3mp/2\}$ to the edges of mG ; thus, g is an edge labeling.

As G is a 3-regular graph with a perfect matching, then every vertex in G is incident to exactly one edge from $E_1(G)$ and exactly two edges from $E_2(G)$. For a given vertex v , let $e_{v,1}^1, e_{v,2}^2, e_{v,3}^2$ be the three edges incident to the vertex v , where e^j means that the edge belongs to the subset $E_j(G)$, $j = 1, 2$.

For the vertex-weight of $v_i \in V(G_i)$, we have

$$\begin{aligned} wt_g(v_i) &= g(e_{v,1i}^1) + g(e_{v,2i}^2) + g(e_{v,3i}^2) \\ &= (mf(e_{v,1}^1) + 1 - i) + (m(f(e_{v,2}^2) - 1) + i) \\ &\quad + (m(f(e_{v,3}^2) - 1) + i) \\ &= m(f(e_{v,1}^1) + f(e_{v,2}^2) + f(e_{v,3}^2)) - 2m + 1 \\ &\quad + i = mwt_f(v) - 2m + 1 + i \\ &= m(wt_f(v) - 2) + 1 + i. \end{aligned} \quad (7)$$

As

$$\begin{aligned} \left\{ wt_f(v) = \sum_{uv \in E(G)} f(uv) : v \in V(G) \right\} \\ = \{a, a+1, \dots, a+p-1\}, \end{aligned} \quad (8)$$

we get that the vertex-weights under the labeling g in the components are

$$\begin{array}{ccccccc} G_1: & m(a-2)+2, & m(a-1)+2, & \dots & m(a+p-3)+2, \\ G_2: & m(a-2)+3, & m(a-1)+3, & \dots & m(a+p-3)+3, \\ & \vdots & \vdots & & \vdots \\ G_i: & m(a-2)+1+i, & m(a-1)+1+i, & \dots & m(a+p-3)+1+i, \\ & \vdots & \vdots & & \vdots \\ G_m: & m(a-1)+1, & ma+1, & \dots & m(a+p-2)+1. \end{array} \quad (9)$$

The reader can easily verify that the vertex-weights are distinct and consecutive:

$$\{wt_g(v) : v \in V(mG)\} = \{m(a-2)+2, m(a-2)+3, \dots, m(a+p-2)+1\}. \quad (10)$$

This means that mG has a $(m(a-2)+2, 1)$ -VAE labeling. $\square$

It is possible to extend the result from the previous theorem also for the disjoint union of arbitrary 3-regular graphs having a perfect matching that satisfy certain additional conditions.

We will follow the notation used in the proof of Theorem 9.

Theorem 10. Let G_i be an $(a, 1)$ -antimagic 3-regular graph of order p having a perfect matching, $i = 1, 2, \dots, m$. Let the set of all edge labels belonging to the perfect matching under the $(a, 1)$ -VAE labeling f_i of a graph G_i be the same for every i , $i = 1, 2, \dots, m$.

Then, the disjoint union $\bigcup_{i=1}^m G_i$ is also a $(b, 1)$ -antimagic graph.

Proof. Let G_i , $i = 1, 2, \dots, m$, be a 3-regular graph of order p having a perfect matching, $i = 1, 2, \dots, m$, and note that G_i is not necessarily isomorphic to G_j for $i \neq j$.

For each G_i , $i = 1, 2, \dots, m$, there exists an $(a, 1)$ -VAE labeling f_i such that the set of all edge labels belonging to the perfect matching under the $(a, 1)$ -VAE labeling f_i of a graph G_i is the same for every graph G_i . This means that

$$\begin{aligned} f_i : E_1(G_i) &\longrightarrow \{t_1, t_2, \dots, t_{p/2}\} \\ E_2(G_i) &\longrightarrow \frac{\{1, 2, \dots, 3p/2\}}{\{t_1, t_2, \dots, t_{p/2}\}}, \\ \left\{ wt_{f_i}(v) = \sum_{uv \in E(G_i)} f_i(uv) : v \in V(G_i) \right\} \\ &= \{a, a+1, \dots, a+p-1\}. \end{aligned} \quad (11)$$

We define the labeling f for the edges of $\bigcup_{i=1}^m G_i$ in the following way:

$$f(e) = \begin{cases} m(f_i(e) - 1) + i, & \text{if } e \in E_2(G_i), \\ mf_i(e) + 1 - i, & \text{if } e \in E_1(G_i). \end{cases} \quad (12)$$

Let $t \in \{1, 2, \dots, 3p/2\}$. We consider two cases.

Case 1. If the number t is assigned by the labeling f_i to an edge from $E_2(G_i)$, then the corresponding edge in $\bigcup_{i=1}^m G_i$ will receive the following label under the labeling f :

$$\begin{array}{ccccccc} m(t-1)+1, & m(t-1)+2, & \dots & m(t-1)+i, & \dots & mt. \\ \text{in } G_1 & \text{in } G_2 & \dots & \text{in } G_i & \dots & \text{in } G_m. \end{array} \quad (13)$$

Case 2. If the number t is assigned by the labeling f_i to an edge from $E_1(G_i)$, then the corresponding edge in $\bigcup_{i=1}^m G_i$ will receive this label under the labeling f :

$$\begin{array}{ccccccc} mt, & mt-1, & \dots & mt+1-i, & \dots & m(t-1)+1. \\ \text{in } G_1 & \text{in } G_2 & \dots & \text{in } G_i & \dots & \text{in } G_m. \end{array} \quad (14)$$

It is easy to see that the edge labels in $\bigcup_{i=1}^m G_i$ are not overlapping; thus, the labeling f is a bijective function which assigns the integer $\{1, 2, \dots, 3mp/2\}$ to the edges of $\bigcup_{i=1}^m G_i$; thus, f is an edge labeling.

Moreover, analogously as in the proof of Theorem 9, we get that the vertex-weights form an arithmetic sequence with a difference 1. This produces the desired result. $\square$

4. $(a, 1)$ -VAE Deficiency of Even Regular Graphs

We start this section by defining a new concept (a, s) -vertex-antimagic edge deficiency for a (p, q) -graph G as follows. It is a parameter to study furthermore those graphs which do not admit any (a, s) -VAE labeling and to measure how close they are away from being an (a, s) -antimagic graph.

Definition 11. The (a, s) -vertex-antimagic edge deficiency (or (a, s) -VAE deficiency for short) is defined as the min k such that the edge labeling $f : E(G) \rightarrow \{1, 2, \dots, q+k\}$ is (a, s) -VAE. The (a, s) -VAE deficiency of the graph G is denoted by

$d_s(G)$. Note that $d_s(G) = 0$ if G is (a, s) -antimagic and $d_s(G) = +\infty$ if no such k exists for the graph G to be (a, s) -antimagic.

In the following, we determine completely the $(a, 1)$ -VAE deficiency of paths and cycles. Also, as a corollary, the $(a, 1)$ -VAE deficiency of Hamiltonian regular graphs of even degree is obtained. First, we have the following two lemmas giving $(a, 1)$ -VAE labelings for odd cycles and paths; see also [8].

Lemma 12. The cycle C_{2n+1} is $(a, 1)$ -antimagic for $n \geq 1$.

Proof. Given a notation for C_{2n+1} with $V(C_{2n+1}) = \{v_i : i = 1, 2, \dots, 2n+1\}$ and $E(C_{2n+1}) = \{v_i v_{i+1} : i = 1, 2, \dots, 2n\} \cup \{v_1 v_{2n+1}\}$, we give an edge labeling f for $E(C_{2n+1})$ with

$$\begin{aligned} f(v_1 v_{2n+1}) &= n+1, \\ f(v_i v_{i+1}) &= \begin{cases} \frac{i+1}{2}, & \text{if } i = 1, 3, \dots, 2n-1, \\ n+1 + \frac{i}{2}, & \text{if } i = 2, 4, \dots, 2n \end{cases} \end{aligned} \quad (15)$$

and it implies the vertex-weight at v_i as follows:

$$wt_f(v_i) = n+1+i, \quad \text{for } i = 1, 2, \dots, 2n+1. \quad (16)$$

Hence, C_{2n+1} is $(n+2, 1)$ -antimagic. $\square$

Lemma 13. The path P_{2n+1} is $(a, 1)$ -antimagic for $n \geq 1$.

Proof. Given a notation for P_{2n+1} with $V(P_{2n+1}) = \{v_i : i = 1, 2, \dots, 2n+1\}$ and $E(P_{2n+1}) = \{v_i v_{i+1} : i = 1, 2, \dots, 2n\}$, we give an edge labeling f for $E(P_{2n+1})$ such that

$$f(v_i v_{i+1}) = \begin{cases} \frac{i}{2}, & \text{if } i = 2, 4, \dots, 2n, \\ n + \frac{i+1}{2}, & \text{if } i = 1, 3, \dots, 2n-1 \end{cases} \quad (17)$$

and it implies the vertex-weight at v_i as follows:

$$wt_f(v_i) = \begin{cases} n+i, & \text{if } i = 1, 2, \dots, 2n, \\ n, & \text{if } i = 2n+1. \end{cases} \quad (18)$$

Hence, P_{2n+1} is $(n, 1)$ -antimagic. $\square$

Here, we have a general observation that every graph G of order p , where $p \equiv 2 \pmod{4}$, can not be made $(a, 1)$ -antimagic.

Lemma 14. Let G be a graph of order p , where $p \equiv 2 \pmod{4}$. Then, $d_1(G) = +\infty$.

Proof. Let G have q edges and $p = 4k+2$ vertices. Assign labels $e_1, e_2, \dots, e_q$ to edges of graph G and suppose the existence of an $(a, 1)$ -VAE labeling with the associated vertex-weights $a, a+1, \dots, a+(4k+1)$. Consider the sum of all vertex-weights:

$$\begin{aligned} a + (a+1) + \dots + a + (4k+1) \\ = (4k+2)a + (4k+1)(2k+1) \end{aligned} \quad (19)$$

which is also equal to $2(e_1 + e_2 + \dots + e_q)$. Note that $2(e_1 + e_2 + \dots + e_q)$ and $(4k + 2)a$ are both even, but $(4k + 1)(2k + 1)$ is odd, a contradiction. $\square$

In the following lemmas, we are dealing with the $(a, 1)$ -VAE deficiency of cycles and paths.

Lemma 15. Consider $d_1(C_{4n}) = 1$ for $n \geq 1$.

Proof. First, we find the missing value x in the set of edge labels $\{1, 2, \dots, 4n\}$. Note that

$$\begin{aligned} 2(1 + 2 + \dots + (4n + 1) - x) \\ = a + (a + 1) + \dots + (a + 4n - 1) \end{aligned} \quad (20)$$

and it implies $a = (4n^2 + 7n + 1 - x)/2n = 2n + 3 + (n + 1 - x)/2n \in \mathbb{Z}$ and thus $(n + 1 - x)/2n \in \mathbb{Z}$. Suppose that $(n + 1 - x)/2n = t$, and then one has $-n \leq -2nt \leq 3n - 1$ since $1 \leq x \leq 4n$. Therefore, t must be 0 or -1 , and it implies that the missing value x must be $n + 1$ or $3n + 1$. We show that for both missing values there exist $(a, 1)$ -VAE labelings.

Let the vertex set and the edge set of C_{4n} be $V(C_{4n}) = \{v_i : i = 1, 2, \dots, 4n\}$ and $E(C_{4n}) = \{v_i v_{i+1} : i = 1, 2, \dots, 4n - 1\} \cup \{v_1 v_{4n}\}$.

Case 1. If the missing value is $n + 1$, then the $(a, 1)$ -VAE labeling can be defined as follows:

$$\begin{aligned} f_1(v_1 v_{4n}) &= 4n + 1, \\ f_1(v_i v_{i+1}) &= \begin{cases} \frac{i+1}{2}, & \text{if } i = 1, 3, \dots, 2n-1, \\ \frac{i+3}{2}, & \text{if } i = 2n+1, 2n+3, \dots, 4n-1, \\ 2n+1 + \frac{i}{2}, & \text{if } i = 2, 4, \dots, 4n-2. \end{cases} \end{aligned} \quad (21)$$

Case 2. If the missing value is $3n + 1$, then, for example, consider the following edge labeling:

$$\begin{aligned} f_2(v_1 v_{4n}) &= 4n + 1, \\ f_2(v_i v_{i+1}) &= \begin{cases} \frac{i+1}{2}, & \text{if } i = 1, 3, \dots, 4n-1, \\ 2n + \frac{i}{2}, & \text{if } i = 2, 4, \dots, 2n, \\ 2n+1 + \frac{i}{2}, & \text{if } i = 2n+2, 2n+4, \dots, 4n-2. \end{cases} \end{aligned} \quad (22)$$

Then, we find that the vertex-weights form the set $\{2n+3, 2n+4, \dots, 6n+2\}$ for Case 1 and the set $\{2n+2, 2n+3, \dots, 6n+1\}$ for Case 2. Hence, $d_1(C_{4n}) = 1$ as required. $\square$

Lemma 16. Consider $d_1(P_{4n}) = 1$ for $n \geq 1$.

Proof. First, we suppose that $d_1(P_{4n}) \leq 1$ and we want to find the missing value x in the set of edge labels $\{1, 2, \dots, 4n - 1\}$. Then,

$$\begin{aligned} 2(1 + 2 + \dots + 4n - x) \\ = a + (a + 1) + \dots + (a + 4n - 1) \end{aligned} \quad (23)$$

and it implies that $a = (4n^2 + 3n - x)/2n = 2n + 1 + (n - x)/2n$ and thus $(n - x)/2n \in \mathbb{Z}$. Suppose that $(n - x)/2n = t$, and then $1 - n \leq -2nt \leq 3n - 1$, since $1 \leq x \leq 4n - 1$. Therefore, t must be 0 or -1 and it implies that the missing value is $x = n$ or $3n$. As in the proof of the previous lemma we show that for both cases it is possible to find required $(a, 1)$ -VAE labelings.

We denote the vertices and the edges of P_{4n} such that $V(P_{4n}) = \{v_i : i = 1, 2, \dots, 4n\}$ and $E(P_{4n}) = \{v_i v_{i+1} : i = 1, 2, \dots, 4n - 1\}$.

Case 1. If the missing value is n , then we define the edge labeling f_3 in the following way:

$$\begin{aligned} f_3(v_i v_{i+1}) &= \begin{cases} 2n + \frac{i+1}{2}, & \text{if } i = 1, 3, \dots, 4n-1, \\ \frac{i}{2}, & \text{if } i = 2, 4, \dots, 2n-2, \\ 1 + \frac{i}{2}, & \text{if } i = 2n, 2n+2, \dots, 4n-2. \end{cases} \end{aligned} \quad (24)$$

Case 2. If the missing value is $3n$, then we define the labeling f_4 such that

$$\begin{aligned} f_4(v_i v_{i+1}) &= \begin{cases} 2n + \frac{i-1}{2}, & \text{if } i = 1, 3, \dots, 2n-1, \\ 2n + \frac{i+1}{2}, & \text{if } i = 2n+1, 2n+3, \dots, 4n-1, \\ \frac{i}{2}, & \text{if } i = 2, 4, \dots, 4n-2. \end{cases} \end{aligned} \quad (25)$$

Then, we obtain that the vertex-weights attain the values from the sets $\{2n+1, 2n+2, \dots, 6n\}$ for Case 1 and $\{2n, 2n+3, \dots, 6n-1\}$ for Case 2. Thus, $d_1(P_{4n}) = 1$. $\square$

To summarize, we have the following $(a, 1)$ -VAE deficiency for paths P_m and cycles C_m .

Theorem 17. Let $m \geq 2$. Then,

$$d_1(P_m) = \begin{cases} 0, & \text{if } m \equiv 1, 3 \pmod{4}, \\ 1, & \text{if } m \equiv 0 \pmod{4}, \\ +\infty, & \text{if } m \equiv 2 \pmod{4}. \end{cases} \quad (26)$$

Theorem 18. Let $m \geq 3$. Then,

$$d_1(C_m) = \begin{cases} 0, & \text{if } m \equiv 1, 3 \pmod{4}, \\ 1, & \text{if } m \equiv 0 \pmod{4}, \\ +\infty, & \text{if } m \equiv 2 \pmod{4}. \end{cases} \quad (27)$$

Therefore, as a corollary of Theorem 18, we immediately have the following result.

Theorem 19. *Let G be a $2r$ -regular, $r \geq 2$, Hamiltonian graph of order p . Then,*

$$d_1(G) = \begin{cases} 0, & \text{if } p \equiv 1, 3 \pmod{4}, \\ 1, & \text{if } p \equiv 0 \pmod{4}, \\ +\infty, & \text{if } p \equiv 2 \pmod{4}. \end{cases} \quad (28)$$

Proof. By Theorem 18, Lemma 12, Theorem 2, and also Petersen's 2-factorization of regular graphs of even degree, we get that a Hamiltonian even regular graph of odd order is $(a, 1)$ -antimagic. On the other hand, by Lemma 15 and Theorem 2, we get that the Hamiltonian even regular graphs of order $4n$ have the $(a, 1)$ -VAE deficiency equal to 1. Finally, by Lemma 14, we complete the proof. $\square$

As a corollary, we have the following example of $(a, 1)$ -VAE deficiency for the Cartesian product of two cycles $C_m \square C_n$, $m, n \geq 3$. Note that $C_m \square C_n$ always contains a Hamiltonian cycle.

Corollary 20. *Let $m, n \geq 3$. Then,*

$$d_1(C_m \square C_n) = \begin{cases} 0, & \text{if } mn \equiv 1, 3 \pmod{4}, \\ 1, & \text{if } mn \equiv 0 \pmod{4}, \\ +\infty, & \text{if } mn \equiv 2 \pmod{4}. \end{cases} \quad (29)$$

Note that similarly one may have the formula for $(a, 1)$ -VAE deficiency of the higher dimensional toroidal grids, that is, the Cartesian product of t cycles $C_{m_1} \square C_{m_2} \square \cdots \square C_{m_t}$, $m_i \geq 3$, for each i .

5. Concluding Remarks and Further Studies

Notice that in 2009 Holden et al. [18] raised a conjecture for the existence of $(a, 1)$ -VAE labeling of 2-regular graphs as follows: a 2-regular graph of odd order possesses an $(a, 1)$ -VAE labeling if and only if it is not one of $C_4 \cup C_3$, $C_4 \cup 3C_3$, or $C_5 \cup 2C_3$. Note that the terminology of the labeling they made is the strong vertex magic total labeling, which is exactly equivalent to the $(a, 1)$ -VAE labeling. Therefore, it is natural to ask for the following.

Problem 1. Determine the $(a, 1)$ -VAE deficiency for $C_4 \cup C_3$, $C_4 \cup 3C_3$, and $C_5 \cup 2C_3$.

Moreover, as a generalization of the above result, we obtain in the last section the $(a, 1)$ -VAE deficiency of a Hamiltonian regular graph of even degree, and we are concerned with the following situation. Note that Swaminathan and Jeyanthi [19] pointed out the following: mC_n is $(a, 1)$ -antimagic if and only if m, n are odd. Therefore, it is natural to ask for the following.

Problem 2. Determine the $(a, 1)$ -VAE deficiency for the 2-regular graph mC_n .

If Problem 2 is answered, then the $(a, 1)$ -VAE deficiency for an arbitrary regular graph of even degree containing a 2-factor mC_n is answered. More generally, we have the following.

Problem 3. Determine the $(a, 1)$ -VAE deficiency for a general 2-regular graph.

If Problem 3 is answered, then the $(a, 1)$ -VAE deficiency for an arbitrary regular graph of even degree is answered. As for 3-regular graphs and general odd regular graphs, we ask for the following.

Problem 4. Determine the $(a, 1)$ -VAE deficiency for 3-regular Generalized Petersen Graphs $P(n, k)$ and Möbius Ladder Graphs M_n .

Problem 5. Determine the $(a, 1)$ -VAE deficiency for a general 3-regular graph.

Problem 6. Determine the $(a, 1)$ -VAE deficiency for a general odd regular graph.

It is not hard to check that K_4 does not admit any $(a, 1)$ -VAE labeling. However, with the aid of computer programs, we have found that tK_4 admits an $(a, 1)$ -VAE labeling for $2 \leq t \leq 9$. This leads to the following conjecture.

Conjecture 21. tK_4 , the disjoint union of t copies of K_4 , admits the $(a, 1)$ -VAE labeling for $t \geq 2$.

Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

Acknowledgment

Tao-Ming Wang's research is supported partially by the Grant no. MOST 103-2115-M-029-001 from the Ministry of Science and Technology of Taiwan.

References

- [1] G. Chartrand and L. Lesniak, *Graphs & Digraphs*, Chapman & Hall/CRC Press, 1996.
- [2] J. A. Gallian, "A dynamic survey of graph labeling," *The Electronic Journal of Combinatorics*, vol. 16, article DS6, 2013.
- [3] N. Hartsfield and G. Ringel, *Pearls in Graph Theory*, Academic Press, San Diego, Calif, US, 1990.
- [4] N. Alon, G. Kaplan, A. Lev, Y. Roditty, and R. Yuster, "Dense graphs are antimagic," *Journal of Graph Theory*, vol. 47, no. 4, pp. 297–309, 2004.
- [5] R. Bodendiek and G. Walther, "Arithmetisch antimagische Graphen," in *Graphentheorie III*, K. Wagner and R. Bodendiek, Eds., BI-Wiss, Mannheim, Germany, 1993.
- [6] R. Bodendiek and G. Walther, "On (a, d) -antimagic parachutes," *Ars Combinatoria*, vol. 42, pp. 129–149, 1996.
- [7] R. Bodendiek and G. Walther, " (a, d) -antimagic parachutes II," *Ars Combinatoria*, vol. 46, pp. 33–63, 1997.

- [8] R. Bodendiek and G. Walther, "On arithmetic antimagic edge labelings of graphs," *Mitteilungen der Mathematischen Gesellschaft in Hamburg*, vol. 17, pp. 85–99, 1998.
- [9] M. Bača and I. Holländer, "On $(a; d)$ -antimagic prisms," *Ars Combinatoria*, vol. 48, pp. 297–306, 1998.
- [10] M. Bača, "Antimagic labellings of antiprisms," *Journal of Combinatorial Mathematics and Combinatorial Computing*, vol. 35, pp. 217–224, 2000.
- [11] M. Miller, M. Bača, and Y. Lin, "On two conjectures concerning $(a; d)$ -antimagic labelings of antiprisms," *Journal of Combinatorial Mathematics and Combinatorial Computing*, vol. 37, pp. 251–254, 2001.
- [12] M. Bača, "Consecutive-magic labeling of generalized Petersen graphs," *Utilitas Mathematica*, vol. 58, pp. 237–241, 2000.
- [13] T. Nicholas, S. Somasundaram, and V. Vilfred, "On (a, d) -Antimagic special trees, unicyclic graphs and complete bipartite graphs," *Ars Combinatoria*, vol. 70, pp. 207–220, 2004.
- [14] J. A. MacDougall, M. Miller, Slamin, and W. D. Wallis, "Vertex magic total labelings of graphs," *Utilitas Mathematica*, vol. 61, pp. 3–21, 2002.
- [15] T.-M. Wang and G.-H. Zhang, "E-super vertex magic regular graphs of odd degree," *Electronic Notes in Discrete Mathematics*. In press.
- [16] J. Ivančo and A. Semaničová, "Some constructions of super-magic graphs using antimagic graphs," *SUT Journal of Mathematics*, vol. 42, no. 2, pp. 177–186, 2006.
- [17] J. Petersen, "Die Theorie der regulären graphs," *Acta Mathematica*, vol. 15, no. 1, pp. 193–220, 1891.
- [18] J. Holden, D. McQuillan, and J. M. McQuillan, "A conjecture on strong magic labelings of 2-regular graphs," *Discrete Mathematics*, vol. 309, no. 12, pp. 4130–4136, 2009.
- [19] V. Swaminathan and P. Jeyanthi, "Super vertex magic labeling," *Indian Journal of Pure and Applied Mathematics*, vol. 34, no. 6, pp. 935–939, 2003.