Research Article

¹H and ¹³C NMR Assignments of Cytotoxic 3*S*-1,2,3,4-Tetrahydro- β -carboline-3-carboxylic Acid from the Leaves of *Cichorium endivia*

Fu-Xin Wang, An-Jun Deng, Jin-Feng Wei, Hai-Lin Qin, and Ai-Ping Wang

Institute of Materia Medica, Chinese Academy of Medical Sciences and Peking Union Medical College, Beijing 100050, China

Correspondence should be addressed to Hai-Lin Qin, qinhailin@imm.ac.cn and Ai-Ping Wang, wangaping@imm.ac.cn

Received 17 September 2012; Accepted 26 November 2012

Academic Editor: Feng Wei

Copyright © 2012 Fu-Xin Wang et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

An amino acid, 3*S*-1,2,3,4-tetrahydro- β -carboline-3-carboxylic acid, was isolated for the first time from the leaves of *Cichorium endivia*. The complete assignment of its ¹H and ¹³C NMR spectroscopic data was carried out also for the first time based on extensive 1D and 2D NMR experiments. Cytotoxicity of this isolated compound against HCT-8 and HepG2 human cancer cell lines was evaluated for the first time, with moderate activities being found.

1. Introduction

Cichorium endivia L. is a popular vegetable from the family of Compositae and is widely cultivated and consumed all over the world. Its popularity is also attributed to the healthy properties mainly due to supply of antioxidant activity [1, 2]. However, phytochemical investigation on this plant is very rare up to now, to the best of our knowledge, only a few papers had reported a few compounds, including five ones by our group [3]. The ongoing research aims at confirming the bioactive compounds from this popular vegetable, and a prevailing and known amino acid, 3*S*-1,2,3,4-tetrahydro- β -carboline-3-carboxylic acid (**1**), was isolated for the first time. By way of the literature survey, it can be learned that the complete assignment of the NMR data of **1** was very deficient up to now, with no practical conducting being obtained due to the poor solubility of **1** in most prevalent solvents and causing the citation of literatures an obvious state of chaos [4, 5]. In this paper, we describe the complete assignment of the ¹H and ¹³C NMR spectroscopic data of **1** based on the determining of optimized solvent and extensive 1D and 2D NMR experiments. An investigation focusing on the cytotoxicity of compound **1** against HCT-8 and HepG2 human cancer cell lines showed that **1** inhibits the cells growth by a moderate reduction in viability of subjects.

2. Results and Discussion

Compound **1** was isolated as an amorphous pale-yellow powder (MeOH/H₂O). Its positive-ion ESI-MS spectrum showed the quasimolecular ion peaks at *m/z* 217.1 [M+H]⁺ and 239.1 [M+Na]⁺, and its molecular formula was established to be C₁₂H₁₂N₂O₂ by the quasimolecular ion peak in the positive mode HRESI-MS experiment at *m/z* 217.0967 [M+H]⁺. The IR spectrum showed strong absorptions at $\nu_{\max}$ 3284, 3019, 1642, 1598, 1452, 1409, and 740 cm⁻¹, indicating the presence of 1,2-disubstituted benzene moiety and labile hydrogen. It was preliminarily identified as 3*S*-1,2,3,4-tetrahydro- β -carboline-3-carboxylic acid by comparison of the ¹H NMR spectroscopic data obtained in DMSO-*d*₆ with the literature values [4, 5], but some obvious errors or inconsistency were evident, including the coupling constants and data ownership (Table 1). Whereas the obtainment of ¹³C NMR spectrum in the NMR solvent of DMSO-*d*₆ was very difficult due to the above-mentioned poor solubility, compound **1** was then recorded the 1D and 2D NMR spectra within D₂O + drops of F₃CCOOD, which proved to be a good solvent for **1**. The ¹H NMR spectroscopic data also clearly revealed the existence of 1,2-disubstituted benzene moiety, with four diagnostic signals from an aromatic ABCD spin system resonating at δ_{H} 7.13 (1H, t, *J* = 7.6 Hz, H-6),

TABLE 1: ^1H and ^{13}C NMR spectroscopic data for compound **1**.

Number	$\delta_{\text{H}}^{\text{ab}}$	$\delta_{\text{C}}^{\text{ac}}$	$\delta_{\text{H}}^{\text{de}}$	$\delta_{\text{H}}^{\text{d}}$ [4]	$\delta_{\text{H}}^{\text{d}}$ [5]	$\delta_{\text{C}}^{\text{d}}$ [5]
1a	4.38 d (15.6)	40.5 t	4.15 d (15.3)	—	—	40.3
1b	4.54 d (15.6)		4.23 d (15.9)	4.22 d (4.8)	—	
3	4.27 dd (10.4, 5.6)	55.0 d	3.60 m	3.14	—	55.3
4a	3.13 dd (10.8, 16.4)	21.7 t	2.81 dd-like	2.83 ddd (10.5, 5.0, 2.4)	—	18.0
4b	3.38 dd (5.6, 16.4)		3.13 br d-like	3.69 dd (10.5, 5.0)	—	
4a'		104.9 s				104.3
4b'		125.6 ^f s				128.5
5	7.53 d (8.0)	118.2 d	7.32 d (7.5)	7.33 d (8.0)	7.38 d (8.2)	118.5
6	7.13 t (7.6)	120.0 d	7.06 t (7.5)	7.08 t (8.0)	7.06 t (8.2)	117.5
7	7.22 t (8.0)	122.9 d	6.97 t (7.8)	6.99 t (7.5)	6.96 t (8.2)	121.1
8	7.43 d (8.0)	111.9 d	7.43 d (7.5)	7.44 d (7.5)	7.44 d (8.2)	111.8
8a		136.7 s				136.1
9a		125.4 ^f s				—
COOH		171.3 s	10.91 s	10.93 s	—	165.6
9-NH				—	10.66 s	

^a D₂O + drops of F₃CCOOD; ^b 400 MHz. ^c 100 MHz. ^d in DMSO-d₆; ^e 300 MHz. ^f Assignments may be interchanged.

7.22 (1H, t, J = 8.0 Hz, H-7), 7.43 (1H, d, J = 8.0 Hz, H-8), and 7.53 (1H, d, J = 8.0 Hz, H-5), which correlated to the aromatic carbon signals at δ_{C} 120.0 (C-6), 122.9 (C-7), 111.9 (C-8), and 118.2 (C-5), respectively, in the HSQC spectrum. In addition, five well-resolved and characteristic signals at δ_{H} 4.38 (1H, d, J = 15.6 Hz, H-1_a), 4.54 (1H, d, J = 15.6 Hz, H-1_b), 4.27 (1H, dd, J = 10.4, 5.6 Hz, H-3), 3.13 (1H, dd, J = 10.8, 16.4 Hz, H-4_a), and 3.38 (1H, dd, J = 5.6, 16.4 Hz, H-4_b) were also examined in the ^1H NMR spectrum, which were, conveniently according to their coupling constants and with the aid of ^1H , ^1H -COSY spectrum, assigned to one AB spin system from an isolated methylene group and one ABX system from one methylene and one methine correlated together by sp^3 hybridized bond. These three functional groups were obviously deshielded and were correlated to their corresponding carbon signals at δ_{C} 40.5 (C-1), 55.0 (C-3), and 21.7 (C-4), respectively, in the HSQC spectrum. Simply, these three groups were arranged to either sp^2 hybridized carbons or nitrogen atoms when considering the ^1H and ^{13}C NMR chemical shifts and examining the ^{13}C and DEPT NMR data which exhibited twelve carbon signals with two aliphatic methylenes, one aliphatic methine, four sp^2 hybridized methines, and five sp^2 hybridized quaternary carbons being categorized (Table 1). The above NMR data were compatible with a benzene moiety, a tetrasubstituted ethylene, and a carbonyl except for the above-mentioned three aliphatic carbons. The long range ^1H , ^{13}C -correlations from δ_{H} 7.53 to δ_{C} 104.9, 125.6, 136.7, and 122.9, from δ_{H} 7.43 to δ_{C} 122.9, 120.0, and 125.6, from δ_{H} 7.22 to δ_{C} 111.9, 136.7, and 118.2, from δ_{H} 7.13 to δ_{C} 118.2, 125.6, 122.9, and 111.9, from δ_{H} 4.38 and 4.54 to δ_{C} 125.4, 104.9, and 55.0, from δ_{H} 3.13 and 3.20 to δ_{C} 104.9, 55.0, 171.3, and 125.6/125.4, and from δ_{H} 4.27 to δ_{C} 171.3, 40.5, 21.7, and

104.9 established the constitutional formula of **1** as indicated by Figure 1. The complete assignment of the NMR data is listed in Table 1.

On evaluation of compound **1** for its cytotoxic effects on two human cancer cell lines, cell growth was measured using a sulforhodamine B (SRB) assay. Results of means of three replicates are expressed as the percentage of viability compared to negative control. Compound **1** exhibited moderate cytotoxicities against HCT-8 and HepG2 cell-lines in the evaluation, with viability of HepG2 and HCT-8 cells being 80.42% and 80.22% after treatment for 48 hours, and 76.14% and 71.48% after 72 hours, respectively, when using a concentration of 140 $\mu\text{g/mL}$. The viabilities at other time points were relatively lower.

3. Experimental

3.1. General Experimental Procedures. IR spectra were obtained on a Nicolet 5700 spectrometer. 1D and 2D NMR spectra were recorded on a Mercury-400 or a MERCURY-300 NMR spectrometer. Chemical shifts (δ) were given in ppm using tetramethylsilane (TMS) as internal standard (δ 0.00). ESI-MS and HRESI-MS were measured on an Agilent 1100 series LC-MSD-Trap-SL spectrometer. RP-18 (YMC-GEL, ODS-A, 12 nm, S-50 mm; YMC Co., Kyoto, Japan) were used for column chromatography. Solvents were of analytical grade and were purchased from Beijing Chemical Company (Beijing, China).

3.2. Plant Material. *Cichorium endivia* was purchased from Beijing Xinfadi agricultural products wholesale market on July 2009 and authenticated by Associate Professor Ma Lin

FIGURE 1: The structure of compound 1 and the key HMBC ($\rightarrow$: H $\rightarrow$ C) correlations.

(Institute of Material Medica, Chinese Academy of Medical Sciences and Peking Union Medical College). A voucher specimen was deposited in New Drug Safety Evaluation Center, Institute of Materia Medica, Peking Union Medical College & Chinese Academy of Medical Sciences, China.

3.3. Extraction and Isolation. The air-dried and pulverized *C. endivia* (5.8 kg) was extracted three times under reflux conditions with 95% EtOH (80 L, 2 h; 70 L, 1 h; 68 L, 1 h). The combined EtOH extracts were evaporated in vacuum to yield a dark-green residue (1233 g, crude EtOH extract), which was suspended in 80% aq. EtOH. The resulting suspension was extracted with petroleum ether (60–90°C, 2 L, 1.5 L, 1.5 L, and 1.5 L). Evaporation of the aq. layers in vacuum yielded also a dark-green residue (940 g, 80% EtOH extract). The residue was redissolved in water and subsequently partitioned with EtOAc in separatory funnel exhaustively. The rest of water soluble fraction was loaded on a column filled with Daion HP-20 and eluted with H₂O, 60% EtOH, and 95% EtOH, respectively. The eluates of 60% EtOH were evaporated in vacuum which yielded a black residue (40 g).

The 60% EtOH fraction was redissolved in solvent of *n*-BuOH and washed with aq. 5% NaHCO₃ then H₂O, respectively. Evaporation of *n*-BuOH under reduced pressure gave 5.5 g of brown-green residue, which was submitted to an ODS-A column eluted with MeOH-H₂O of decreasing polarity (40%–100%) to yield compound 1 (108 mg) as pale-yellow powder.

3.3.1. 3S-1,2,3,4-Tetrahydro-β-carboline-3-carboxylic Acid (1). Pale-yellow powder. IR (KBr) $\nu_{\max}$ (cm⁻¹): 3284, 3019, 2849, 1642, 1598, 1452, 1409, 1271, 1221, and 740; ¹H and ¹³C NMR spectroscopic data are listed in Table 1. ESI-MS (positive mode) *m/z*: 217.1 [M+H]⁺ and 239.1 [M+Na]⁺; HRESI-MS (positive mode) *m/z*: 217.0967 [M+H]⁺ (calcd for C₁₂H₁₂N₂O₂, 217.0972).

3.4. Cytotoxicity Assay. HCT-8 and HepG2 cells were cultivated in RPMI1640 medium containing 0.22% sodium bicarbonate, 10% fetal bovine serum (FBS), 100 U/mL penicillin, and 100 μg/mL streptomycin. The cells were incubated in 5% CO₂-air at 37°C. Compound 1 was dissolved in phosphate-buffered saline (PBS) at a concentration of 1.4 mg/mL and

was diluted to the required concentration with RPMI1640 medium immediately before use.

The cell viability was measured by using sulforhodamine B (SRB) assay. Briefly, the cells were seeded in 96-well plates (1 × 10⁴ cells/well) and routinely cultured for 24 h. Compound 1 was added to in-serial concentrations (from 14 μg/mL to 140 μg/mL), while PBS was added alone to control wells as a negative control, and incubation was continued for an additional 48 h. SRB (1 mg/mL) was added to each well after the plates were fixed using TCA (0.4% m/v). After 20 minutes of incubation, each well was washed by acid (1% v/v) three times. Then wells were added into Tris (100 mmol/L), respectively. The absorbance of each well was recorded on a microplate spectrophotometer at 515 nm.

Acknowledgments

The authors thank Associate Professor L. Ma for the sample authentication. The authors are also thankful to the State Key Laboratory of Bioactive Substance and Function of Natural Medicine and Drug Safety Evaluation Center, Institute of Materia Medica, for providing cells and equipment for research.

References

- [1] M. S. DuPont, Z. Mondin, G. Williamson, and K. R. Price, "Effect of variety, processing, and storage on the flavonoid glycoside content and composition of lettuce endive," *Journal of Agricultural and Food Chemistry*, vol. 48, no. 9, pp. 3957–3964, 2000.
- [2] R. Llorach, A. Martínez-Sánchez, F. A. Tomás-Barberán, M. I. Gil, and F. Ferreres, "Characterisation of polyphenols and antioxidant properties of five lettuce varieties and escarole," *Food Chemistry*, vol. 108, pp. 1028–1038, 2008.
- [3] C. J. Chen, A. J. Deng, C. Liu, R. Shi, H. L. Qin, and A. P. Wang, "Hepatoprotective activity of *Cichorium endivia* L. extract and its chemical constituents," *Molecules*, vol. 16, pp. 9049–9066, 2011.
- [4] J. P. Peng, Y. Q. Qiao, and X. S. Yao, "Nitrogen-containing compounds from *Allium macrostemon* Bunge. and *Allium chinese* G. Don," *Chinese Journal of Medicinal Chemistry*, vol. 5, p. 134, 1995.
- [5] S. Yahara, N. Uda, E. Yoshio, and E. Yae, "Steroidal alkaloid glycosides from tomato (*Lycopersicon esculentum*)," *Journal of Natural Products*, vol. 67, no. 3, pp. 500–502, 2004.