<http://www.e-journals.net>

ISSN: 0973-4945; CODEN ECJHAO

E-Journal of Chemistry

Vol. 1, No. 2, pp 132-136, April 2004

Kinetics of Oxidation of Some Essential Amino Acids by *N*-Chlorosaccharin in Aqueous Acetic Acid Medium

N. A. MOHAMED FAROOK*, G. A. SEYED DAMEEM,
A. MURUGESAN and M. KANAGARAJ

P. G & Research Department of Chemistry
Khadir Mohideen College,
Adirampattinam - 614 701 India
nafarook@hotmail.com

Received 15 March 2004; Accepted 30 March 2004

Abstract: The kinetics of oxidation of some essential amino acids namely, valine, leucine and phenylalanine, (AA) by *N*-chlorosaccharin (NCSA) in aqueous acetic acid medium in the presence of perchloric acid have been investigated. The observed rate of oxidation is first order in [AA], [NCSA] and of inverse fractional order in $[H^+]$. The main product of the oxidation is the corresponding aldehyde. The ionic strength on the reaction rate has no significant effect. The effect of changing the dielectric constant of the medium on the rate indicates the reaction to be of dipole-dipole type. Hypochlorous acid has been postulated as the reactive oxidizing species. The reaction constants involved in the mechanism are derived. The activation parameters are computed with respect to slow step of the mechanism.

Keywords: Oxidation, Kinetics, Amino acids, *N*-Chlorosaccharin

Introduction

The chemistry of reactions of *N*-halo compounds form a separate branch, which is a great synthetic importance.¹ *N*-Halo compounds have been extensively employed as oxidizing agents for organic substrates²⁻³. In the recent development, *N*-halo compounds are the source of positive halogen and have been exploited as oxidant for a variety of substrates in both acidic and alkaline media. The nature of active oxidizing species and mechanism

depends on the nature of the halogen atom, the groups attached to the nitrogen and the reaction conditions. Although a lot of works have been reported on the oxidation of organic compounds by *N*-halo compounds,⁴⁻⁷ it is to be noted that no systematic kinetic investigation on the oxidation of amino acids by *N*-chlorosaccharin has yet been reported in the literature. Here we report the results of the kinetics of the oxidation of valine, l eucine and phenylalanine (AA) with *N*-chlorosaccharin (NCSA) in aqueous acetic acid medium in the presence of perchloric acid.

Experimental

All the chemicals used were of A.R. grade. Double-distilled water was used throughout. *N*-chlorosaccharin⁸ were prepared using reported procedures. Standard solution of NCSA was prepared afresh. Perchloric acid (AnalaR) was used as source of hydrogen ions. Sodium perchlorate (Merck) was used to keep the ionic strength constant.

The reaction was carried out under pseudo-first order condition ($[AA] \gg [NCSA]$). The reaction was followed by potentiometrically by setting up a cell made up of the reaction mixture into which the platinum electrode and reference electrode (SCE) were dipped. The emf of the cell was measured periodically using an Equip-Tronic potentiometer, while the reaction mixture was continuously stirred. The pseudo-first order rate constants computed from the plots of $\log (E_t - E_\infty)$ against time were reproducible within $\pm 3\%$.

Results and Discussion

Stoichiometry and reaction products

Different sets of reaction mixtures containing different quantities of NCSA and AA at constant concentration of perchloric acid and sodium perchlorate were allowed to react for 24 h at 30°C and then analyzed. The remaining NCSA was estimated. The oxidation products were identified as aldehyde and saccharin by their spot tests; it was confirmed by noting the mixed melting point, chemical methods and TLC techniques. The results are in good agreement with 1:1 stoichiometry.

Reaction order

The kinetics of oxidation of amino acids by NCSA was investigated at several initial concentrations of the reactants in 50%-80% acetic acid-water mixture in the presence of perchloric acid. At constant $[H^+]$ with [substrate] in excess, the plot of $\log (E_t - E_\infty)$ versus time is linear, indicating a first order dependence of rate on [NCSA]. Plots of $\log k_{obs}$ versus $\log [AA]$ are linear with a slope of nearly unity for all the substrates (Figure 1) and a plot of k_{obs} versus $[AA]$ is also linear passing through the origin indicating a first order dependence on $[AA]$ (Figure 2). The pseudo-first order rate constants decrease with increase in $[H^+]$ in the range $0.5 - 1.6 \text{ mol dm}^{-3}$ (Table 1). The linearity of the plots of $\log [NCSA]$ versus time indicates the order in [NCSA] as unity.

The rate increases with increase in dielectric constant of the medium. A plot of $\log k_{\text{obs}}$ versus $[D-1/2D+1]$, where D is the dielectric constant of the medium, is linear with positive slope indicating a dipole- dipole type of interaction. The reaction have been studied in the temperature range 30-50°C and using Arrhenius equation the thermodynamic parameters have been calculated (Table 2). The oxidation of amino acids by NCSA under nitrogen atmosphere fails to induce polymerization of added acrylonitrile, thus ruling out radical oxidation. The retarding effect of added saccharin suggests that the pre-equilibrium step involves hydrolysis of NCSA to give saccharin and HOCl (eq. 2).

Of the four probable oxidizing species in the reaction, *viz.* NCSA, NCSAH^+ , HOCl and H_2OCl^+ , HOCl is considered as the active oxidizing species since the rate is an inverse function of [saccharin] and $[\text{H}^+]$.

Figure 1. Plots of $\log [\text{AA}]$ versus $\log k_1$ (A=Valine ; B=leucine ; C=Phenylalanine)

Table 1. Effect of varying [amino acid], [NCSA] and $[\text{H}^+]$ on the rate of reaction at 30°C^a.

$10^2 [\text{AA}]$ mol dm ⁻³	$10^3 [\text{NCSA}]$ mol dm ⁻³	$[\text{H}^+]$ mol dm ⁻³	$10^5 \times k_{\text{obs}} \text{ s}^{-1}$		
			Val	Leu	Phe
2.0	1.0	0.5	2.52	2.64	3.36
4.0	1.0	0.5	5.05	5.34	6.71
6.0	1.0	0.5	7.58	7.92	10.1
8.0	1.0	0.5	9.99	10.6	13.4
2.0	3.0	0.5	2.66	2.66	3.33
2.0	5.0	0.5	2.64	2.64	3.35
2.0	7.0	0.5	2.63	2.63	3.34
2.0	1.0	0.8	2.15	2.28	2.51
2.0	1.0	1.2	1.74	1.85	2.06
2.0	1.0	1.6	1.31	1.42	1.78

^aGeneral conditions: $[\text{NaClO}_4] = 0.5 \text{ mol dm}^{-3}$,
Solvent composition: 50% Acetic acid – 50% Water (v/v).

Figure 2. Plots of [AA] versus k_1 (A=Valine ; B=Leucine ; C=Phenylalanine)

Table 2. Activation parameters for the oxidation of amino acids by NCSA.^a

Parameters	Valine	Leucine	Phenylalanine
$10^5 k_{\text{obs}} (\text{s}^{-1})$	2.52	2.64	3.36
$E_a (\text{kJ mol}^{-1})$	45.7	47.6	49.2
$\Delta H^* (\text{kJ mol}^{-1})$	43.2	45.1	46.7
$\Delta S^* (\text{JK}^{-1} \text{mol}^{-1})$	40.2	41.2	43.2
$\Delta G^* (\text{kJ mol}^{-1})$	61.2	65.2	72.4

^aGeneral conditions: $[\text{NaClO}_4] = 0.5 \text{ mol dm}^{-3}$,

Solvent composition: 50% Acetic acid – 50% Water (v/v)

$[\text{AA}] = 0.02 \text{ mol dm}^{-3}$, $[\text{NCSA}] = 0.001 \text{ mol dm}^{-3}$.

The retarding effect of $[\text{H}^+]$ on the rate of reaction can be explained by considering the equilibrium (3) between the substrate (AA) and H^+ .

This effect coupled with the solvent effect does point out that the reaction with unprotonated amino acid should be more predominant.

Mechanism

The first order dependence each on [AA] and [NCSA] reveals that overall rate may involve the interaction of HOCl and amino acid in the rate determining step to give a mono ester similar to that reported by Komal Reddy and Sundaram for *N*-bromoacetamide oxidations^{9,10}. The mono ester involving *O*-Cl linkage may then undergo a series of fast steps as shown in Scheme 1 to give aldehyde and ammonia.

Scheme 1

The rate law based on **scheme 1** is represented by eq. (8)

$$\frac{-d[\text{NCSA}]}{dt} = \frac{k_1 k_3 [\text{NCSA}] [\text{SH}^+]}{k_2 [\text{H}^+] [\text{C}_6\text{H}_4\text{SO}_2\text{CONH}]} \quad (8)$$

Equation (8) clearly points out the observed results, *i.e.* first order in [NCSA], [AA] and inverse first order in [H⁺] negligible influence of changes in ionic strength and the effect of dielectric constant of the medium on the rate of oxidation.

Acknowledgment

The author N.A.Mohamed Farook is thankful to UGC for financial support to complete this project

References

1. (a) Patrocino F A *J Organomet. Chem.* 2000 **603** 220 (b) Dhurn P S, Mohe N U and Salunkhe M M *Synth. Commun.* 2001 **31** 3653 (c) Caribano V, Rodriguez J F, Santose M, Sanz-Tejedor M A, Carreno M C, Gonzalez G and Garcia-Ruano J L *Synthesis* 2001 **14** 2175 (d) Bandgar B P, Uppalla L S and Sadavarte V S *Syn. Lett.* 2001 **11** 1715
2. Thenraja D, Subramaniam P and Srinivasan C *J. Chem. Soc. Perkin Trans* 2002 **2** 2125
3. Mukaiyama T, Mastsuo J I, Lida D and Kitagawa H *Chem. Lett.* 2001 **8** 846
4. Hambly G F and Chan T H *Tetrahedron Lett.* 1986 **27** 2563
5. Antelo J M, Arce F, Crugeiras J O and Parajo M J. *Phys. Org. Chem.* 1997 **10** 631
6. Karunakaran C and Ganapathy K *Indian J. Chem.* 1990 **29A** 133
7. Harihar A L, Kembhavi M R and Nandibewoor S T *J. Indian Chem. Soc.* 1999 **76** 128
8. Chattaway F D *J. Chem. Soc.* 1905 **87** 1884
9. Komal Reddy M and Sundaram E V *Indian J. Chem.* 1986 **25A** 471
10. Gopalakrishnan G and Hogg J L *J. Org. Chem.* 1985 **50** 1206