http://www.e-journals.net

ISSN: 0973-4945; CODEN ECJHAO

E-Journal of Chemistry

Vol. 3, No.3, pp 159-163, July 2006

Infrared Irradiation Synthesis of Substituted 5-Oxo-1, 2, 3, 4, 5, 6, 7, 8-octahydroquinoline Derivatives under Solvent-free Conditions

SHU-XIANG WANG¹, SHAO-BO GUO¹, MING-ZHU GAO²,
JI-TAI LI¹ and YUN-FEI DUAN¹

¹College of Chemistry and Environmental Science,
Hebei University, Baoding 071002, Hebei Province P. R. China.

²Luck Film Corporation, P.R. China.

Received 18 May 2006; Accepted 23 June 2006.

Abstract: Infrared irradiation promoted the synthesis of substituted 5-oxo-1, 2, 3, 4, 5, 6, 7, 8-octahydroquinoline derivatives. The condensations of 5, 5-dimethyl-1, 3-cyclohexane-dione, 1, 3-diaryl-2-propen-1-one and ammonium acetate in the absence of solvent under infrared irradiation for 3 h gave the title compounds in 67.6%~86.5% yields

Keywords: substituted 5-oxo-1, 2, 3, 4, 5, 6, 7, 8-octahydroquinoline derivatives; synthesis; solvent-free; infrared irradiation.

Introduction

1, 4-Dihydropyridines derivatives (1, 4-DHPs) possess a variety of biological activities. For example, they have been used as effective calcium channel modulators for the treatment of cardiovascular disorders¹. 4-Substituted 1, 4-dihydropyridines are analogues of NADH coenzymes and can be used as medicine². Much effort has been devoted to the synthesis of 1, 4-DHPs with different substituents³ or heteroatom⁴. The substituted 5-oxo-1, 2, 3, 4, 5, 6, 7, 8-octahydroquinoline derivatives are these chemical modifications rings. Therefore, preparation of it has attracted considerable attention in recent years.

Synthetic chemistry requires the consumption of energy and material. One of the greatest environmentally problematic aspects of organic synthetic chemistry is the use of solvent. The solvent has to be produced in another chemical process, which is involved in

consuming energy and resources. Within the reaction the vast amount of solvent has to be heated or cooled, consuming large amount of energy, and has to be removed after completion of the reaction from the reaction mixture. A move away from the use of solvents in organic synthesis has led in some cases to improved results and more benign synthetic procedures⁵.

Infrared irradiation is a secure, cheap energy and can be easily obtained. Therefore, infrared irradiation in organic synthesis has been the focus of considerable attention in recent years. Guillermo⁶ reported a synthesis of 2-alkylbenzimidazoles and benzimidazol-2-one by using a natural clay and infrared irradiation in solvent-free conditions. Alcerreca⁷ reported that several benzaldehydes were condensed with barbituric acid in absence of solvent under infrared irradiation, affording the corresponding 5-benzylidene barbituric acids. Obrador⁸ also synthesized benzylidenemalonitriles, benzylidenecyanoacetamides and benzylidenecyanoacetic acids under infrared irradiation.

All of the results stated above spur us to study the possibility of synthesis of substituted 5-oxo-1, 2, 3, 4, 5, 6, 7, 8-octahydroquinoline derivatives (3) in the absence of solvent under infrared irradiation. Herein, we wish to report the results of the condensations of 5, 5-dimethyl-1, 3-cyclohexane-dione, 1, 3-diaryl-2-propen- 1-one and ammonium acetate in solvent-free under infrared irradiation (Scheme 1).

Scheme 1

Experimental

Materials and Measurements

Melting points are uncorrected. 5, 5-Dimethyl-1, 3-cyclohexane-dione was purchased from Fluka and was used without further purification. IR spectra were recorded on Bio-Rad FTS-40 spectrometer (KBr). ¹HNMR spectra were measured on Bruker AVANCE 400(400 MHz) spectrometer in CDCl₃ using TMS as internal standard. The reaction was performed under W-10 IR lamp (with a nominal power of 500 W; Beijing Zhen-xing lab instrument Co., Ltd.).

General Procedure

A dry 5 mL round-bottomed flask was charged with 1 (1 mmol), 2 (1.25 mmol) and ammonium acetate (2.5mmol). The mixture was irradiated at a distance of 10 cm from an infrared lamp for 3 h, and then cooled to room temperature. The solid was transferred with a small amount ethanol, and then filtered. The obtained solid was washed with water (3×10 mL). The crude product was purified by recrystallization from 95% ethanol to give 3. The authenticity of the products was established by comparing their melting points, IR, ¹HNMR and elemental analysis.

3d: $\nu_{\text{max}}/\text{cm}^{-1}$ 3240, 3020, 2980, 1610, 1580, 1500, 830, 760, 690; δ_{H} 1.13(3H, s, CH₃), 1.24(3H, s, CH₃), 2.28(1H, d, $J=16$ Hz, CH), 2.35(1H, d, $J=16$ Hz, CH), 2.40(1H, d, $J=16$ Hz, CH), 2.51(1H, d, $J=16$ Hz, CH), 4.83(1H, d, $J=5.2$ Hz, CH), 5.37(1H, dd, $J=5.2$ Hz, $J'=1.2$ Hz, CH), 5.84(1H, s, NH), 7.23-7.53(9H, m, ArH). Anal. Calcd for C₂₃H₂₂ClNO: C, 75.91; H, 6.10; N, 3.85. Found(%): C, 76.04; H, 5.94; N, 3.68.

3e: $\nu_{\max}/\text{cm}^{-1}$ 3250, 3020, 2990, 2980, 1620, 1590, 1500, 810, 690; δ_{H} 1.13(3H, s, CH₃), 1.14(3H, s, CH₃), 2.21 (1H, d, $J=16$ Hz, CH), 2.27(1H, d, $J=16$ Hz, CH), 2.32(1H, d, $J=16$ Hz, CH), 2.41(1H, d, $J=16$ Hz, CH), 4.67 (1H, d, $J=5.2$ Hz, CH), 5.26(1H, dd, $J=5.2$ Hz, $J'=1.2$ Hz, CH), 5.83(1H, s, NH), 5.91(2H, s, OCH₂O), 6.72-6.86(3H, m, ArH), 7.29-7.45(4H, m, ArH). Anal. Calcd for C₂₄H₂₂ClNO₃: C, 70.67; H, 5.44; N, 3.43. Found(%): C, 70.86, H, 5.23; N, 3.15.

3f: $\nu_{\max}/\text{cm}^{-1}$ 3240, 3060, 2950, 1660, 1590, 1490, 820, 760, 690; δ_{H} 1.06(3H, s, CH₃), 1.13(3H, s, CH₃), 2.24(1H, d, $J=12$ Hz, CH), 2.31(1H, d, $J=12$ Hz, CH), 2.34(1H, d, $J=16$ Hz, CH), 2.42(1H, d, $J=16$ Hz, CH), 4.75 (1H, d, $J=5.2$ Hz, CH), 5.27(1H, dd, $J=5.2$ Hz, $J'=1.2$ Hz, CH), 5.97(1H, s, NH), 7.14-7.39(4H, m, ArH), 7.41-7.44(5H, m, ArH). Anal. Calcd for C₂₃H₂₂ClNO: C, 75.79; H, 6.13; N, 3.99. Found(%): C, 75.92, H, 6.09; N, 3.85.

3g: $\nu_{\max}/\text{cm}^{-1}$ 3220, 3060, 2960, 1660, 1580, 1490, 820, 770, 690; δ_{H} 1.14(6H, s, CH₃), 2.23(1H, d, $J=12$ Hz, CH), 2.25(1H, d, $J=12$ Hz, CH), 2.38(1H, d, $J=16$ Hz, CH), 2.43(1H, d, $J=16$ Hz, CH), 5.16(1H, d, $J=5.2$ Hz, CH), 5.25(1H, dd, $J=5.2$ Hz, $J'=1.2$ Hz, CH), 5.96(1H, s, NH), 7.17-7.35(3H, m, ArH), 7.37(5H, m, ArH). Anal. Calcd for C₂₃H₂₁Cl₂NO: C, 69.17; H, 5.41; N, 3.71. Found(%): C, 69.35, H, 5.31; N, 3.52.

3h: $\nu_{\max}/\text{cm}^{-1}$ 3320, 3040, 2950, 2870, 1660, 1580, 1490, 810, 770; δ_{H} 1.05(3H, s, CH₃), 1.14(3H, s, CH₃), 2.22(1H, d, $J=12$ Hz, CH), 2.28(1H, d, $J=12$ Hz, CH), 2.30(1H, d, $J=16$ Hz, CH), 2.41(1H, d, $J=16$ Hz, CH), 4.70(1H, d, $J=4.8$ Hz, CH), 5.28(1H, dd, $J=4.8$ Hz, $J'=1.2$ Hz, CH), 5.80(1H, s, NH), 7.09-7.26(4H, m, ArH), 7.28-7.35(4H, m, ArH). Anal. Calcd for C₂₄H₂₄ClNO: C, 76.03; H, 6.30; N, 4.00. Found(%): C, 76.28, H, 6.40; N, 3.71

3i: $\nu_{\max}/\text{cm}^{-1}$ 3230, 3050, 2950, 2870, 1660, 1580, 1490, 810, 760, 695; δ_{H} 1.04(3H, s, CH₃), 1.12(3H, s, CH₃), 2.18 (1H, d, $J=16$ Hz, CH), 2.25(1H, d, $J=16$ Hz, CH), 2.31(1H, d, $J=16$ Hz, CH), 2.41(1H, d, $J=16$ Hz, CH), 4.72 (1H, d, $J=5.2$ Hz, CH), 5.31(1H, dd, $J=5.2$ Hz, $J'=1.2$ Hz, CH), 5.95(1H, s, NH), 7.08-7.28(4H, m, ArH), 7.36-7.44(5H, m, ArH). Anal. Calcd for C₂₄H₂₅NO: C, 83.82; H, 7.41; N, 4.14. Found(%): C, 83.93; H, 7.34; N, 4.08.

Results and Discussion

The effects of the reaction conditions on yields of compounds **3** are summarized in Table 1. Firstly, the mixture of 5, 5-dimethyl-1, 3-cyclohexane-dione (**2**), 1, 3-phenol -2-propen-1-one (**1a**) and ammonium acetate was chosen as the model reaction to detect the proper distance from IR lamp for this reaction. When the distance is 5 cm, 10 cm and 20 cm, the yields of **3a** are 40.4%, 83.0% and 76.5% respectively. At a distance of 10cm, reaction of **1a**, **2** and ammonium acetate for 2 h, 3 h, 4 h and 5 h, afford **3a** in 53.7%, 83.0%, 82.4% and 83.1% yield respectively.

From the results above, the reaction conditions we chose are: the reaction time is 3 h and the distance is 10 cm. Using the present method we did a series of experiments for synthesis of substituted 5-oxo-1, 2, 3, 4, 5, 6, 7, 8-octahydroquinoline derivatives (**3a-3i**), the results were summarized in Table 2.

Wang⁹ reported the synthesis of **3a** in DMF. However, we stirred **2**, **1a** and ammonium acetate in the same condition which resulted in a poor yield of **3a**. We prolonged the reaction time from 2 h to 5 h; the yield of **3a** was only increased to 28.6% (Table 2, **3a**⁴). Using this procedure, the yield of **3a** was lower than that of reported.

In addition, when the distance is 10 cm from IR lamp we measured the reaction temperature, which was 80 °C. To compare the effectiveness of infrared irradiation method with stirring

method, we stirred **1**, **2** and ammonium acetate in the absence of solvent at 80 °C for 3 h. The yield of **3a** was 73.8% (Table 2, **3a^B**), and lower than that of infrared irradiation method.

Table 1. The Effects of the Reaction Conditions on the Yield of Substituted 5-Oxo-1, 2, 3, 4, 5, 6, 7, 8-Octahydroquinoline Derivatives

Entry	Distance from IR lamp/cm	Time/h	Yield*/%
1	5	3	40.4
2	10	3	83.0
3	20	3	76.5
4	10	2	53.7
5	10	3	83.0
6	10	4	82.4
7	10	5	83.1

*Yields of pure isolated product

^aThe reaction temperature of different distance from IR lamp

Table 2 The Substituted 5-Oxo-1,2,3,4,5,6,7,8-Octahydroquinoline Derivatives Synthesized by 5,5-Dimethyl-1,3-Cyclohexane-Dione, Ammonium Acetate and 1,3-Diaryl-2-Propen-1-One in Absence of Solvent Under Infrared Irradiation

Entry	Ar	Ar'	Yield*/%	M.P./°C [Ref ⁹]
3a	C ₆ H ₅	C ₆ H ₅	83.0	206-208[206-208]
3a^A	C ₆ H ₅	C ₆ H ₅	28.6	206-208
3a^B	C ₆ H ₅	C ₆ H ₅	73.8	206-208
3b	4-ClC ₆ H ₄	C ₆ H ₅	85.4	224-226[224-226]
3c	3,4-(CH ₃ O) ₂ C ₆ H ₃	C ₆ H ₅	76.9	236-238[236-238]
3d	C ₆ H ₅	4-ClC ₆ H ₄	78.5	212-214[250-252]
3e	3,4-(CH ₃ O) ₂ C ₆ H ₃	4-ClC ₆ H ₄	86.5	250-252[176-177]
3f	3-ClC ₆ H ₄	C ₆ H ₅	79.6	225-227
3g	2,4-ClC ₆ H ₄	C ₆ H ₅	70.4	204-206
3h	4-CH ₃ C ₆ H ₄	4-ClC ₆ H ₄	67.6	234-236
3i	4-CH ₃ C ₆ H ₄	C ₆ H ₅	75.1	213-215

* Yields of pure isolated product.

^A: Stirring for 5 h at 80 °C using DMF as solvent.

^B: Stirring for 3 h at 80 °C in the absence of solvent.

Varying the substituents on the benzene ring of 1, 3-diaryl-2-propen-1-one did not detrimentally affect the yields of compound **3**. 1, 3-Diaryl-2-propen-1-one containing electron-withdrawing groups and electron-donating groups on the benzene were reacted well to give the corresponding 5-oxo-1, 2, 3, 4, 5, 6, 7, 8-octahydroquinoline derivatives (**3b-3i**) in good yields.

Conclusions

We have found a clean and easy method for the preparation of the substituted 5-oxo-1, 2, 3, 4, 5, 6, 7, 8-octahydroquinoline derivatives. This method provides many obvious advantages such as the avoidance of discharging harmful organic solvent, the simplicity of the work-up, good yields, being environmentally friendly and using the safe and easily obtained IR lamp. The results showed that infrared irradiation can be used as a valuable means for synthesis of organic compounds.

References

- 1 Suárez M, Verdecia Y, Illescas B, Martínez-Alvarez R, Alvarez A, Ochoa E, Seoane C, Kayali N and Martín N, *Tetrahedron.*, 2003, **59**(46), 9179
- 2 Mauzerall D and Westheimer F H, *J. Am. Chem. Soc.*, 1955, **77**(8), 2261.
- 3 Stout D M and Meyers A I, *Chem. Rev.*, 1982, **82**(2), 223.
- 4 (a) Chorvat R J and Rorig K J, *J. Org. Chem.*, 1988, **53**(24), 5779. (b) Guzman A, Romero M Maddox M L and Muchowski J M, *J. Org. Chem.*, 1990, **55**(22), 5793. (c) Kappe C O, *Tetrahedron.*, 1993, **49**(32), 6937.
- 5 (a) Kidwai W, Sapra P, Bhushan K R and Misra P, *Synthesis*, 2001, **10**, 1509. (b) Kidwai W and Sapra P, *Synth. Commu.*, 2002, **32**(11), 1639.
- 6 Guillermo Penieres C, Imelda Bonifas A, José Guadalupe López C, José Guadalupe Garcia E and Cecilio Alvarez T, *Synth. Commu.*, 2000, **30**(12), 2191.
- 7 Alcerrca G, Sanabria R, Miranda R, Arroyo G, Tamariz J and Delgado F, *Synth. Commu.*, 2000, **30**(7), 1295.
- 8 Obrador E, Castro M, Tamariz J, Zepeda G, Miranda R and Delgado F, *Synth. Commu.*, 1998, **28**(24), 4649.
- 9 Xiang-Shan Wang , Da-Qing Shi and Shu-Jiang Tu, *Synth. Commu.*, 2002, **32**, 3449.