Synthesis and Antimicrobial Activity of Newer Quinazolinones

J. A. PATEL, B. D. MISTRY* and K. R. DESAI**

*Department of Chemistry, B. K. M. Science College, Valsad-396001,
Affiliated with

**Department of Chemistry,
Veer Narmad South Gujarat
University, Surat-395007, India.

Received 2 January 2006; Accepted 14 April 2006

Abstract: 2-alkyl-6-bromo-3,1-benzoxazine-4-one (2) is synthesized by treating p-Bromoanthranilic acid and Acetylchloride or Benzoylchloride. Reaction of 2-alkyl-6-bromo-3,1-benzoxazine-4-one (2). with hydrazinehydrate furnish the corresponding 3-Amino-2-methyl-6-bromoquinazoline-4(3H)-one (3) which on reaction with benzaldehyde afford N,N - arylidene derivative (4). Reaction of 4 with various diazonium salts yields 6-bromo-2-alkyl/aryl-3{[phenyl(phenyldiazenyl)methylene]amino}quinazolin-4(3H)-one .

Key words: Synthesis, quinazolidinone, antibacterial activity

Introduction

Quinazoline nucleus possessed the potent pharmacodynamic nucleus. In addition several quinazoline derivatives posses diverse biological activities Viz. anticonvulsant¹, hyponotics², antiinflammatory^{3,4}, diuretics antihypertensive^{5,6,7}, antitubercular etc. quinazoline system posses variable sites at 2 and 3 positions substituted by various hetrocyclic moieties afford the potent anticonvulsant agent. Furthermore SAR study of different quinazolinone derivatives have revealed that the substitution at 3-position has significant role on hypotensive activity.

Experimental

All the melting points were determined in open capillary and are uncorrected. The purity is checked by TLC. IR Spectras were recorded in KBr on shimatzu F.T. - IR 8300 spectrophotometer. Analytical data were also confirmed from its H - NMR Spectra.

The starting compound 2-alkyl/aryl-3-amino-6-bromo quinazolinone has been prepared according to known method^{8,9}

3-amino-2-alkyl/aryl-6-bromoquinazolin-4(3H)-one **3** has been prepared by known method.^{8,9}

Synthesis of 6-bromo-2-alkyl/aryl-3-[(1E)-phenylmethylene]amino]quinazolin-4(3H)-one **4:**

3-Amino-2-alkyl/aryl-7-bromoquinazolin-4(3H)-one **3** (0.01 mole) and benzaldehyde (0.01 mole) were refluxed in absolute ethanol (40ml) for 3 hrs. The excess solvent was then distilled off and the resulting solid washed with water, dried and recrystallized from ethanol.

*Synthesis of 6-bromo-2-alkyl/aryl-3-[[phenyl (phenyldiazenyl) methylene] amino] quinazoline-4(3H)-one **5** :*

Appropriate amine (0.02 Mole) in glacial Acetic acid (2ml) & HCl (1.5ml) was diazotised with NaNO₂ (0.2 g in 2 ml H₂O) at 0–5 °C. The resultant diazoniumchloride solution was added with stirring to a solution of (**4**) (0.02 mole) in pyridine. (3 ml) in the cold. The reaction mixture. was left over night at ambient temperature. There after, it was poured into cold water (250 ml) with stirring when a dark coloured solid separated out. It was filtered, washed, repeatedly with water containing HCl and finally with water and recrystallized from methanol to furnish 5a₁–a₁₆. (Table 1)

Table 1.Elemental analysis of synthesized compounds **5a₁–a₁₆**

No.	R ₁	R ₂	M.F.	Yield %	M.P. (°C)	%Analysis Calc.(Found)		
						C	H	N
5a ₁	CH ₃	-H	C ₂₇ H ₁₈ N ₅ OBr	76	160°C	63.79 (63.20)	3.57 (4.28)	13.78 (12.40)
5a ₂	CH ₃	2-OCH ₃	C ₂₈ H ₂₀ N ₅ O ₂ Br	55	110– 112°C	62.46 (62.75)	3.74 (3.94)	13.01 (13.50)
5a ₃	CH ₃	3-OCH ₃	C ₂₈ H ₂₀ N ₅ O ₂ Br	58	113– 115°C	62.46 (62.75)	3.74 (3.94)	13.01 (13.50)
5a ₄	CH ₃	4-OCH ₃	C ₂₈ H ₂₀ N ₅ O ₂ Br	52	120°C	62.46 (62.75)	3.74 (3.94)	13.01 (13.50)
5a ₅	CH ₃	2-Br	C ₂₇ H ₁₇ N ₅ OBr ₂	56	145°C	55.22 (55.50)	2.92 (3.02)	11.93 (12.05)
5a ₆	CH ₃	3-Br	C ₂₇ H ₁₇ N ₅ OBr ₂	62	130– 135°C	55.22 (55.50)	2.92 (3.02)	11.93 (12.05)
5a ₇	CH ₃	4-Br	C ₂₇ H ₁₇ N ₅ OBr ₂	80	155°C	55.22 (55.50)	2.92 (3.02)	11.93 (12.05)
5a ₈	CH ₃	2-Cl	C ₂₇ H ₁₇ N ₅ OBr Cl	66	136– 140°C	59.74 (60.03)	3.16 (3.33)	12.90 (13.15)
5a ₉	CH ₃	3-Cl	C ₂₇ H ₁₇ N ₅ OBr Cl	68	148°C	59.74 (60.03)	3.16 (3.33)	12.90 (13.15)
5a ₁₀	CH ₃	4-Cl	C ₂₇ H ₁₇ N ₅ OBr Cl	56	142°C	59.74 (60.03)	3.16 (3.33)	12.90 (13.15)
5a ₁₁	CH ₃	2-NO ₂	C ₂₇ H ₁₇ N ₆ O ₃ Br	68	162– 165°C	58.60 (58.61)	3.10 (3.28)	15.19 (15.75)
5a ₁₂	CH ₃	3-NO ₂	C ₂₇ H ₁₇ N ₆ O ₃ Br	63	168°C	58.60 (58.61)	3.10 (3.28)	15.19 (15.75)
5a ₁₃	CH ₃	4-NO ₂	C ₂₇ H ₁₇ N ₆ O ₃ Br	67	156°C	58.60 (58.61)	3.10 (3.28)	15.19 (15.75)
5a ₁₄	CH ₃	2-CH ₃	C ₂₈ H ₂₀ N ₅ OBr	52	128°C (d)	64.38 (64.81)	3.86 (4.07)	13.41 (13.71)
5a ₁₅	CH ₃	3-CH ₃	C ₂₈ H ₂₀ N ₅ OBr	54	140(d)	64.38 (64.81)	3.86 (4.07)	13.41 (13.71)
5a ₁₆	CH ₃	4-CH ₃	C ₂₈ H ₂₀ N ₅ OBr	55	156(d)	64.38 (64.81)	3.86 (4.07)	13.41 (13.71)

Table –2. Elemental analysis of synthesized compounds **5b₁-b₁₆**

No.	R ₁	R ₂	M.F.	Yield %	M.P. (°C)	%Analysis Calc.(Found)		
						C	H	N
5b ₁	C ₆ H ₅	-H	C ₂₂ H ₁₆ N ₅ OBr	68	179–180°C	59.21 (59.20)	3.61 (3.59)	15.69 (15.40)
5b ₂	C ₆ H ₅	2-OCH ₃	C ₂₃ H ₁₈ N ₅ O ₂ Br	66	120°C	58.00 (58.25)	3.81 (3.94)	14.70 (14.88)
5b ₃	C ₆ H ₅	3-OCH ₃	C ₂₃ H ₁₈ N ₅ O ₂ Br	65	124°C	58.00 (58.25)	3.81 (3.94)	14.70 (14.88)
5b ₄	C ₆ H ₅	4-OCH ₃	C ₂₃ H ₁₈ N ₅ O ₂ Br	60	130–132°C	58.00 (58.25)	3.81 (3.94)	14.70 (14.88)
5b ₅	C ₆ H ₅	2-Br	C ₂₂ H ₁₅ N ₅ OBr ₂	78	125–127°C	50.31 (50.12)	2.88 (2.75)	13.33 (13.55)
5b ₆	C ₆ H ₅	3-Br	C ₂₂ H ₁₅ N ₅ OBr ₂	73	148°C	50.31 (50.12)	2.88 (2.75)	13.33 (13.55)
5b ₇	C ₆ H ₅	4-Br	C ₂₂ H ₁₅ N ₅ OBr ₂	74	136–137°C	50.31 (50.12)	2.88 (2.75)	13.33 (13.55)
5b ₈	C ₆ H ₅	2-Cl	C ₂₂ H ₁₅ N ₅ OBrCl	56	145°C	54.96 (54.87)	3.14 (3.33)	14.57 (14.76)
5b ₉	C ₆ H ₅	3-Cl	C ₂₂ H ₁₅ N ₅ OBrCl	52	140–141°C	54.96 (54.87)	3.14 (3.33)	14.57 (14.76)
5b ₁₀	C ₆ H ₅	4-Cl	C ₂₂ H ₁₅ N ₅ OBrCl	58	151°C	54.96 (54.87)	3.14 (3.33)	14.57 (14.76)
5b ₁₁	C ₆ H ₅	2-NO ₂	C ₂₂ H ₁₅ N ₆ O ₃ Br	82	170–172°C	53.78 (53.91)	3.08 (3.28)	17.11 (17.35)
5b ₁₂	C ₆ H ₅	3-NO ₂	C ₂₂ H ₁₅ N ₆ O ₃ Br	76	165°C	53.78 (53.91)	3.08 (3.28)	17.11 (17.35)
5b ₁₃	C ₆ H ₅	4-NO ₂	C ₂₂ H ₁₅ N ₆ O ₃ Br	78	162°C	53.78 (53.91)	3.08 (3.28)	17.11 (17.35)
5b ₁₄	C ₆ H ₅	2-CH ₃	C ₂₃ H ₁₈ N ₅ OBr	62	132–134°C (d)	60.01 (60.23)	3.94 (4.07)	15.21 (15.43)
5b ₁₅	C ₆ H ₅	3-CH ₃	C ₂₃ H ₁₈ N ₅ OBr	56	138°C (d)	60.01 (60.23)	3.94 (4.07)	15.21 (15.43)
5b ₁₆	C ₆ H ₅	4-CH ₃	C ₂₃ H ₁₈ N ₅ OBr	58	142°C (d)	60.01 (60.23)	3.94 (4.07)	15.21 (15.43)

Results and Discussion

Synthesized compounds have been confirmed by elemental analysis, IR Spectra and ¹HNMR Spectra.

Compound 5a₁ shows IR absorption bands at 1716–1670 cm⁻¹ (C=O stretching), 1676–1670 cm⁻¹ (C=N), 1340 cm⁻¹ (Ar-CH₃), 650 cm⁻¹ (C-Br stretching).

Compound 5b₁ shows IR absorption bands at 1734–1684 cm⁻¹ (C=O stretching) 1684 – 1654 cm⁻¹ (C=N), 652 cm⁻¹ (C-Br stretching).

^1H NMR of compound 5a₁: δ 2.45 (s, 3H, Ar-CH₃), 8.66 (d, 1H, C₅-H), 8.25 (d, 1H, C₇-H), 8.12 (d, 1H, C₈-H), 7.45–7.98 (m, 9H, Ar-H).

^1H NMR of compound 5b₁: δ 8.65 (d, 1H, C₅-H), 8.3 (d, 1H, C₇-H), 8.265 (d, 1H, C₈-H), 7.4–8.2 (m, 14H, Ar-H).

Antibacterial Activity

The synthesized compounds were tested against gram positive bacteria staphylococcus aureus and Bacillus cereus and gram negative bacteria E. coli and Pseudomonas aeruginosa and a yeast candida albicans.

Table 3. Antimicrobial Activity of Newly Synthesised Compounds, Zone of Inhibition (Mm)

Sample No.	E.coli	Pseudomonas aeruginosa	S.aureus	Bacillus cereus	Candida	M.I.C. for mycobacterium tuberculosis $\mu\text{g/ml}$
5a ₁	8	9	8	9	00	>100
5a ₂	6	6	6	6	00	>100
5a ₃	8	9	6	6	00	>100
5a ₄	9	9	6	6	00	>100
5a ₅	10	8	11	6	00	>100
5a ₆	8	7	8	8	00	>100
5a ₇	7	7	8	8	00	>100
5a ₈	6	6	6	6	00	>100
5a ₉	6	6	6	6	00	>100
5a ₁₀	8	8	8	8	00	>100
5a ₁₁	6	9	8	8	00	>100
5a ₁₂	7	8	8	8	00	>100
5a ₁₃	9	8	8	7	00	>100
5a ₁₄	9	8	6	6	00	>100
5a ₁₅	8	9	6	6	00	>100
5a ₁₆	8	8	6	6	00	>100
5b ₁	9	8	9	9	00	>100
5b ₂	7	9	7	7	00	>100
5b ₃	8	8	6	6	00	>100
5b ₄	11	12	9	9	00	>100
5b ₅	8	7	8	8	00	>100
5b ₆	9	8	8	8	00	>100
5b ₇	8	8	6	6	00	>100
5b ₈	9	9	8	8	00	>100
5b ₉	8	8	9	9	00	>100
5b ₁₀	11	12	10	10	00	>100
5b ₁₁	8	8	9	9	00	>100
5b ₁₂	7	7	7	7	00	>100
5b ₁₃	9	8	8	9	00	>100
5b ₁₄	6	6	6	6	00	>100
5b ₁₅	6	6	6	7	00	>100
5a ₁₆	6	6	6	8	00	>100

The stock solutions of compounds were prepared at a concentration of 5mg/ml & from stock solution the disc were prepared at a concentration of 100µg/ml. The testing was done on muller hinton agar plates by swabbing the agar plates with respective cultures, and placing the disc on it and incubating at 37°C for 24 hrs. the above results were obtained.

The compounds were also tested against mycobacterium tuberculosis at a concentration of 100µg/ml.

The L.J. medium was prepared from dehydrated powder following the instructions given on the dehydrated medium bottle. The compound were then added into the medium in the desired concentration and then inspissated then the prepared L.J. medium was inoculated with the pure culture of mycobacterium tuberculosis and incubated at 37°C for 7 weeks and then the results were read.

Standard drug tetracycline and Greseofulvin were also screened under similar conditions for comparison. Results are shown in Table 3.

Acknowledgement

We are grateful to the Department of Chemistry, B. K. M. Science College, Valsad. (Veer Narmad South Gujarat University, Surat) for providing the necessary facilities for the research work. Thanks to I.I.T. Bombay, for providing N.M.R. Spectras.

References

- 1 Archana, Srivastva V K, Chandra Ramesh and Ashokkumar, *Indian J. Chem*, 2002, **41B**, 2371.
- 2 Buyuktimkin S, *Chem Abstr*, 1987, **107**, 51388t.
- 3 Ka'si.R, Pande K, Bhalla T N, Barthwal J P, Gupta G P and Parmar S S, *J. Pharma Sci.*, 1990, **79(4)**, 317.
- 4 Atulkumar, Singh S, Saxena A K and Shanker K, *Indian J. Chem*, 1988, **27B**, 443.
- 5 Ashokkumar, Tyagi Mirdula and Srivastva V K, *Indian J. Chem*, 2003, **42B**, 2142.
- 6 Pandey V K, Tulsi Sarah and Tulsi Zehra, *Indian J. Chem*, 2004, **3B**, 180.
- 7 Tyagi Mirdula, Srivastva V K, Chandra Ramesh and Ashokkumar, *Indian J. Chem*, 2002, **41B**, 2367.
- 8 Wasfy A A F, *Indian J. Chem*, 2003, **42B**, 3102.
- 9 Yassin F A, Eissa A M F and Wasfy A A F, *Indian J. Chem*, 1994, **33B**, 1193.
- 10 Dalal B Guirguis, *Indian J. Chem*, 2000, **39B**, 264.